



National Comprehensive
Cancer Network®

NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®)

Non-Small Cell Lung Cancer

Version 3.2024 — March 12, 2024

NCCN.org

NCCN Guidelines for Patients® available at www.nccn.org/patients

Continue

National
Comprehensive
Cancer
Network®

NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

[NCCN Guidelines Index](#)
[Table of Contents](#)
[Discussion](#)

Gregory J. Riely, MD, PhD/Chair † ‡
Memorial Sloan Kettering Cancer Center

Douglas E. Wood, MD/Vice Chair ¶
Fred Hutchinson Cancer Center

David S. Ettinger, MD/Immediate Past Chair †
The Sidney Kimmel Comprehensive
Cancer Center at Johns Hopkins

Dara L. Aisner, MD, PhD ≠
University of Colorado Cancer Center

Wallace Akerley, MD †
Huntsman Cancer Institute at the University of Utah

Jessica R. Bauman, MD ‡ †
Fox Chase Cancer Center

Ankit Bharat, MD ¶ £
Robert H. Lurie Comprehensive Cancer
Center of Northwestern University

Debora S. Bruno, MD, MS †
Case Comprehensive Cancer Center/
University Hospitals Seidman Cancer Center
and Cleveland Clinic Taussig Cancer Institute

Joe Y. Chang, MD, PhD §
The University of Texas
MD Anderson Cancer Center

Lucian R. Chirieac, MD ≠ £
Dana-Farber/Brigham and Women's
Cancer Center

Malcolm DeCamp, MD ¶
University of Wisconsin Carbone Cancer Center

Aakash P. Desai, MD
O'Neal Comprehensive Cancer Center at UAB

Thomas J. Dilling, MD, MS §
Moffitt Cancer Center

Jonathan Dowell, MD †
UT Southwestern Simmons
Comprehensive Cancer Center

Gregory A. Durm, MD, MS †
Indiana University Melvin and Bren Simon
Comprehensive Cancer Center

NCCN
Kristina Gregory, RN, MSN
Lisa Hang, PhD

Scott Gettinger, MD † ‡
Yale Cancer Center/Smilow Cancer Hospital

Travis E. Grotz, MD ¶
Mayo Clinic Comprehensive Cancer Center

Matthew A. Gubens, MD, MS †
UCSF Helen Diller Family
Comprehensive Cancer Center

Aditya Juloori, MD §
The UChicago Medicine Comprehensive Cancer
Center

Rudy P. Lackner, MD ¶
Fred & Pamela Buffett Cancer Center

Michael Lanuti, MD ¶
Mass General Cancer Center

Jules Lin, MD ¶
University of Michigan Rogel Cancer Center

Billy W. Loo, Jr., MD, PhD §
Stanford Cancer Institute

Christine M. Lovly, MD, PhD †
Vanderbilt-Ingram Cancer Center

Fabien Maldonado, MD £
Vanderbilt-Ingram Cancer Center

Erminia Massarelli, MD, PhD, MS †
City of Hope National Medical Center

Daniel Morgensztern, MD †
Siteman Cancer Center at Barnes-Jewish Hospital
and Washington University School of Medicine

Trey C. Mullikin, MD §
Duke Cancer Institute

Thomas Ng, MD ¶
The University of Tennessee
Health Science Center

Dawn Owen, MD, PhD §
Mayo Clinic Comprehensive Cancer Center

Dwight H. Owen, MD, MSc †
The Ohio State University Comprehensive
Cancer Center - James Cancer Hospital
and Solove Research Institute

Sandip P. Patel, MD ‡ † ‡
UC San Diego Moores Cancer Center

Tejas Patil, MD †
University of Colorado Cancer Center

Patricio M. Polanco, MD ¶
UT Southwestern Simmons
Comprehensive Cancer Center

Jonathan Riess, MD ‡
UC Davis Comprehensive Cancer Center

Theresa A. Shapiro, MD, PhD ≠ ‡
The Sidney Kimmel Comprehensive
Cancer Center at Johns Hopkins

Aditi P. Singh, MD †
Abramson Cancer Center at the
University of Pennsylvania

James Stevenson, MD †
Case Comprehensive Cancer Center/
University Hospitals Seidman Cancer Center
and Cleveland Clinic Taussig Cancer Institute

Alda Tam, MD ϕ
The University of Texas
MD Anderson Cancer Center

Tawee Tanvetyanon, MD, MPH †
Moffitt Cancer Center

Jane Yanagawa, MD ¶
UCLA Jonsson Comprehensive Cancer Center

Stephen C. Yang, MD ¶
The Sidney Kimmel Comprehensive
Cancer Center at Johns Hopkins

Edwin Yau, MD, PhD †
Roswell Park Comprehensive Cancer Center

Continue

[NCCN Guidelines Panel Disclosures](#)

‡ Hematology/Hematology oncology	¶ Surgery/Surgical oncology
‡ Internal medicine	ϕ Diagnostic/Interventional radiology
† Medical oncology	£ Pulmonary medicine
≠ Pathology	* Discussion Section Writing Committee
¥ Patient advocacy	
§ Radiation oncology/Radiotherapy	



National
Comprehensive
Cancer
Network®

NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

[NCCN Guidelines Index](#)
[Table of Contents](#)
[Discussion](#)

[NCCN Non-Small Cell Lung Cancer Panel Members](#) [Summary of Guidelines Updates](#)

[Lung Cancer Prevention and Screening \(PREV-1\)](#)

[Clinical Presentation and Risk Assessment \(DIAG-1\)](#)

[Initial Evaluation and Clinical Stage \(NSCL-1\)](#)

[Evaluation and Treatment:](#)

- [Stage IA \(T1abc, N0\) \(NSCL-2\)](#)
- [Stage IB \(peripheral T2a, N0\), Stage I \(central T1abc–T2a, N0\), Stage II \(T1abc–2ab, N1; T2b, N0\), Stage IIB \(T3, N0\), and Stage IIIA \(T3, N1\) \(NSCL-3\)](#)
- [Stage IIB \(T3 invasion, N0\) and Stage IIIA \(T4 extension, N0–1; T3, N1; T4, N0–1\) \(NSCL-5\)](#)
- [Stage IIIA \(T1–2, N2\); Stage IIIB \(T3, N2\); Separate Pulmonary Nodule\(s\) \(Stage IIB, IIIA, IV\) \(NSCL-8\)](#)
- [Multiple Lung Cancers \(N0–1\) \(NSCL-12\)](#)
- [Stage IIIB \(T1–2, N3\); Stage IIIC \(T3, N3\) \(NSCL-13\)](#)
- [Stage IIIB \(T4, N2\); Stage IIIC \(T4, N3\); Stage IVA, M1a: Pleural or Pericardial Effusion \(NSCL-14\)](#)
- [Stage IVA, M1b \(NSCL-15\)](#)

[Surveillance After Completion of Definitive Therapy \(NSCL-17\)](#)

[Therapy for Recurrence and Metastasis \(NSCL-18\)](#)

[Systemic Therapy for Advanced or Metastatic Disease \(NSCL-19\)](#)

[Principles of Pathologic Review \(NSCL-A\)](#)

[Principles of Surgical Therapy \(NSCL-B\)](#)

[Principles of Radiation Therapy \(NSCL-C\)](#)

[Principles of Image-Guided Thermal Ablation Therapy \(NSCL-D\)](#)

[Perioperative Systemic Therapy \(NSCL-E\)](#)

[Concurrent Chemoradiation Regimens \(NSCL-F\)](#)

[Cancer Survivorship Care \(NSCL-G\)](#)

[Principles of Molecular and Biomarker Analysis \(NSCL-H\)](#)

[Emerging Biomarkers to Identify Novel Therapies for Patients with Metastatic NSCLC \(NSCL-I\)](#)

[Molecular and Biomarker-Directed Therapy for Advanced or Metastatic Disease \(NSCL-J\)](#)

[Systemic Therapy for Advanced or Metastatic Disease \(NSCL-K\)](#)

Clinical Trials: NCCN believes that the best management for any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

Find an NCCN Member Institution:
<https://www.nccn.org/home/member-institutions>.

NCCN Categories of Evidence and Consensus: All recommendations are category 2A unless otherwise indicated.

See [NCCN Categories of Evidence and Consensus](#).

NCCN Categories of Preference: All recommendations are considered appropriate.

See [NCCN Categories of Preference](#).

[Staging \(ST-1\)](#)

[Abbreviations \(ABBR-1\)](#)

The NCCN Guidelines® are a statement of evidence and consensus of the authors regarding their views of currently accepted approaches to treatment. Any clinician seeking to apply or consult the NCCN Guidelines is expected to use independent medical judgment in the context of individual clinical circumstances to determine any patient's care or treatment. The National Comprehensive Cancer Network® (NCCN®) makes no representations or warranties of any kind regarding their content, use or application and disclaims any responsibility for their application or use in any way. The NCCN Guidelines are copyrighted by National Comprehensive Cancer Network®. All rights reserved. The NCCN Guidelines and the illustrations herein may not be reproduced in any form without the express written permission of NCCN. ©2024.



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

Terminologies in all NCCN Guidelines are being actively modified to advance the goals of equity, inclusion, and representation.

Updates in Version 3.2024 of the NCCN Guidelines for Non-Small Cell Lung Cancer from Version 2.2024 include:

[NSCL-18](#)

- Footnote II added: An FDA-approved biosimilar is an appropriate substitute.

Updates in Version 2.2024 of the NCCN Guidelines for Non-Small Cell Lung Cancer from Version 1.2024 include:

[NSCL-31](#)

- Subsequent Therapy
 - Asymptomatic
 - ◊ *Repotrectinib (if not previously given)* or lorlatinib
 - Symptomatic; Brain
 - ◊ Entrectinib (if previously treated with crizotinib or certinib) or repotrectinib (if previously treated with crizotinib or ceritinib *or entrectinib*) or lorlatinib (if not previously given)
 - Symptomatic; Systemic -- Multiple lesions
 - ◊ *Repotrectinib (if not previously given)* or lorlatinib

[NSCL-J 1 of 6](#)

- *EGFR* S768I, L861Q, and/or G719X
 - Subsequent therapy
 - ◊ Amivantamab-vmjw + carboplatin + pemetrexed (nonsquamous) added (as per NSCL-24/NSCL-22)

[NSCL-J 6 of 6](#)

- Reference updated:
 - 32 Drilon A, et al. Repotrectinib in ROS1 fusion-positive non-small-cell lung cancer. *N Engl J Med* 2024;390:118-131.

Updates in Version 1.2024 of the NCCN Guidelines for Non-Small Cell Lung Cancer from Version 5.2023 include:

[DIAG-2](#)

- Footnote i modified
 - First sentence modified: FDG-PET/CT performed skull base to *mid-thigh knees or whole body*. (also applies to DIAG-3, footnote k on NSCL-3, NSCL-5, NSCL-8, NSCL-11, NSCL-13 through NSCL-15)
 - Third sentence modified: A *false-positive* FDG-PET/CT scan finding can be caused by infection or inflammation, including absence of lung cancer with localized infection, presence of lung cancer with associated (eg, postobstructive) infection, and presence of lung cancer with related inflammation (eg, nodal, parenchymal, pleural). (also applies to DIAG-3)
 - Last sentence added: If a false-negative FDG-PET/CT is due to low tumor avidity and/or low cellularity is suspected, follow-up CT or biopsy are reasonable options. (also applies to DIAG-3)
- Footnote j modified: ~~If empiric therapy is contemplated without tissue confirmation~~ *Prior to treatment*, multidisciplinary evaluation that ~~at least~~ includes *treating physicians and specialists in obtaining tissue diagnosis* (thoracic surgery, interventional pulmonology, and interventional radiology) is required to determine the safest and most efficient approach for biopsy, or to provide consensus that a biopsy is too risky or difficult, *that a clinical diagnosis of lung cancer is appropriate, and that treatment is warranted* and that the patient can proceed with therapy without tissue confirmation. (Jsseldijk MA, et al. *J Thorac Oncol* 2019;14:583-595.) (also applies to DIAG-3, footnote o on NSCL-2, NSCL-3)



Updates in Version 1.2024 of the NCCN Guidelines for Non-Small Cell Lung Cancer from Version 5.2023 include:

DIAG-A 1 of 3

- Bullet 1, including sub-bullets replaced with the following
 - ▶ The decision to pursue preoperative biopsy of a potential stage I lung cancer should be informed by the pre-test probability of malignancy.
 - ◊ Factors that might be considered in pre-test probability assessment include risk factors, radiologic appearance (including comparison to prior chest imaging if available or FDG-PET/CT if performed), and current or prior residence in regions with prevalent endemic infectious lung disease (ie, fungal, mycobacterial), among other potential factors.
 - ◊ Patients with very high pre-test probability of stage IA lung cancer do not require a biopsy before surgery. A biopsy adds time, costs, and procedural risk and may not be needed for treatment decisions.
 - ◊ If a preoperative tissue diagnosis has not been obtained, then an intraoperative diagnosis (ie, wedge resection, needle biopsy) is necessary before lobectomy, bilobectomy, or pneumonectomy.
 - ◊ Situations in which a preoperative biopsy may be appropriate:
 - A non-lung cancer diagnosis that can be diagnosed by minimally invasive biopsy is at least moderately likely (eg, granulomatous nodule due to endemic fungus).
 - Suspected stage IB or higher lung cancer in patients who may be candidates for systemic therapy prior to surgery.
 - An intraoperative diagnosis appears difficult or very risky.
 - To establish the diagnosis prior to stereotactic ablative radiotherapy (SABR)
- Footnote 2 added: Prior to treatment, multidisciplinary evaluation that includes treating physicians and specialists in obtaining tissue diagnosis (thoracic surgery, interventional pulmonology, and interventional radiology) is required to determine the safest and most efficient approach for biopsy, or to provide consensus that a biopsy is too risky or difficult, that a clinical diagnosis of lung cancer is appropriate, and treatment is warranted.

DIAG-A 2 of 3

- Bullet 4; sub-bullet 1
 - ▶ Last diamond added: Left anterior mediastinotomy/Chamberlain

DIAG-A 3 of 3

- Bullet 1; sub-bullet 2
 - ▶ Diamond 2 modified: Patients with ~~peripheral (outer one-third)~~ *pulmonary* nodules may benefit from navigational bronchoscopy (*including robotic*), radial EBUS, or transthoracic needle aspiration (TTNA).
 - ▶ Diamond 3; entry 3 modified with the addition of the 3P nodal station
 - ▶ Diamond 6 modified: Patients with lung cancer with an associated pleural effusion should undergo thoracentesis and cytology. A negative *pleural* cytology result ~~on initial thoracentesis~~ does not exclude pleural involvement. ~~An additional thoracentesis and/or Thoracoscopic evaluation of the pleura should be considered before starting curative intent therapy if pleural fluid is a lymphocytic exudate with negative pleural fluid cytology.~~

NSCL-2

- Medically inoperable category modified (also applies to NSCL-3)
 - ▶ Medically inoperable, *high surgical risk as determined by thoracic surgeon, and those who decline surgery after thoracic surgical consultation*
- Footnote n modified: Image-guided thermal ablation (IGTA) therapy (eg, cryotherapy, microwave, radiofrequency) may be an option for select patients ~~not receiving SABR or definitive RT.~~ (also applies to NSCL-12, NSCL-16, NSCL-18, NSCL-22, NSCL-23, NSCL-28, NSCL-29, NSCL-31)



Updates in Version 1.2024 of the NCCN Guidelines for Non-Small Cell Lung Cancer from Version 5.2023 include:

[NSCL-3](#)

- Following Pretreatment Evaluation
 - Negative mediastinal nodes changed to No nodal disease
 - ◊ Consider adjuvant chemotherapy for high-risk stages IB, II
 - ◊ Treatment row for N1 removed, as the category changed to no nodal disease
 - Positive mediastinal nodes changed to N1 or N2 disease
 - ◊ Stage IIB added with a link to NSCL-8
- Footnote r modified: Examples of high-risk factors may include poorly differentiated tumors (including lung neuroendocrine tumors [excluding well-differentiated neuroendocrine tumors]), ~~vascular invasion, wedge resection, visceral pleural involvement, and unknown lymph node status (Nx).~~ These factors independently may not be an indication and may be considered when determining treatment with adjuvant chemotherapy. (also applies to NSCL-9)

[NSCL-4](#)

- Footnote l added to Adjuvant Treatment with a link to the Principles of Surgical Therapy.

[NSCL-5](#)

- Proximal airway or mediastinum changed to Trachea/carina or mediastinum

[NSCL-6](#)

- Footnote v added: For patients who have received sequential chemoradiation, durvalumab can be considered as consolidation immunotherapy. (also applies to NSCL-7, NSCL-9, NSCL-13, NSCL-14)

[NSCL-7](#)

- Initial Treatment options modified
 - Surgery (preferred) ~~after preoperative systemic therapy, if planned~~
 - ~~Systemic therapy~~ or concurrent chemoradiation ~~or chemotherapy~~
 - ◊ Margins negative after surgery; Following resection after Margins positive
 - Adjuvant Systemic Therapy (NSCL-E) added
- Footnote x added: Resectability should be determined by thoracic surgery evaluation prior to initiation of any therapy.

[NSCL-9](#)

- Mediastinal biopsy categories modified and NSCL-9 now covers 2 pages (NSCL-9 and NSCL-10)
 - T1–3, N0–1 (including T3 with multiple nodules in same lobe) changed to 3 categories
 - ◊ T1, N0
 - Categories of operable and medically inoperable added
 - Operable
 - Surgical exploration and resection + mediastinal lymph node dissection or systematic lymph node sampling ~~after preoperative systemic therapy, if planned~~
 - Adjuvant Treatment (NSCL-4) added
 - Medically inoperable, high surgical risk as determined by thoracic surgeon, and those who decline surgery after thoracic surgical consultation
 - Definitive RT, preferably SABR added
 - ◊ T2a–3, N0
 - Categories of operable and medically inoperable added
 - Operable
 - Adjuvant Treatment (NSCL-4) changed to Adjuvant Systemic Therapy (NSCL-E)
 - Medically inoperable changed as noted above
 - Definitive RT, preferably SABR added
 - Consider adjuvant chemotherapy for high-risk stage II added



Updates in Version 1.2024 of the NCCN Guidelines for Non-Small Cell Lung Cancer from Version 5.2023 include:

NSCL-10

- Mediastinal biopsy categories modified and cover 2 pages (NSCL-9 and NSCL-10)
 - ▶ T1-3, N1 nodes positive, M0
 - ◊ Categories of operable and medically inoperable added, as per NSCL-9
 - Operable
 - Surgical resection + mediastinal lymph node dissection or systematic lymph node sampling after preoperative systemic therapy, if planned
 - Adjuvant Treatment (NSCL-E) added
 - Medically inoperable, high surgical risk as determined by thoracic surgeon, and those who decline surgery after thoracic surgical consultation
 - Definitive concurrent chemoradiation (category 1) added
 - Durvalumab (category 1 stage III; category 2A stage II) added
- T3 (invasion), N2 nodes positive, M0 combined with T1–2, T3 (other than invasive), N2 nodes positive, M0
New category is T1–3, N2 nodes positive, M0
 - ▶ ~~Induction~~ Systemic therapy followed by surgery ± RT
 - ◊ Adjuvant Systemic Therapy (NSCL-E) added
- Footnote y added: Selected patients with N2 disease (fit, single station non-bulky N2, requiring only lobectomy) may be considered for systemic therapy followed by surgery.

NSCL-11

- Separate pulmonary nodule(s), same lobe (T3, N0–1), or ipsilateral non-primary lobe (T4, N0–1)
 - ▶ Margins negative after surgery
 - ◊ Adjuvant Treatment modified: ~~Chemotherapy (category 1)~~ *Adjuvant Systemic Therapy (NSCL-E)* or Sequential chemotherapy + *Consider RT*

NSCL-19

- Footnote II modified: ~~If there is insufficient tissue to allow testing for all of Complete genotyping for EGFR, KRAS, ALK, ROS1, BRAF, NTRK1/2/3, MET, RET, and ERBB2 (HER2) via repeat biopsy and/or plasma testing should be done. Combinations of tissue and plasma testing, either concurrently or in sequence are acceptable. Concurrent testing can improve time to test results and should be considered in the appropriate clinical situation. Negative results (meaning absence of definitive driver mutation) by one method suggests the use of a complementary method. If a clinically actionable marker is found, it is reasonable to start therapy based on the identified marker. If these are not feasible, Treatment is guided by available results and, if unknown, these patients are treated as though they do not have driver oncogenes. (also applies to NSCL-20)~~

NSCL-21

- First-line therapy, Other recommended
 - ▶ Osimertinib + pemetrexed + (cisplatin or carboplatin) (nonsquamous) (category 1) added
- Footnote tt modified: If systemic therapy regimen contains an immune checkpoint inhibitor, physicians should be aware of the long half-life of such drugs and data reporting adverse events when using osimertinib in combination with or following checkpoint inhibitors. *The rate of side effects (pneumonitis) is higher within 3 months.* Schoenfeld AJ, et al. Ann Oncol 2019;30:839-844; Oshima Y, et al. JAMA Oncol 2018;4:1112-1115; Oxnard GR, et al. Ann Oncol 2020;31:507-516; *Gettinger S, et al. J Thorac Oncol 2018;13:1363-1372.* (also applies NSCL-24)

NSCL-22

- Subsequent therapy
 - ▶ Amivantamab-vmjw + carboplatin + pemetrexed (nonsquamous) (category 1) (preferred) added
- Footnote ww modified: Consider a biopsy at time of progression to rule out SCLC transformation (approximately 6%) and *biopsy or plasma testing* to evaluate mechanisms of resistance. (also applies to NSCL-23)



Updates in Version 1.2024 of the NCCN Guidelines for Non-Small Cell Lung Cancer from Version 5.2023 include:

[NSCL-25](#)

- First-line therapy
 - Amivantamab-vmjw + carboplatin + pemetrexed (nonsquamous) (category 1) (preferred) added

[NSCL-26](#)

- First-line therapy
 - Systemic therapy as per NSCL-K changed to Systemic therapy based on PD-L1 status (NSCL-37, NSCL-38)
- Footnote removed: Adagrasib and sotorasib have a similar mechanism of action and it is not recommended to switch between these agents at the time of progression (reworded and added to NSCL-J 1 of 6)

[NSCL-28](#)

- Lorlatinib
 - Qualifier if ALK G1202R removed from the algorithm
 - Footnote ggg added: Lorlatinib is an option for resistant mutations, such as *ALK* G1202R and L1196M (except compound L1196M/G1202R)

[NSCL-30](#)

- Repotrectinib added as a preferred first-line therapy option. (continues on to NSCL-31)
- Footnote iii modified: Entrectinib *or repotrectinib* may be better for patients with brain metastases.

[NSCL-31](#)

- Brain: Entrectinib *or repotrectinib* (if previously treated with crizotinib or ceritinib)

[NSCL-37](#)

- Footnote nnn added: Atezolizumab monotherapy is a treatment option for patients with PS 3, regardless of PD-L1 status. (also applies to NSCL-38, footnote f on NSCL-J 2 of 6, NSCL-J 3 of 6, footnote j on NSCL-K 1 of 5, NSCL-K 2 of 5)

[NSCL-B 2 of 6](#)

- Bullet 2 modified: Sublobar resection - Segmentectomy and wedge resection *should be strongly considered for peripheral T1ab, N0 tumors*
- Bullet 5 modified: Segmentectomy (preferred) or wedge resection is appropriate in selected patients ~~for the following reasons: with poor pulmonary reserve or other major comorbidity that contraindicates lobectomy~~
- The following content was removed
 - Poor pulmonary reserve or other major comorbidity that contraindicates lobectomy
 - Peripheral nodule ≤2 cm with at least one of the following:
 - ◊ Pure AIS histology
 - ◊ Nodule has ≥50% ground-glass appearance on CT
 - ◊ Radiologic surveillance confirms a long doubling time (≥400 days)
- Bullet 6 modified: *Minimally invasive surgery* (VATS or ~~minimally invasive surgery~~ (including robotic-assisted approaches) should be strongly considered for patients with no anatomic or surgical contraindications, as long as there is no compromise of standard oncologic and dissection principles of thoracic surgery. *Robotic surgery should only be initiated by surgeons who have completed and maintained proficiency in the technique.*
- Bullet 8 added: Studies of robotic-assisted pulmonary resection show non-inferiority to traditional VATS approaches when performed by experienced robotic surgeons.



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

Updates in Version 1.2024 of the NCCN Guidelines for Non-Small Cell Lung Cancer from Version 5.2023 include:

[NSCL-B 3 of 6](#)

- Margins and Nodal Assessment
 - ▶ Bullet 2 modified: N1 and N2 node resection and mapping should be a routine component of lung cancer resections—a minimum of *one N1 and three N2 stations* sampled or complete lymph node dissection.
 - ▶ Bullet 3 modified: Formal ipsilateral mediastinal lymph node dissection is indicated for patients undergoing resection for ~~stage IIIA (N2)~~ *stage IIIA (N2)* disease.
 - ▶ Bullet 6 modified: Consider referral to a radiation oncologist for ~~resected stage IIIA-N2~~ *resected stage IIIA-N2* disease.
- The Role of Surgery in Patients with ~~Stage IIIA (N2)~~ *Stage IIIA (N2)* NSCLC

[NSCL-B 6 of 6](#)

- References added:
 - 1 Altorki N, Wang X, Kozono D, et al. Lobar or sublobar resection for peripheral stage IA non-small-cell lung cancer. *N Engl J Med* 2023;388:489-498
 - 2 Lampridis S, Maraschi A, Le Reun C, et al. Robotic versus video-assisted thoracic surgery for lung cancer: short-term outcomes of a propensity matched analysis. *Cancer (Basel)* 2023;15:2391.
 - 3 Ma J, Li X, Zhao S, et al. Robot-assisted thoracic surgery versus video-assisted thoracic surgery for lung lobectomy or segmentectomy in patients with non-small cell lung cancer: a meta-analysis. *BMC Cancer* 2021;21:498.

[NSCL-C 1 of 11](#)

- General Principles
 - ▶ Bullet 4, sentence 2 modified: More advanced technologies are appropriate when needed to deliver curative RT safely. These technologies include (but are not limited to) 4D-CT ~~and/or FDG-PET/CT simulation~~, *FDG-PET/CT and/or MRI simulation*, IMRT/VMAT, IGRT, motion management, and proton therapy (<https://www.astro.org/Daily-Practice/Reimbursement/Model-Policies/Model-Policies>).
 - ▶ Bullet 4, last sentence added: In retrospective studies, intensity-modulated proton therapy (IMPT) has also been shown to reduce the toxicities as compared with 3D-based passive scattering proton therapy in stage III NSCLC.

[NSCL-C 2 of 11](#)

- Radiation Therapy Simulation, Planning, and Delivery
 - ▶ Bullet 6 modified: IGRT—including (but not limited to) orthogonal pair planar imaging and/or volumetric imaging (such as CBCT, CT on rails, *or MRI*)—is recommended when using SABR, 3D-CRT/IMRT, and proton therapy with steep dose gradients around the target, when OARs are in close proximity to high-dose regions, and when using complex motion management techniques.

[NSCL-C 3 of 11](#)

- Early-Stage NSCLC (Stage I, selected node-negative Stage IIA)
 - ▶ Bullet 1 modified: SABR (also known as SBRT) has achieved good primary tumor control rates and overall survival, higher than conventionally fractionated radiotherapy. Although SABR is not proven equivalent to lobectomy, some prospective series have demonstrated similar overall and cancer-specific survival *with reduced acute toxicity*.
 - ▶ Bullet 2 modified: SABR is also an appropriate option for patients with high surgical risk (able to tolerate sublobar resection but not lobectomy [eg, age ≥75 years, poor lung *or cardiac* function]).
 - ▶ Bullet 3 modified: More modestly hypofractionated or dose-intensified conventionally fractionated ~~3D-CRT regimens~~ *highly conformal radiation (IMRT with IGRT preferred)* are less preferred alternatives and may be considered if referral for SABR is not feasible.
- SABR for Node-Negative Early-Stage NSCLC
 - ▶ Bullet 2, sub-bullet 1, sentence 3 modified: For centrally located tumors (defined variably as within 2 cm of the proximal bronchial tree and/or abutting mediastinal pleura) and even ultra-central tumors (defined as abutting the proximal bronchial tree *or, in some definitions, other critical mediastinal structures as well*), 4 to 10 fraction risk-adapted SABR regimens appear to be effective and safe, while 54 to 60 Gy in 3 fractions is unsafe and should be avoided.

[Continued](#)
UPDATES



Updates in Version 1.2024 of the NCCN Guidelines for Non-Small Cell Lung Cancer from Version 5.2023 include:

[NSCL-C 4 of 11](#)

- Locally Advanced NSCLC (Stage II–III)
 - ▶ Bullets 4 and 5 modified: stage IIIA changed to N2.
 - ▶ Last bullet added: In patients with completely resected pN1 receiving adjuvant systemic therapy, PORT is not recommended. PORT may be considered for these patients if they are unable to receive adjuvant systemic therapy.

[NSCL-C 5 of 11](#)

- Palliative RT for Advanced/Metastatic NSCLC
 - ▶ Bullet 2, last sentence added: SABR/SRS has been found in randomized clinical trials to produce better pain and tumor control of spine and non-spine bone metastases than conventionally fractionated palliative RT, and is appropriate especially for patients with longer expected survival.

[NSCL-C 7 of 11](#)

- Table 2. Commonly Used Doses for SABR
 - ▶ Row 4: Example Indications modified: Central ~~or peripheral~~ tumors
 - ▶ Row 5 added: Total dose = 50–60 Gy; # of Fractions = 5; Example Indications = Peripheral tumors

[NSCL-C 9 of 11 through NSCL-C 11 of 11](#)

- References added
 - ▶ 31 Ball D, Mai GT, Vinod S, et al. Stereotactic ablative radiotherapy versus standard radiotherapy in stage 1 non-small-cell lung cancer (TROG 09.02 CHISEL): a phase 3, open-label, randomised controlled trial. *Lancet Oncol* 2019;20:494-503.
 - ▶ 112 Sahgal A, et al. Stereotactic body radiotherapy versus conventional external beam radiotherapy in patients with painful spinal metastases: an open-label, multicenter, randomised, controlled, phase 2/3 trial. *Lancet Oncol* 2021;22:1023-1033.

[NSCL-E 1 of 5](#)

- Neoadjuvant Systemic Therapy
 - ▶ Bullet 2 modified: Test for PD-L1 status, *EGFR* mutations, and *ALK* rearrangements (stages IB–IIIA, IIIB [T3,N2]). *PD-L1 status can be incorporated with other clinical factors to determine patients who may benefit from induction chemotherapy and immunotherapy.*
 - ▶ Bullet 3 added: Clinical trials for neoadjuvant nivolumab + chemotherapy excluded patients harboring *EGFR* mutations and *ALK* rearrangements. Thus, exclusion of these biomarkers, at a minimum, is recommended prior to consideration for neoadjuvant nivolumab + chemotherapy.

[NSCL-E 3 of 5](#)

- Adjuvant Chemotherapy
 - ▶ Bullet 1 modified: Stage IB (T2a, N0) removed.
 - ▶ Bullet 3 modified: Test for PD-L1 status, *EGFR* mutations, and *ALK* rearrangements (stages ~~IB~~–IIIA, IIIB [T3,N2]).

[NSCL-E 4 of 5](#)

- Pembrolizumab
 - ▶ Sub-bullet 2 modified: For patients with completely resected stage ~~IB~~–IIIA or stage IIIB (T3, N2) NSCLC who received previous neoadjuvant pembrolizumab + chemotherapy (category 1).
- Footnote c added: In general, perioperative therapy should be given as a single regimen and change of immunotherapy is not recommended.



Updates in Version 1.2024 of the NCCN Guidelines for Non-Small Cell Lung Cancer from Version 5.2023 include:

[NSCL-F 1 of 2](#)

- Additional cycles of pemetrexed removed from bullet and added to a footnote.
- Additional cycles of paclitaxel and carboplatin removed from bullets and added to a footnote.
- Footnote e modified into 2 new footnotes
 - ▶ Footnote e: If using durvalumab, additional chemotherapy after radiation is not recommended. *If not using durvalumab, an additional 4 cycles of pemetrexed 500 mg/m² may be used.*
 - ▶ Footnote f: If using durvalumab, additional chemotherapy after radiation is not recommended. *If not using durvalumab, an additional 2 cycles every 21 days of paclitaxel 200 mg/m² and carboplatin AUC 6 may be used.*
- Footnote g added: In patients with tumors that are positive for *EGFR* exon 19 deletion of exon 21 L858R mutations there is risk of toxicity when TKI is administered in temporal proximity to IO therapy.
- Footnote h added: For patients who have received sequential chemoradiation, durvalumab can be considered as consolidation immunotherapy.

NSCL-H

- Terminology modified to use circulating tumor DNA (ctDNA) and remove cell-free DNA (cf DNA)

[NSCL-H 1 of 8](#)

- Molecular Diagnostic Studies in NSCLC
 - ▶ Bullet 3, sub-bullet 6 added: Understanding what constitutes an informative versus a null result. A positive finding for a known oncogenic driver is considered an informative result. Absence of a known oncogenic driver, including results that show only "passenger" alterations, might be considered null depending on the context of the specimen limitations and clinical situation. Specifically, technical issues related to tumor cellularity in tissue testing or burden of disease in circulating tumor DNA (ctDNA) testing are considerations in interpretation of findings. Additional considerations are elaborated in sections below.
- Testing Methodologies
 - ▶ Sub-bullet 1, diamond 1, entry 1, sentence 2 modified: NCCN acknowledges that many currently available NGS-based assays used to fully genotype NSCLC are larger than the 50-gene limit threshold utilized by Current Procedural Terminology (CPT) coding convention; as a result, *utilization of panels with greater than 50 genes for panels larger than 50 genes*, it may be practical to follow these recommendations.
 - ▶ Sub-bullet 1, diamond 1, entry 2 added: Although broad-based genomic testing approaches are preferred, in some clinical situations rapid testing may be warranted and should be followed up with broad-based genomic testing.

[NSCL-H 2 of 8](#)

- Specimen Selection
 - ▶ Sub-bullet 2, diamond 2 modified: If an assay has a technical failure related to an insufficient quantity, consideration of an alternate testing modality or *procurement of additional tissue may be warranted is recommended to achieve broad molecular profiling.*

[NSCL-H 6 of 8](#)

- PD-L1
 - ▶ Sub-bullet 2, diamond 1, entry 2 modified: While some clones for PD-L1 IHC are FDA-approved for specific indications, use of multiple IHC tests is not *necessary routinely recommended*, provided any individual IHC test has been internally validated for comparability for categorical results against the FDA-approved clone.

**Updates in Version 1.2024 of the NCCN Guidelines for Non-Small Cell Lung Cancer from Version 5.2023 include:****NSCL-H 8 of 8**

- ~~Plasma Cell-Free/Circulating Tumor DNA (ctDNA) Testing~~
 - ▶ Sub-bullet 2 added: ctDNA is not routinely recommended in settings other than advanced/metastatic disease. For stages I–III, tissue-based testing is preferred. Metastatic disease confined to the thorax may have a higher yield with tissue-based testing.
 - ▶ Sub-bullet 3 modified: Some laboratories offer testing for molecular alterations examining nucleic acids in peripheral circulation, most commonly in processed plasma (sometimes referred to as "liquid biopsy").
 - ▶ Sub-bullet 4 added:
Studies have demonstrated ctDNA and tissue testing to have very high specificity. Both ctDNA and tissue testing have appreciable false-negative rates, supporting the complementarity of these approaches, and data support complementary testing to reduce turnaround time and increase yield of targetable alteration detection.
 - ◊ Limitations of ctDNA testing that can impact interpretation include:
 - Low tumor fraction/ctDNA; some assays include a measure of ctDNA fraction, which can aid in identification of situations in which low ctDNA fraction might suggest compromised sensitivity
 - The presence of mutations from sites other than the target lesion, most commonly clonal hematopoiesis of indeterminate potential (CHIP) or post-chemotherapy marrow clones. KRAS and TP53 can be seen in either of these circumstances
 - The inherent ability of the assay to detect fusions or other genomic variation of relevance
 - ◊ Limitations of tissue testing that can impact interpretation include:
 - Low tumor percent in a sample not sufficiently mitigated by tumor enrichment or high analytic sensitivity methods
 - The inherent ability of the assay to detect fusions or other genomic variation of relevance
- Bullets removed:
 - ▶ Published guidelines elaborating standards for analytical performance characteristics of ctDNA have not been established, and in contrast to tissue-based testing, no guidelines exist regarding the recommended performance characteristics of this type of testing.
 - ▶ Cell-free tumor DNA testing can identify alterations that are unrelated to a lesion of interest, for example, clonal hematopoiesis of indeterminate potential (CHIP)
 - ▶ Studies have demonstrated cell-free tumor DNA testing to generally have very high specificity, but significantly compromised sensitivity, with up to a 30% false-negative rate; however, data support complementary testing to reduce turnaround time and increase yield of targetable alteration detection.
 - ▶ The use of cell-free/circulating tumor DNA testing can be considered in specific clinical circumstances, most notably:
 - ◊ If a patient is medically unfit for invasive tissue sampling
 - ◊ In the initial diagnostic setting, if following pathologic confirmation of a NSCLC diagnosis there is insufficient material for molecular analysis, cell-free/circulating tumor DNA can be used; however, follow-up tissue-based analysis for all patients in which an oncogenic driver is not identified should be planned (see NSCL-18 for oncogenic drivers with available targeted therapy options).
 - ◊ In the initial diagnostic setting, if tissue-based testing does not completely assess all recommended biomarkers owing to tissue quantity or testing methodologies available, consider repeat biopsy and/or cell-free/circulating tumor DNA testing.
 - ◊ In the initial diagnostic setting, if the feasibility of timely tissue-based testing is uncertain, concurrent cfDNA testing may aid in biomarker evaluation for treatment selection, provided negative results are considered per above limitations.

NSCL-I

- Footnote b added: In patients with *EGFR* mutant NSCLC who develop high-level *MET* amplifications, administration of these agents with continuation of osimertinib is acceptable.



Updates in Version 1.2024 of the NCCN Guidelines for Non-Small Cell Lung Cancer from Version 5.2023 include:

[NSCL-J 1 of 6](#)

- *EGFR* Exon 19 Deletion or Exon 21 L858R
 - First-line therapy
 - ◊ Osimertinib + pemetrexed + (cisplatin or carboplatin) (nonsquamous) added
 - Subsequent therapy
 - ◊ Amivantamab-vmjw + carboplatin + pemetrexed (nonsquamous) added
- *EGFR* Exon 20 Insertion Mutation
 - First-line therapy
 - ◊ Amivantamab-vmjw + carboplatin + pemetrexed (nonsquamous) added
- *ROS1* Rearrangement
 - First-line therapy and Subsequent therapy
 - ◊ Repotrectinib added
- Footnote d added: For agents with a similar mechanism of action, it is not recommended to switch between these drugs at the time of progression.

[NSCL-J 2 of 6](#)

- Adenocarcinoma, Large Cell, NSCLC NOS
 - Cemiplimab-rwlc + pemetrexed + (carboplatin or cisplatin) (category 1) moved from other recommended to preferred (also applies to NSCL-J 3 of 6, NSCL-K 1 of 5)
- Squamous Cell Carcinoma
 - Cemiplimab-rwlc + paclitaxel + (carboplatin or cisplatin) (category 1) moved from other recommended to preferred (also applies to NSCL-J 3 of 6, NSCL-K 2 of 5)

[NSCL-J 5 of 6](#)

- References added:
 - 7 Planchard D, et al. Osimertinib with or without chemotherapy in *EGFR*-mutated advanced NSCLC. *N Eng J Med* 2023;389:1935-1948.
 - 11 Passaro A, et al. Amivantamab plus chemotherapy with and without lazertinib in *EGFR*-mutant advanced NSCLC after disease progression on osimertinib: primary results from the phase III MARIPOSA-2 study. *Ann Oncol* 2023;S0923-7534:04281-04283.
 - 14 Zhou C, et al. Amivantamab plus chemotherapy in NSCLC with *EGFR* Exon 20 insertions. *N Engl J Med* 2023;389:2039-2051.

[NSCL-J 6 of 6](#)

- Reference added:
 - 32 Cho BC, et al. OA03.06 Repotrectinib in patients with *ROS1* fusion-positive (*ROS1*+) NSCLC: Update from the pivotal phase 1/2 TRIDENT-1 trial. *J Thorac Oncol* 2023;18:S50-S51.

[NSCL-K 1 of 5](#)

- Category designation for tremelimumab-actl-combination regimens changed from category 2A to category 1. (also applies to NSCL-K 2 of 5)

[NSCL-K 3 of 5](#)

- Footnote k added: Monitoring During Subsequent Therapy or Maintenance Therapy: Response assessment with CT of known or high-risk sites of disease with or without contrast every 6–12 weeks. Timing of CT scans within Guidelines parameters is a clinical decision. (also applies to NSCL-K 4 of 5)

[NSCL-K 4 of 5](#)

- Footnote d added: Contraindications for treatment with PD-1/PD-L1 inhibitors may include active or previously documented autoimmune disease and/or current use of immunosuppressive agents; some oncogenic drivers (ie, *EGFR* exon 19 deletion or L858R, *ALK* rearrangements) have been shown to be associated with less benefit from PD-1/PD-L1 inhibitors.

**LUNG CANCER PREVENTION AND SCREENING**

- Lung cancer is a unique disease in that the major etiologic agent is an addictive product that is made and promoted by an industry. Approximately 85% to 90% of cases are caused by voluntary or involuntary (second-hand) cigarette smoking. Reduction of lung cancer mortality will require effective public health policies to prevent initiation of smoking, U.S. Food and Drug Administration (FDA) oversight of tobacco products, and other tobacco control measures.
- Persistent smoking is associated with second primary cancers, treatment complications, drug interactions, other tobacco-related medical conditions, diminished quality of life, and reduced survival.
- Reports from the Surgeon General on both active smoking (http://www.cdc.gov/tobacco/data_statistics/sgr/2004/pdfs/executivesummary.pdf) and second-hand smoke show that both cause lung cancer. The evidence shows a 20% to 30% increase in the risk for lung cancer from second-hand smoke exposure associated with living with a person who smokes (<http://www.ncbi.nlm.nih.gov/books/NBK44324>). Every person should be informed of the health consequences, addictive nature, and mortal threat posed by tobacco consumption and exposure to tobacco smoke, and effective legislative, executive, administrative, or other measures should be contemplated at the appropriate governmental level to protect all persons from exposure to tobacco smoke.
- Further complicating this problem, the delivery system of lung carcinogens also contains the highly addictive substance, nicotine. Reduction of lung cancer mortality will require widespread implementation of Agency for Healthcare Research and Quality (AHRQ) Guidelines (<http://www.ahrq.gov/professionals/clinicians-providers/guidelines-recommendations/tobacco/index.html>) to identify, counsel, and treat patients with nicotine habituation.
- Patients who smoke or who formerly smoked have significant risk for the development of lung cancer; chemoprevention agents are not yet established for these patients. When possible, these patients should be encouraged to enroll in chemoprevention trials.
- Lung cancer screening using low-dose CT (LDCT) is recommended in select patients at high risk for lung cancer who smoke or formerly smoked (see the [NCCN Guidelines for Lung Cancer Screening](#)).
- See the [NCCN Guidelines for Smoking Cessation](#).

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.





NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

CLINICAL PRESENTATION

RISK ASSESSMENT^b



^a Multidisciplinary evaluation including thoracic surgeons, thoracic radiologists, and pulmonologists to determine the likelihood of a cancer diagnosis and the optimal diagnostic or follow-up strategy.

^b Risk calculators can be used to quantify individual patient and radiologic factors but do not replace evaluation by a multidisciplinary diagnostic team with substantial experience in the diagnosis of lung cancer.

^c [Principles of Diagnostic Evaluation \(DIAG-A 1 of 3\)](#).

^d The most important radiologic factor is change or stability compared with a previous imaging study.


Note: All recommendations are category 2A unless otherwise indicated.
Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

FINDINGS



^c [Principles of Diagnostic Evaluation \(DIAG-A 1 of 3\)](#).

^d The most important radiologic factor is change or stability compared with a previous imaging study.

^e Low risk = minimal or absent history of smoking or other known risk factors.

^f High risk = history of smoking or other known risk factors. Known risk factors include history of lung cancer in a first-degree relative or exposure to asbestos, radon, or uranium.

^g Non-solid (ground-glass) nodules may require longer follow-up to exclude indolent adenocarcinoma.

^h Adapted from Fleischner Society Guidelines: MacMahon H, Naidich DP, Goo JM, et al. Guidelines for management of incidental pulmonary nodules detected on CT images: From the Fleischner Society 2017. Radiology 2017;284:228-243. ©Radiological Society of North America. Fleischner Society Guidelines do not direct whether or not contrast is necessary or if an LDCT is appropriate. LDCT is preferred unless there is a reason for contrast enhancement for better diagnostic resolution.

ⁱ FDG-PET/CT performed skull base to mid-thigh. A positive FDG-PET/CT result is defined as a standardized uptake value (SUV) in the lung nodule greater than the baseline mediastinal blood pool. A false-positive FDG-PET/CT scan finding can be caused by infection or inflammation, including absence of lung cancer with localized infection, presence of lung cancer with associated (eg, postobstructive) infection, and presence of lung cancer with related inflammation (eg, nodal, parenchymal, pleural). A false-negative FDG-PET/CT scan can be caused by a small nodule, low cellular density (nonsolid nodule or ground-glass opacity [GGO]), or low tumor avidity for FDG (eg, adenocarcinoma in situ [AIS; previously known as bronchoalveolar carcinoma], carcinoid tumor). If a false-negative FDG-PET/CT is due to low tumor avidity and/or low cellularity is suspected, follow-up CT or biopsy are reasonable options.

^j Prior to treatment, multidisciplinary evaluation that includes treating physicians and specialists in obtaining tissue diagnosis (thoracic surgery, interventional pulmonology, and interventional radiology) is required to determine the safest and most efficient approach for biopsy, or to provide consensus that a biopsy is too risky or difficult, that a clinical diagnosis of lung cancer is appropriate, and that treatment is warranted.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.




NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

FINDINGS

FOLLOW-UP^{c,d,g,h}



^c [Principles of Diagnostic Evaluation \(DIAG-A 1 of 3\)](#).

^d The most important radiologic factor is change or stability compared with a previous imaging study.

^g Non-solid (ground-glass) nodules may require longer follow-up to exclude indolent adenocarcinoma.

^h Adapted from Fleischner Society Guidelines: MacMahon H, Naidich DP, Goo JM, et al. Guidelines for management of incidental pulmonary nodules detected on CT images: From the Fleischner Society 2017. Radiology 2017;284:228-243. ©Radiological Society of North America. Fleischner Society Guidelines do not direct whether or not contrast is necessary or if an LDCT is appropriate. LDCT is preferred unless there is a reason for contrast enhancement for better diagnostic resolution.

ⁱ FDG-PET/CT performed skull base to mid-thigh. A positive FDG-PET/CT result is defined as an SUV in the lung nodule greater than the baseline mediastinal blood pool. A false-positive FDG-PET/CT scan finding can be caused by infection or inflammation, including absence of lung cancer with localized infection, presence of lung cancer with associated (eg, postobstructive) infection, and presence of lung cancer with related inflammation (eg, nodal, parenchymal, pleural). A false-negative FDG-PET/CT scan can be caused by a small nodule, low cellular density (nonsolid nodule or GGO), or low tumor avidity for FDG (eg, AIS [previously known as bronchoalveolar carcinoma], carcinoid tumor). If a false-negative FDG-PET/CT is due to low tumor avidity and/or low cellularity is suspected, follow-up CT or biopsy are reasonable options.

^j Prior to treatment, multidisciplinary evaluation that includes treating physicians and specialists in obtaining tissue diagnosis (thoracic surgery, interventional pulmonology, and interventional radiology) is required to determine the safest and most efficient approach for biopsy, or to provide consensus that a biopsy is too risky or difficult, that a clinical diagnosis of lung cancer is appropriate, and that treatment is warranted.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



PRINCIPLES OF DIAGNOSTIC EVALUATION

- The decision to pursue preoperative biopsy of a potential stage I lung cancer should be informed by the pre-test probability of malignancy.
 - ▶ Factors that might be considered in pre-test probability assessment include risk factors, radiologic appearance (including comparison to prior chest imaging if available or FDG-PET/CT if performed), and current or prior residence in regions with prevalent endemic infectious lung disease (ie, fungal, mycobacterial), among other potential factors.
 - ▶ Patients with very high pre-test probability of stage IA lung cancer do not require a biopsy before surgery. A biopsy adds time, costs, and procedural risk and may not be needed for treatment decisions.
 - ▶ If a preoperative tissue diagnosis has not been obtained, then an intraoperative diagnosis (ie, wedge resection, needle biopsy) is necessary before lobectomy, bilobectomy, or pneumonectomy.
 - ▶ Situations in which a preoperative biopsy may be appropriate:
 - ◊ A non-lung cancer diagnosis that can be diagnosed by minimally invasive biopsy is at least moderately likely (eg, granulomatous nodule due to endemic fungus).
 - ◊ Suspected stage IB or higher lung cancer in patients who may be candidates for systemic therapy prior to surgery.
 - ◊ An intraoperative diagnosis appears difficult or very risky.¹
 - ◊ To establish the diagnosis prior to stereotactic ablative radiotherapy (SABR).²

¹ Patients require tissue confirmation of non-small cell lung cancer (NSCLC) before a lobectomy, bilobectomy, or pneumonectomy. If a preoperative or intraoperative tissue diagnosis appears risky or unreliable, multidisciplinary evaluation that at least includes interventional radiology, thoracic surgery, and interventional pulmonology is recommended to determine the safest and most efficient approach for biopsy, or to provide consensus that a biopsy is too risky or difficult and that the patient can proceed with anatomic resection without tissue confirmation (IJsseldijk MA, et al. J Thorac Oncol 2019;14:583-595).

² Prior to treatment, multidisciplinary evaluation that includes treating physicians and specialists in obtaining tissue diagnosis (thoracic surgery, interventional pulmonology, and interventional radiology) is required to determine the safest and most efficient approach for biopsy, or to provide consensus that a biopsy is too risky or difficult, that a clinical diagnosis of lung cancer is appropriate, and that treatment is warranted.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



PRINCIPLES OF DIAGNOSTIC EVALUATION

- A preoperative bronchoscopy may also be preferred for tissue diagnosis and/or mediastinal staging (endobronchial ultrasound [EBUS]).
- If a bronchoscopy has not been previously performed for diagnosis or staging, bronchoscopy should be performed during the planned surgical resection, rather than as a separate procedure.
 - ▶ Bronchoscopy is required before surgical resection ([NSCL-2](#)).
 - ▶ A separate bronchoscopy may not be needed for treatment decisions before the time of surgery and adds time, costs, and procedural risk.
 - ▶ A preoperative bronchoscopy may be appropriate if a central tumor requires pre-resection evaluation for biopsy, surgical planning (eg, potential sleeve resection), or preoperative airway preparation (eg, coring out an obstructive lesion).
 - ▶ A preoperative bronchoscopy may also be preferred for tissue diagnosis and/or mediastinal staging (EBUS).
- Invasive mediastinal staging is recommended before surgical resection for most patients with clinical stage I or II lung cancer ([NSCL-2](#)). For patients undergoing EBUS/endoscopic ultrasound (EUS) staging, this most commonly should be a separate procedure to allow for pathologic evaluation.
 - ▶ Patients having mediastinoscopy should preferably undergo invasive mediastinal staging (mediastinoscopy) as the initial step before the planned resection (during the same anesthetic procedure), rather than as a separate procedure.
 - ▶ A separate staging procedure adds time, costs, coordination of care, inconvenience, and an additional anesthetic risk.
 - ▶ Preoperative invasive mediastinal staging may be appropriate for a strong clinical suspicion of N2 or N3 nodal disease or when intraoperative cytology or frozen section analysis is not available.
- In patients with suspected non-small cell lung cancer (NSCLC), many techniques are available for tissue diagnosis.
 - ▶ Diagnostic tools that should be routinely available include:
 - ◊ Sputum cytology
 - ◊ Bronchoscopy with biopsy and transbronchial needle aspiration (TBNA)
 - ◊ Image-guided transthoracic needle core biopsy (preferred) or fine-needle aspiration (FNA)
 - ◊ Thoracentesis
 - ◊ Mediastinoscopy
 - ◊ Video-assisted thoracic surgery (VATS) and open surgical biopsy
 - ◊ Left anterior mediastinotomy/Chamberlain
 - ▶ Diagnostic tools that provide important additional strategies for biopsy include:
 - ◊ EBUS-guided biopsy
 - ◊ EUS-guided biopsy
 - ◊ Navigational bronchoscopy
 - ◊ Robotic bronchoscopy

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

**PRINCIPLES OF DIAGNOSTIC EVALUATION**

- **The preferred diagnostic strategy for an individual patient depends on the size and location of the tumor, the presence of mediastinal or distant disease, patient characteristics (such as pulmonary pathology and/or other significant comorbidities), and local experience and expertise.**
 - **Factors to be considered in choosing the optimal diagnostic step include:**
 - ◊ **Anticipated diagnostic yield (sensitivity)**
 - ◊ **Diagnostic accuracy including specificity and particularly the reliability of a negative diagnostic study (ie, true negative)**
 - ◊ **Adequate volume of tissue specimen for diagnosis and molecular testing**
 - ◊ **Invasiveness and risk of procedure**
 - ◊ **Efficiency of evaluation**
 - **Access and timeliness of procedure**
 - **Concomitant staging is beneficial, because it avoids additional biopsies or procedures. It is preferable to biopsy the pathology that would confer the highest stage (ie, to biopsy a suspected metastasis or mediastinal lymph node rather than the pulmonary lesion). Therefore, FDG-PET/CT imaging is frequently best performed before a diagnostic biopsy site is chosen in cases of high clinical suspicion for aggressive, advanced-stage tumors.**
 - ◊ **Technologies and expertise available**
 - ◊ **Tumor viability at proposed biopsy site from FDG-PET/CT imaging**
 - **Decisions about the optimal diagnostic steps for suspected stage I to III lung cancer should be made by thoracic radiologists, interventional radiologists, thoracic surgeons, and pulmonologists who devote a significant portion of their practice to thoracic oncology. Multidisciplinary evaluation should also include a pulmonologist or thoracic surgeon with expertise in advanced bronchoscopic techniques for diagnosis. The least invasive biopsy with the highest yield is preferred as the first diagnostic study.**
 - ◊ **Patients with central masses and suspected endobronchial involvement should undergo bronchoscopy.**
 - ◊ **Patients with pulmonary nodules may benefit from navigational bronchoscopy (including robotic), radial EBUS, or transthoracic needle aspiration (TTNA).**
 - ◊ **Patients with suspected nodal disease should be biopsied by EBUS, EUS, navigational bronchoscopy, or mediastinoscopy.**
 - **EBUS provides access to nodal stations 2R/2L, 3P, 4R/4L, 7, 10R/10L, 11–13, and other hilar nodal stations if necessary.**
 - **An EBUS-TBNA negative for malignancy in a clinically (FDG-PET/CT and/or CT) positive mediastinum should undergo subsequent mediastinoscopy prior to surgical resection.**
 - **EUS-guided biopsy provides additional access to stations 3P, 5, 7, 8, and 9 lymph nodes if these are clinically suspicious.**
 - **TTNA and anterior mediastinotomy (ie, Chamberlain procedure) provide additional access to anterior mediastinal (stations 5 and 6) lymph nodes if these are clinically suspicious. If TTNA is not possible due to proximity to aorta, VATS biopsy is also an option.**
 - ◊ **EUS also provides reliable access to the left adrenal gland.**
 - ◊ **Rapid on-site evaluation (ROSE), when available, helps to increase diagnostic and molecular yield.**
 - ◊ **Patients with lung cancer with an associated pleural effusion should undergo thoracentesis and cytology. A negative pleural cytology result does not exclude pleural involvement. Thorascopic evaluation of the pleura should be considered before starting curative intent therapy if pleural fluid is a lymphocytic exudate with negative pleural fluid cytology.**
 - ◊ **Patients suspected of having a solitary site of metastatic disease should have tissue confirmation of that site if feasible.**
 - ◊ **Patients suspected of having metastatic disease should have confirmation from one of the metastatic sites if feasible.**
 - ◊ **Patients who may have multiple sites of metastatic disease—based on a strong clinical suspicion—should have biopsy of the primary lung lesion or mediastinal lymph nodes if it is technically difficult or very risky to biopsy a metastatic site.**

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

PATHOLOGIC DIAGNOSIS OF NSCLC

INITIAL EVALUATION

- Pathology review^a
- H&P (include performance status + weight loss)^b
- CT chest and upper abdomen with contrast, including adrenals
- CBC, platelets
- Chemistry profile
- Smoking cessation advice, counseling, and pharmacotherapy
- ▶ Use the 5 A's Framework:
<http://www.ahrq.gov/clinic/tobacco/5steps.htm>
- Integrate palliative care^c
[NCCN Guidelines for Palliative Care](#)
- For tools to aid in the optimal assessment and management of NSCLC in older adults, see the [NCCN Guidelines for Older Adult Oncology](#)

NSCLC →

CLINICAL STAGE^d

Stage IA, peripheral^e (T1abc, N0)[Pretreatment
Evaluation \(NSCL-2\)](#)

Stage IB, peripheral^e (T2a, N0);
Stage I, central^e (T1abc–T2a, N0);
Stage II (T1abc–T2ab, N1; T2b, N0);
Stage IIB (T3, N0)^f; Stage IIIA (T3, N1)

[Pretreatment
Evaluation \(NSCL-3\)](#)

Stage IIB (T3 invasion, N0);
Stage IIIA (T4 extension, N0–1; T3, N1; T4, N0–1)

[Pretreatment
Evaluation \(NSCL-5\)](#)

Stage IIIA (T1–2, N2); Stage IIIB (T3, N2)

[Pretreatment
Evaluation \(NSCL-8\)](#)

Separate pulmonary nodule(s) (Stage IIB, IIIA, IV)

[Pretreatment
Evaluation \(NSCL-8\)](#)

Multiple lung cancers

[Treatment \(NSCL-12\)](#)

Stage IIIB (T1–2, N3); Stage IIIC (T3, N3)

[Pretreatment
Evaluation \(NSCL-13\)](#)

Stage IIIB (T4, N2); Stage IIIC (T4, N3)

[Pretreatment
Evaluation \(NSCL-14\)](#)Stage IVA (M1a)^c (pleural or pericardial effusion)[Pretreatment
Evaluation \(NSCL-14\)](#)Stage IVA (M1b)^c[Pretreatment
Evaluation \(NSCL-15\)](#)Stage IVB (M1c)^c disseminated metastases[Systemic
Therapy \(NSCL-19\)](#)^a [Principles of Pathologic Review \(NSCL-A\)](#).^b Enhanced frailty or geriatric assessments may predict complications better following treatment modalities, particularly surgery. A preferred frailty assessment system has not been established.^c Temel JS, et al. N Engl J Med 2010;363:733-742.^d For patients where more than one treatment modality (surgery, radiation therapy [RT], or systemic therapy) is usually considered, a multidisciplinary evaluation should be performed.^e Based on the CT of the chest: Peripheral = outer third of lung; Central = inner two thirds of lung.^f T3, N0 related to size or satellite nodules.**Note:** All recommendations are category 2A unless otherwise indicated.**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.




NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

CLINICAL ASSESSMENT

PRETREATMENT EVALUATION^g

INITIAL TREATMENT



^g Testing is not listed in order of priority and is dependent on clinical circumstances, institutional processes, and judicious use of resources.

^h Methods for evaluation include mediastinoscopy, mediastinotomy, EBUS, EUS, and CT-guided biopsy. An EBUS-TBNA negative for malignancy in a clinically (FDG-PET/CT and/or CT) positive mediastinum should undergo subsequent mediastinoscopy prior to surgical resection.

ⁱ There is low likelihood of positive mediastinal lymph nodes when these nodes are CT and FDG-PET/CT negative in peripheral tumors (outer third of lung) ≤ 3 cm. Thus, pretreatment pathologic mediastinal evaluation is optional in these settings. Invasive mediastinal staging is recommended for central tumors.

^j In patients who are medically inoperable, while mediastinal biopsy is generally preferred, the risks in selected patients may outweigh the benefits.

^k FDG-PET/CT performed skull base to mid-thigh. Positive FDG-PET/CT scan findings for distant disease need pathologic or other radiologic confirmation. If FDG-PET/CT scan is positive in the mediastinum, lymph node status needs pathologic confirmation.

^l [Principles of Surgical Therapy \(NSCL-B\)](#).

^m [Principles of Radiation Therapy \(NSCL-C\)](#).

ⁿ Image-guided thermal ablation (IGTA) therapy (eg, cryotherapy, microwave, radiofrequency) may be an option for select patients.

[Principles of Image-Guided Thermal Ablation Therapy \(NSCL-D\)](#).

^o Prior to treatment, multidisciplinary evaluation that includes treating physicians and specialists in obtaining tissue diagnosis (thoracic surgery, interventional pulmonology, and interventional radiology) is required to determine the safest and most efficient approach for biopsy, or to provide consensus that a biopsy is too risky or difficult, that a clinical diagnosis of lung cancer is appropriate, and that treatment is warranted.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

CLINICAL ASSESSMENT

PRETREATMENT EVALUATION^g

INITIAL TREATMENT

Stage IB
(peripheral
T2a, N0)
Stage I
(central
T1abc–T2a, N0)
Stage II
(T1abc–2ab, N1;
T2b, N0)
Stage IIB
(T3, N0)^f
Stage IIIA
(T3, N1)

- Evaluate for perioperative therapy^p
- PFTs (if not previously done)
- Bronchoscopy
- Pathologic mediastinal lymph node evaluation^h
- FDG-PET/CT scan^k (if not previously done)
- Brain MRI with contrast^q (Stage II, IIIA) (Stage IB [optional])

No nodal
disease

Operable

Surgical exploration and resection^l +
mediastinal lymph node dissection or
systematic lymph node sampling after
preoperative systemic therapy, if planned^p

[Adjuvant
Treatment
\(NSCL-4\)](#)

Medically
inoperable, high
surgical risk as
determined by
thoracic surgeon,^l
and those who
decline surgery after
thoracic surgical
consultation

Definitive RT,
preferably
SABR^{m,o}

Consider
adjuvant
chemotherapy^p
for high-risk
stage II^r

[Surveillance
\(NSCL-17\)](#)

N1 or N2
disease

[Stage IIB \(NSCL-8\)](#)
[Stage IIIA/IIIB \(NSCL-8\)](#) or
[Stage IIIB/IIIC \(NSCL-13\)](#)

^f T3, N0 related to size or satellite nodules.

^g Testing is not listed in order of priority and is dependent on clinical circumstances, institutional processes, and judicious use of resources.

^h Methods for evaluation include mediastinoscopy, mediastinotomy, EBUS, EUS, and CT-guided biopsy. An EBUS-TBNA negative for malignancy in a clinically (FDG-PET/CT and/or CT) positive mediastinum should undergo subsequent mediastinoscopy prior to surgical resection.

^k FDG-PET/CT performed skull base to mid-thigh. Positive FDG-PET/CT scan findings for distant disease need pathologic or other radiologic confirmation. If FDG-PET/CT scan is positive in the mediastinum, lymph node status needs pathologic confirmation.

^l [Principles of Surgical Therapy \(NSCL-B\)](#).

^m [Principles of Radiation Therapy \(NSCL-C\)](#).

^o Prior to treatment, multidisciplinary evaluation that includes treating physicians and specialists in obtaining tissue diagnosis (thoracic surgery, interventional pulmonology, and interventional radiology) is required to determine the safest and most efficient approach for biopsy, or to provide consensus that a biopsy is too risky or difficult, that a clinical diagnosis of lung cancer is appropriate, and that treatment is warranted.

^p [Perioperative Systemic Therapy \(NSCL-E\)](#).

^q If MRI is not possible, CT of head with contrast.

^r Examples of high-risk factors may include poorly differentiated tumors (including lung neuroendocrine tumors [excluding well-differentiated neuroendocrine tumors]). These factors independently may not be an indication and may be considered when determining treatment with adjuvant chemotherapy.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.




NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

FINDINGS AT SURGERY

ADJUVANT TREATMENT¹



Footnotes, [NSCL-4A](#)

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



National
Comprehensive
Cancer
Network®

NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

[NCCN Guidelines Index](#)
[Table of Contents](#)
[Discussion](#)

FOOTNOTES

^l [Principles of Surgical Therapy \(NSCL-B\).](#)

^m [Principles of Radiation Therapy \(NSCL-C\).](#)

^p [Perioperative Systemic Therapy \(NSCL-E\).](#)

^s [Concurrent Chemoradiation Regimens \(NSCL-F\).](#)

^t R0 = no residual tumor, R1 = microscopic residual tumor, R2 = macroscopic residual tumor.

^u Increasing size is an important variable when evaluating the need for adjuvant chemotherapy.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

CLINICAL ASSESSMENT

PRETREATMENT EVALUATION

CLINICAL EVALUATION

Stage IIB (T3 invasion, N0)
Stage IIIA (T4 extension,
N0–1; T3, N1; T4, N0–1)

- Evaluate for perioperative therapy^p
- PFTs (if not previously done)
- Bronchoscopy
- Pathologic mediastinal lymph node evaluation^h
- Brain MRI with contrast^q
- MRI with contrast of spine + thoracic inlet for superior sulcus lesions abutting the spine, subclavian vessels, or brachial plexus
- FDG-PET/CT scan^k (if not previously done)

Superior sulcus tumor → [Treatment \(NSCL-6\)](#)

Chest wall → [Treatment \(NSCL-7\)](#)

Trachea/carina or mediastinum → [Treatment \(NSCL-7\)](#)

Stage IIIA (T4, N0–1) → [Treatment \(NSCL-7\)](#)

Unresectable disease → [Treatment \(NSCL-7\)](#)

Positive mediastinal nodes → [Stage IIIA/IIIB \(NSCL-8\)](#)

Metastatic disease → [Treatment for Metastasis limited sites \(NSCL-15\) or distant disease \(NSCL-18\)](#)

^h Methods for evaluation include mediastinoscopy, mediastinotomy, EBUS, EUS, and CT-guided biopsy. An EBUS-TBNA negative for malignancy in a clinically (FDG-PET/CT and/or CT) positive mediastinum should undergo subsequent mediastinoscopy prior to surgical resection.

^k FDG-PET/CT performed skull base to mid-thigh. Positive FDG-PET/CT scan findings for distant disease need pathologic or other radiologic confirmation. If FDG-PET/CT scan is positive in the mediastinum, lymph node status needs pathologic confirmation.

^p [Perioperative Systemic Therapy \(NSCL-E\)](#).

^q If MRI is not possible, CT of head with contrast.


Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

CLINICAL PRESENTATION**INITIAL TREATMENT****ADJUVANT TREATMENT**^l [Principles of Surgical Therapy \(NSCL-B\)](#).^m [Principles of Radiation Therapy \(NSCL-C\)](#).^s [Concurrent Chemoradiation Regimens \(NSCL-F\)](#).^v For patients who have received sequential chemoradiation, durvalumab can be considered as consolidation immunotherapy.^w MRI with contrast of spine + thoracic inlet for superior sulcus lesions abutting the spine, subclavian vessels, or brachial plexus.**Note:** All recommendations are category 2A unless otherwise indicated.**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.




NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

CLINICAL PRESENTATION^x

INITIAL TREATMENT

ADJUVANT TREATMENT

^l [Principles of Surgical Therapy \(NSCL-B\)](#).^m [Principles of Radiation Therapy \(NSCL-C\)](#).^p [Perioperative Systemic Therapy \(NSCL-E\)](#).^s [Concurrent Chemoradiation Regimens \(NSCL-F\)](#).^t R0 = no residual tumor, R1 = microscopic residual tumor, R2 = macroscopic residual tumor.^v For patients who have received sequential chemoradiation, durvalumab can be considered as consolidation immunotherapy.^x Resectability should be determined by thoracic surgery evaluation prior to initiation of any therapy.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.




NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

CLINICAL ASSESSMENT

PRETREATMENT EVALUATION

MEDIASTINAL BIOPSY FINDINGS AND RESECTABILITY



^h Methods for evaluation include mediastinoscopy, mediastinotomy, EBUS, EUS, and CT-guided biopsy. An EBUS-TBNA negative for malignancy in a clinically (FDG-PET/CT and/or CT) positive mediastinum should undergo subsequent mediastinoscopy prior to surgical resection.

^k FDG-PET/CT performed skull base to mid-thigh. Positive FDG-PET/CT scan findings for distant disease need pathologic or other radiologic confirmation. If FDG-PET/CT scan is positive in the mediastinum, lymph node status needs pathologic confirmation.

^p [Perioperative Systemic Therapy \(NSCL-E\)](#).

^q If MRI is not possible, CT of head with contrast.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.




NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

MEDIASTINAL BIOPSY FINDINGS

INITIAL TREATMENT


ADJUVANT TREATMENT

^l [Principles of Surgical Therapy \(NSCL-B\)](#).^m [Principles of Radiation Therapy \(NSCL-C\)](#).ⁿ Image-guided thermal ablation (IGTA) therapy (eg, cryotherapy, microwave, radiofrequency) may be an option for select patients.^o [Principles of Image-Guided Thermal Ablation Therapy \(NSCL-D\)](#).^p Prior to treatment, multidisciplinary evaluation that includes treating physicians and specialists in obtaining tissue diagnosis (thoracic surgery, interventional pulmonology, and interventional radiology) is required to determine the safest and most efficient approach for biopsy, or to provide consensus that a biopsy is too risky or difficult, that a clinical diagnosis of lung cancer is appropriate, and that treatment is warranted.^r [Perioperative Systemic Therapy \(NSCL-E\)](#).^r Examples of high-risk factors may include poorly differentiated tumors (including lung neuroendocrine tumors [excluding well-differentiated neuroendocrine tumors]). These factors independently may not be an indication and may be considered when determining treatment with adjuvant chemotherapy.**Note:** All recommendations are category 2A unless otherwise indicated.**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

^l [Principles of Surgical Therapy \(NSCL-B\)](#).^m [Principles of Radiation Therapy \(NSCL-C\)](#).^p [Perioperative Systemic Therapy \(NSCL-E\)](#).^s [Concurrent Chemoradiation Regimens \(NSCL-F\)](#).^v For patients who have received sequential chemoradiation, durvalumab can be considered as consolidation immunotherapy.^y Selected patients with N2 disease (fit, single station non-bulky N2, requiring only lobectomy) may be considered for systemic therapy followed by surgery.^z Chest CT with contrast and/or FDG-PET/CT to evaluate progression.

Note: All recommendations are category 2A unless otherwise indicated.


Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

CLINICAL PRESENTATION



^h Methods for evaluation include mediastinoscopy, mediastinotomy, EBUS, EUS, and CT-guided biopsy. An EBUS-TBNA negative for malignancy in a clinically (FDG-PET/CT and/or CT) positive mediastinum should undergo subsequent mediastinoscopy prior to surgical resection.

^k FDG-PET/CT performed skull base to mid-thigh. Positive FDG-PET/CT scan findings for distant disease need pathologic or other radiologic confirmation. If FDG-PET/CT scan is positive in the mediastinum, lymph node status needs pathologic confirmation.

^l [Principles of Surgical Therapy \(NSCL-B\)](#).

^m [Principles of Radiation Therapy \(NSCL-C\)](#).

^q If MRI is not possible, CT of head with contrast.

^p [Perioperative Systemic Therapy \(NSCL-E\)](#).

^s [Concurrent Chemoradiation Regimens \(NSCL-F\)](#).

^t R0 = no residual tumor, R1 = microscopic residual tumor, R2 = macroscopic residual tumor.

^{aa} Lesions with different cell types (eg, squamous cell carcinoma, adenocarcinoma) are usually different primary tumors. This analysis may be limited by small biopsy samples. However, lesions of the same cell type are not necessarily metastases. Single contralateral lung nodules with clinical, radiologic, or pathologic features suggestive of a synchronous primary lung cancer (eg, long disease-free survival, ground glass components, different histologic characteristics) that are amenable to local therapy should be considered as probable separate primary cancers and eligible for local therapy ([NSCL-11](#)). Multiple studies suggest that next-generation sequencing (NGS) testing with broad gene coverage may allow for unambiguous determination of clonal relatedness among separate lung nodules.

^{bb} For guidance regarding the evaluation, workup, and management of subsolid pulmonary nodules, please see the diagnostic evaluation of a nodule suspicious for lung cancer ([DIAG-1](#)).

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.




NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

CLINICAL PRESENTATION

INITIAL TREATMENT


^l [Principles of Surgical Therapy \(NSCL-B\)](#).

^m [Principles of Radiation Therapy \(NSCL-C\)](#).

ⁿ IGTA therapy (eg, cryotherapy, microwave, radiofrequency) may be an option for select patients. [Principles of Image-Guided Thermal Ablation Therapy \(NSCL-D\)](#).

^{cc} Multidisciplinary evaluation including thoracic radiology, pulmonary medicine, thoracic surgery, medical oncology, and radiation oncology.

^{dd} Lesions at low risk of becoming symptomatic can be observed (eg, small subsolid nodules with slow growth). However, if the lesion(s) show accelerating growth or increasing solid component or increasing FDG uptake, even while small, treatment should be considered.

^{ee} Lung-sparing resection is preferred, but tumor distribution and institutional expertise should guide individual treatment planning. Patients should be evaluated in a multidisciplinary setting (ie, surgery, radiation oncology, medical oncology, interventional oncology).

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.




NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

CLINICAL ASSESSMENT

PRETREATMENT EVALUATION

INITIAL TREATMENT



^k FDG-PET/CT performed skull base to mid-thigh. Positive FDG-PET/CT scan findings for distant disease need pathologic or other radiologic confirmation. If FDG-PET/CT scan is positive in the mediastinum, lymph node status needs pathologic confirmation.

^m [Principles of Radiation Therapy \(NSCL-C\)](#).

^q If MRI is not possible, CT of head with contrast.

^s [Concurrent Chemoradiation Regimens \(NSCL-F\)](#).

^v For patients who have received sequential chemoradiation, durvalumab can be considered as consolidation immunotherapy.


Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer



^k FDG-PET/CT performed skull base to mid-thigh. Positive FDG-PET/CT scan findings for distant disease need pathologic or other radiologic confirmation. If FDG-PET/CT scan is positive in the mediastinum, lymph node status needs pathologic confirmation.

^m [Principles of Radiation Therapy \(NSCL-C\)](#).

^q If MRI is not possible, CT of head with contrast.

^s [Concurrent Chemoradiation Regimens \(NSCL-F\)](#).

^v For patients who have received sequential chemoradiation, durvalumab can be considered as consolidation immunotherapy.

^{ff} Most pleural (pericardial) effusions with lung cancer are a result of the tumor. In a few patients, however, multiple microscopic examinations of pleural (pericardial) fluid are negative for tumor, and fluid is non-bloody and not an exudate. If these elements and clinical judgment dictate that the effusion is not related to the tumor, the effusion should be excluded as a staging descriptor.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.




NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

CLINICAL ASSESSMENT

PRETREATMENT EVALUATION

INITIAL TREATMENT^{hh}



^k FDG-PET/CT performed skull base to mid-thigh. Positive FDG-PET/CT scan findings for distant disease need pathologic or other radiologic confirmation. If FDG-PET/CT scan is positive in the mediastinum, lymph node status needs pathologic confirmation.

^q If MRI is not possible, CT of head with contrast.

⁹⁹ Including selected patients with stage M1c and limited number and volume of metastatic lesions amenable to definitive local therapy. Clinical trials have included up to 3 to 5 progressing sites.

^{hh} [NCCN Guidelines for Central Nervous System Cancers](#).

Note: All recommendations are category 2A unless otherwise indicated.


Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

TREATMENT OF THORACIC DISEASE



^h Methods for evaluation include mediastinoscopy, mediastinotomy, EBUS, EUS, and CT-guided biopsy. An EBUS-TBNA negative for malignancy in a clinically (PET and/or CT) positive mediastinum should undergo subsequent mediastinoscopy prior to surgical resection.

^l [Principles of Surgical Therapy \(NSCL-B\)](#).

^m [Principles of Radiation Therapy \(NSCL-C\)](#).

ⁿ IGTA therapy (eg, cryotherapy, microwave, radiofrequency) may be an option for select patients. [Principles of Image-Guided Thermal Ablation Therapy \(NSCL-D\)](#).

^S [Concurrent Chemoradiation Regimens \(NSCL-F\)](#).

^z Chest CT with contrast and/or PET/CT to evaluate progression.

ⁱⁱ Typically, RT (including SABR) or surgical resection.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

**SURVEILLANCE AFTER COMPLETION
OF DEFINITIVE THERAPY**

- No evidence of clinical/radiographic disease**
- Stage I–II (primary treatment included surgery ± chemotherapy)
 - H&P and chest CT^{jj} ± contrast every 6 mo for 2–3 y, then H&P and a low-dose non-contrast-enhanced chest CT annually
 - Stage I–II (primary treatment included RT) or stage III or stage IV (oligometastatic with all sites treated with definitive intent)
 - H&P and chest CT^{jj} ± contrast every 3–6 mo for 3 y, then H&P and chest CT ± contrast every 6 mo for 2 y, then H&P and a low-dose non-contrast-enhanced chest CT annually
 - ◊ Residual or new radiographic abnormalities may require more frequent imaging
 - Smoking cessation advice, counseling, and pharmacotherapy
 - FDG-PET/CT^{kk} or brain MRI is not routinely indicated
 - [Cancer Survivorship Care \(NSCL-G\)](#)

Recurrence →


- FDG-PET/CT
- Brain MRI with contrast^q

Locoregional
recurrence[Therapy for Recurrence
and Metastasis \(NSCL-18\)](#)Distant
metastases[Therapy for Recurrence
and Metastasis \(NSCL-18\)](#)^q If MRI is not possible, CT of head with contrast.^{jj} Timing of CT scans within Guidelines parameters is a clinical decision.^{kk} FDG-PET/CT is currently not warranted in the routine surveillance and follow-up of patients with NSCLC. However, many benign conditions (such as atelectasis, consolidation, and radiation fibrosis) are difficult to differentiate from neoplasm on standard CT imaging, and FDG-PET/CT can be used to differentiate true malignancy in these settings. However, if FDG-PET/CT is to be used as a problem-solving tool in patients after RT, histopathologic confirmation of recurrent disease is needed because areas previously treated with RT can remain FDG avid for up to 2 years.**Note:** All recommendations are category 2A unless otherwise indicated.**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

^l [Principles of Surgical Therapy \(NSCL-B\)](#).^m [Principles of Radiation Therapy \(NSCL-C\)](#).ⁿ IGTA therapy (eg, cryotherapy, microwave, radiofrequency) may be an option for select patients. [Principles of Image-Guided Thermal Ablation Therapy \(NSCL-D\)](#).^q If MRI is not possible, CT of head with contrast.^s [Concurrent Chemoradiation Regimens \(NSCL-F\)](#).^{hh} [NCCN Guidelines for Central Nervous System Cancers](#).^{ll} An FDA-approved biosimilar is an appropriate substitute.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.




NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

CLINICAL PRESENTATION

HISTOLOGIC SUBTYPE^a

BIOMARKER TESTINGⁿⁿ



^a [Principles of Pathologic Review \(NSCL-A\)](#).

^c Temel JS, et al. N Engl J Med 2010;363:733-742.

^{mm} Complete genotyping for *EGFR*, *KRAS*, *ALK*, *ROS1*, *BRAF*, *NTRK1/2/3*, *MET*, *RET*, and *ERBB2 (HER2)* via biopsy and/or plasma testing. Combinations of tissue and plasma testing, either concurrently or in sequence are acceptable. Concurrent testing can improve time to test results and should be considered in the appropriate clinical situation. Negative results (meaning absence of definitive driver mutation) by one method suggests the use of a complementary method. If a clinically actionable marker is found, it is reasonable to start therapy based on the identified marker. Treatment is guided by available results and, if unknown, these patients are treated as though they do not have driver oncogenes.

ⁿⁿ [Principles of Molecular and Biomarker Analysis \(NSCL-H\)](#).

^{oo} The NCCN NSCLC Guidelines Panel strongly advises broader molecular profiling with the goal of identifying rare driver mutations for which effective drugs may already be available, or to appropriately counsel patients regarding the availability of clinical trials. Broad molecular profiling is defined as molecular testing that identifies all biomarkers identified in [NSCL-20](#) in either a single assay or a combination of a limited number of assays, and optimally also identifies emerging biomarkers ([NSCL-I](#)). Tiered approaches based on low prevalence of co-occurring biomarkers are acceptable. Broad molecular profiling is a key component of the improvement of care of patients with NSCLC. [Emerging Biomarkers to Identify Patients for Therapies \(NSCL-I\)](#).

^{pp} Lam VK, et al. Clin Lung Cancer 2019;20:30-36.e3; Sands JM, et al. Lung Cancer 2020;140:35-41.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



TESTING RESULTS^{mm,nn}

<i>EGFR</i> exon 19 deletion or exon 21 L858R mutation positive	NSCL-21
<i>EGFR</i> S768I, L861Q, and/or G719X mutation positive	NSCL-24
<i>EGFR</i> exon 20 insertion mutation positive	NSCL-25
<i>KRAS</i> G12C mutation positive	NSCL-26
<i>ALK</i> rearrangement positive	NSCL-27
<i>ROS1</i> rearrangement positive	NSCL-30
<i>BRAF</i> V600E mutation positive	NSCL-32
<i>NTRK1/2/3</i> gene fusion positive	NSCL-33
<i>MET</i>ex14 skipping mutation positive	NSCL-34
<i>RET</i> rearrangement positive	NSCL-35
<i>ERBB2</i> (<i>HER2</i>) mutation positive	NSCL-36
PD-L1 ≥1% and negative for actionable molecular biomarkers above	NSCL-37
PD-L1 <1% and negative for actionable molecular biomarkers above	NSCL-38

^{mm} Complete genotyping for *EGFR*, *KRAS*, *ALK*, *ROS1*, *BRAF*, *NTRK1/2/3*, *MET*, *RET*, and *ERBB2* (*HER2*) via biopsy and/or plasma testing. Combinations of tissue and plasma testing, either concurrently or in sequence are acceptable. Concurrent testing can improve time to test results and should be considered in the appropriate clinical situation. Negative results (meaning absence of definitive driver mutation) by one method suggests the use of a complementary method. If a clinically actionable marker is found, it is reasonable to start therapy based on the identified marker. Treatment is guided by available results and, if unknown, these patients are treated as though they do not have driver oncogenes.

ⁿⁿ [Principles of Molecular and Biomarker Analysis \(NSCL-H\)](#).

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.




NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

EGFR EXON 19 DELETION OR EXON 21 L858R MUTATIONSⁿⁿ

FIRST-LINE THERAPY^{qq}



ⁿⁿ [Principles of Molecular and Biomarker Analysis \(NSCL-H\)](#).

^{qq} [Molecular or Biomarker-Directed Therapy for Advanced or Metastatic Disease \(NSCL-J\)](#).

^{rr} For performance status 0–4.

^{ss} Criteria for treatment with bevacizumab: nonsquamous NSCLC, and no recent history of hemoptysis.

^{tt} An FDA-approved biosimilar is an appropriate substitute for bevacizumab.

^{uu} If systemic therapy regimen contains an immune checkpoint inhibitor, physicians should be aware of the long half-life of such drugs and data reporting adverse events when using osimertinib in combination with or following checkpoint inhibitors. The rate of side effects (pneumonitis) is higher within 3 months. Schoenfeld AJ, et al. *Ann Oncol* 2019;30:839-844; Oshima Y, et al. *JAMA Oncol* 2018;4:1112-1115; Oxnard GR, et al. *Ann Oncol* 2020;31:507-516; Gettinger S, et al. *J Thorac Oncol* 2018;13:1363-1372.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.




NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

EGFR EXON 19 DELETION OR EXON 21 L858R MUTATIONSⁿⁿ

SUBSEQUENT THERAPY^{qq}



ⁿ IGTA therapy (eg, cryotherapy, microwave, radiofrequency) may be an option for select patients. [Principles of Image-Guided Thermal Ablation Therapy \(NSCL-D\)](#).

ⁿⁿ [Principles of Molecular and Biomarker Analysis \(NSCL-H\)](#).

^{qq} [Molecular or Biomarker-Directed Therapy for Advanced or Metastatic Disease \(NSCL-J\)](#).

^{vv} Beware of flare phenomenon in subset of patients who discontinue tyrosine kinase inhibitor (TKI). If disease flare occurs, restart TKI.

^{ww} Clinical trials have included up to 3 to 5 progressing sites.

^{xx} Consider a biopsy at time of progression to rule out small cell lung cancer (SCLC) transformation (approximately 6%) and biopsy or plasma testing to evaluate mechanisms of resistance. [Principles of Molecular and Biomarker Analysis \(NSCL-H\)](#) and [NCCN Guidelines for Small Cell Lung Cancer](#).

^{yy} Afatinib + cetuximab may be considered in patients with disease progression on EGFR TKI therapy.

^{zz} The data in the second-line setting suggest that programmed death cell protein 1 (PD-1)/PD-L1 inhibitor monotherapy is less effective, irrespective of PD-L1 expression, in *EGFR* exon 19 deletion or exon 21 L858R, *ALK*+ NSCLC.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.




NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

EGFR EXON 19 DELETION OR EXON 21 L858R MUTATIONSⁿⁿ

SUBSEQUENT THERAPY^{qq}



ⁿ IGTA therapy (eg, cryotherapy, microwave, radiofrequency) may be an option for select patients. [Principles of Image-Guided Thermal Ablation Therapy \(NSCL-D\)](#).

ⁿⁿ [Principles of Molecular and Biomarker Analysis \(NSCL-H\)](#).

^{qq} [Molecular or Biomarker-Directed Therapy for Advanced or Metastatic Disease \(NSCL-J\)](#).

^{rr} For performance status 0–4.

^{ss} Criteria for treatment with bevacizumab: nonsquamous NSCLC, and no recent history of hemoptysis.

^{vv} Beware of flare phenomenon in subset of patients who discontinue TKI. If disease flare occurs, restart TKI.

^{ww} Clinical trials have included up to 3 to 5 progressing sites.

^{xx} Consider a biopsy at time of progression to rule out SCLC transformation (approximately 6%) and biopsy or plasma testing to evaluate mechanisms of resistance. [NCCN Guidelines for Small Cell Lung Cancer](#).

^{yy} Afatinib + cetuximab may be considered in patients with disease progression on EGFR TKI therapy.

^{zz} The data in the second-line setting suggest that PD-1/PD-L1 inhibitor monotherapy is less effective, irrespective of PD-L1 expression, in EGFR exon 19 deletion or exon 21 L858R, ALK+ NSCLC.

^{aaa} Plasma or tissue-based testing via broad molecular profiling should be considered at progression, for the T790M mutation and other genomic resistance mechanisms. If plasma-based testing is negative, tissue-based testing with rebiopsy material is strongly recommended. Practitioners may want to consider scheduling the biopsy concurrently with plasma testing referral.

^{bbb} Consider osimertinib (regardless of T790M status) for progressive CNS disease or leptomeningeal disease. In the BLOOM study, osimertinib was used at 160 mg once daily for patients with leptomeningeal disease.

^{ccc} In the randomized phase III trial of dacomitinib, patients with brain metastases were not eligible for enrollment. In the setting of brain metastases, consider other options.


Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



EGFR S768I, L861Q, and/or G719X MUTATIONSⁿⁿ

FIRST-LINE THERAPY^{qq}



ⁿⁿ [Principles of Molecular and Biomarker Analysis \(NSCL-H\)](#).

^{qq} [Molecular or Biomarker-Directed Therapy for Advanced or Metastatic Disease \(NSCL-J\)](#).

^{rr} For performance status 0–4.

^{uu} If systemic therapy regimen contains an immune checkpoint inhibitor, physicians should be aware of the long half-life of such drugs and data reporting adverse events when using osimertinib in combination with or following checkpoint inhibitors. The rate of side effects (pneumonitis) is higher within 3 months. Schoenfeld AJ, et al. Ann Oncol 2019;30:839-844; Oshima Y, et al. JAMA Oncol 2018;4:1112-1115; Oxnard GR, et al. Ann Oncol 2020;31:507-516; Gettinger S, et al. J Thorac Oncol 2018;13:1363-1372.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.




NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

EGFR EXON 20 INSERTION MUTATIONⁿⁿ

FIRST-LINE THERAPY^{ddd}

SUBSEQUENT THERAPY^{qq}



ⁿⁿ [Principles of Molecular and Biomarker Analysis \(NSCL-H\)](#).

^{qq} [Molecular or Biomarker-Directed Therapy for Advanced or Metastatic Disease \(NSCL-J\)](#).

^{ddd} Monitoring During Initial Therapy: Response assessment after 2 cycles, then every 2–4 cycles with CT of known or high-risk sites of disease with or without contrast or when clinically indicated. Timing of CT scans within Guidelines parameters is a clinical decision.

^{eee} In general, 4 cycles of initial systemic therapy (ie, with carboplatin or cisplatin) are administered prior to maintenance therapy. However, if patient is tolerating therapy well, consideration can be given to continue to 6 cycles.

^{fff} Monitoring During Subsequent Therapy or Maintenance Therapy: Response assessment with CT of known or high-risk sites of disease with or without contrast every 6–12 weeks. Timing of CT scans within Guidelines parameters is a clinical decision.

Note: All recommendations are category 2A unless otherwise indicated.


Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



KRAS G12C MUTATIONⁿⁿ

FIRST-LINE THERAPY^{ddd}

SUBSEQUENT THERAPY^{qq}



ⁿⁿ [Principles of Molecular and Biomarker Analysis \(NSCL-H\)](#).

^{qq} [Molecular or Biomarker-Directed Therapy for Advanced or Metastatic Disease \(NSCL-J\)](#).

^{ddd} Monitoring During Initial Therapy: Response assessment after 2 cycles, then every 2–4 cycles with CT of known or high-risk sites of disease with or without contrast or when clinically indicated. Timing of CT scans within Guidelines parameters is a clinical decision.

^{eee} In general, 4 cycles of initial systemic therapy (ie, with carboplatin or cisplatin) are administered prior to maintenance therapy. However, if patient is tolerating therapy well, consideration can be given to continue to 6 cycles.

^{fff} Monitoring During Subsequent Therapy or Maintenance Therapy: Response assessment with CT of known or high-risk sites of disease with or without contrast every 6–12 weeks. Timing of CT scans within Guidelines parameters is a clinical decision.

^{ggg} Sotorasib or adagrasib may be used after at least one line of therapy (or second-line and beyond), if no previous *KRAS* G12C-targeted therapy.

Note: All recommendations are category 2A unless otherwise indicated.


Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

ALK REARRANGEMENTⁿⁿ



ⁿⁿ [Principles of Molecular and Biomarker Analysis \(NSCL-H\)](#).
^{qq} [Molecular or Biomarker-Directed Therapy for Advanced or Metastatic Disease \(NSCL-J\)](#).
^{rr} For performance status 0–4.


Note: All recommendations are category 2A unless otherwise indicated.
Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

ALK REARRANGEMENTⁿⁿ



ⁿ IGTA therapy (eg, cryotherapy, microwave, radiofrequency) may be an option for select patients. [Principles of Image-Guided Thermal Ablation Therapy \(NSCL-D\)](#).

ⁿⁿ [Principles of Molecular and Biomarker Analysis \(NSCL-H\)](#).

^{qq} [Molecular or Biomarker-Directed Therapy for Advanced or Metastatic Disease \(NSCL-J\)](#).

^{vv} Beware of flare phenomenon in subset of patients who discontinue TKI. If disease flare occurs, restart TKI.

^{ww} Clinical trials have included up to 3 to 5 progressing sites.

^{zz} The data in the second-line setting suggest that PD-1/PD-L1 inhibitor monotherapy is less effective, irrespective of PD-L1 expression, in *EGFR* exon 19 deletion or exon 21 L858R, *ALK*+ NSCLC.

^{aaa} Plasma or tissue-based testing via broad molecular profiling should be considered at progression for genomic resistance mechanisms. If plasma-based testing is negative, tissue-based testing with rebiopsy material is strongly recommended. Practitioners may want to consider scheduling the biopsy concurrently with plasma testing referral.

^{hhh} Lorlatinib is an option for resistant mutations, such as *ALK* G1202R and L1196M (except compound L1196M/G1202R).

Note: All recommendations are category 2A unless otherwise indicated.


Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

ALK REARRANGEMENTⁿⁿ



ⁿ IGTA therapy (eg, cryotherapy, microwave, radiofrequency) may be an option for select patients. [Principles of Image-Guided Thermal Ablation Therapy \(NSCL-D\)](#).

ⁿⁿ [Principles of Molecular and Biomarker Analysis \(NSCL-H\)](#).

^{qq} [Molecular or Biomarker-Directed Therapy for Advanced or Metastatic Disease \(NSCL-J\)](#).

^{rr} For performance status 0–4.

^{vv} Beware of flare phenomenon in subset of patients who discontinue TKI. If disease flare occurs, restart TKI.

^{ww} Clinical trials have included up to 3 to 5 progressing sites.

^{zz} The data in the second-line setting suggest that PD-1/PD-L1 inhibitor monotherapy is less effective, irrespective of PD-L1 expression, in *EGFR* exon 19 deletion or exon 21 L858R, *ALK*+ NSCLC.

^{aaa} Plasma or tissue-based testing via broad molecular profiling should be considered at progression for genomic resistance mechanisms. If plasma-based testing is negative, tissue-based testing with rebiopsy material is strongly recommended. Practitioners may want to consider scheduling the biopsy concurrently with plasma testing referral.


ⁱⁱⁱ Patients who are intolerant to crizotinib may be switched to ceritinib, alectinib, brigatinib, or lorlatinib.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



ROS1 REARRANGEMENTⁿⁿ



ⁿⁿ [Principles of Molecular and Biomarker Analysis \(NSCL-H\)](#).

^{qq} [Molecular or Biomarker-Directed Therapy for Advanced or Metastatic Disease \(NSCL-J\)](#).

^{rr} For performance status 0–4.

^{vv} Beware of flare phenomenon in subset of patients who discontinue TKI. If disease flare occurs, restart TKI.

^{jjj} Entrectinib or repotrectinib may be better for patients with brain metastases.

Note: All recommendations are category 2A unless otherwise indicated.


Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

ROS1 REARRANGEMENTⁿⁿ



ⁿ IGTA therapy (eg, cryotherapy, microwave, radiofrequency) may be an option for select patients. [Principles of Image-Guided Thermal Ablation Therapy \(NSCL-D\)](#).

ⁿⁿ [Principles of Molecular and Biomarker Analysis \(NSCL-H\)](#).

^{qq} [Molecular or Biomarker-Directed Therapy for Advanced or Metastatic Disease \(NSCL-J\)](#).

^{vv} Beware of flare phenomenon in subset of patients who discontinue TKI. If disease flare occurs, restart TKI.

^{ww} Clinical trials have included up to 3 to 5 progressing sites.

^{aaa} Plasma or tissue-based testing via broad molecular profiling should be considered at progression for genomic resistance mechanisms. If plasma-based testing is negative, tissue-based testing with rebiopsy material is strongly recommended. Practitioners may want to consider scheduling the biopsy concurrently with plasma testing referral.

Note: All recommendations are category 2A unless otherwise indicated.


Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 3.2024


Non-Small Cell Lung Cancer

BRAF V600E MUTATIONⁿⁿ

ⁿⁿ [Principles of Molecular and Biomarker Analysis \(NSCL-H\)](#).^{qq} [Molecular or Biomarker-Directed Therapy for Advanced or Metastatic Disease \(NSCL-J\)](#).^{rr} For performance status 0–4.^{kkk} Single-agent vemurafenib or dabrafenib are treatment options if the combination of dabrafenib + trametinib is not tolerated.**Note:** All recommendations are category 2A unless otherwise indicated.**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NTRK GENE FUSIONⁿⁿ



ⁿⁿ [Principles of Molecular and Biomarker Analysis \(NSCL-H\)](#).

^{qq} [Molecular or Biomarker-Directed Therapy for Advanced or Metastatic Disease \(NSCL-J\)](#).


^{rr} For performance status 0–4.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



METex14 SKIPPING MUTATIONⁿⁿ



ⁿⁿ [Principles of Molecular and Biomarker Analysis \(NSCL-H\)](#).

^{qq} [Molecular or Biomarker-Directed Therapy for Advanced or Metastatic Disease \(NSCL-J\)](#).

^{rr} For performance status 0–4.

^{vv} Beware of flare phenomenon in subset of patients who discontinue TKI. If disease flare occurs, restart TKI.

Note: All recommendations are category 2A unless otherwise indicated.


Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

RET REARRANGEMENTⁿⁿ

ⁿⁿ [Principles of Molecular and Biomarker Analysis \(NSCL-H\)](#).^{qq} [Molecular or Biomarker-Directed Therapy for Advanced or Metastatic Disease \(NSCL-J\)](#).^{rr} For performance status 0–4.^{vv} Beware of flare phenomenon in subset of patients who discontinue TKI. If disease flare occurs, restart TKI.**Note:** All recommendations are category 2A unless otherwise indicated.**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.




NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

ERBB2 (HER2) MUTATIONⁿⁿ

FIRST-LINE THERAPY^{ddd}

SUBSEQUENT THERAPY^{qq}



ⁿⁿ [Principles of Molecular and Biomarker Analysis \(NSCL-H\)](#).

^{qq} [Molecular or Biomarker-Directed Therapy for Advanced or Metastatic Disease \(NSCL-J\)](#).

^{ddd} Monitoring During Initial Therapy: Response assessment after 2 cycles, then every 2–4 cycles with CT of known or high-risk sites of disease with or without contrast or when clinically indicated. Timing of CT scans within Guidelines parameters is a clinical decision.


^{eee} In general, 4 cycles of initial systemic therapy (ie, with carboplatin or cisplatin) are administered prior to maintenance therapy. However, if patient is tolerating therapy well, consideration can be given to continue to 6 cycles.

^{fff} Monitoring During Subsequent Therapy or Maintenance Therapy: Response assessment with CT of known or high-risk sites of disease with or without contrast every 6–12 weeks. Timing of CT scans within Guidelines parameters is a clinical decision.

^{lll} For oncogenic or likely oncogenic *HER2* mutations, refer to definitions at oncokb.org.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.


**PD-L1 POSITIVE (≥1%)^{nn,mmm}****FIRST-LINE THERAPY^{qq,nnn}**ⁿⁿ [Principles of Molecular and Biomarker Analysis \(NSCL-H\)](#).^{qq} [Molecular or Biomarker-Directed Therapy for Advanced or Metastatic Disease \(NSCL-J\)](#).^{mmm} Contraindications for treatment with PD-1/PD-L1 inhibitors may include active or previously documented autoimmune disease and/or current use of immunosuppressive agents; some oncogenic drivers (ie, *EGFR* exon 19 deletion or L858R, *ALK* rearrangements) have been shown to be associated with less benefit from PD-1/PD-L1 inhibitors. If there are contraindications, refer to [NSCL-K 1 of 5 \(adenocarcinoma\)](#) or [NSCL-K 2 of 5 \(squamous cell carcinoma\)](#).ⁿⁿⁿ For patients who require an urgent start to therapy but molecular testing is pending, consider holding immunotherapy for one cycle, unless confirmed that no driver mutations are present.^{ooo} Atezolizumab monotherapy is a treatment option for patients with PS 3, regardless of PD-L1 status.^{ppp} If patient has not received platinum-doublet chemotherapy, refer to "systemic therapy." If patient received platinum chemotherapy and anti-PD-1/PD-L1, refer to "subsequent therapy."**Note: All recommendations are category 2A unless otherwise indicated.****Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.**



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

PD-L1 <1%



^{ddd} Monitoring During Initial Therapy: Response assessment after 2 cycles, then every 2–4 cycles with CT of known or high-risk sites of disease with or without contrast or when clinically indicated. Timing of CT scans within Guidelines parameters is a clinical decision.

^{eee} In general, 4 cycles of initial systemic therapy (ie, with carboplatin or cisplatin) are administered prior to maintenance therapy. However, if patient is tolerating therapy well, consideration can be given to continue to 6 cycles.

^{fff} Monitoring During Subsequent Therapy or Maintenance Therapy: Response assessment with CT of known or high-risk sites of disease with or without contrast every 6–12 weeks. Timing of CT scans within Guidelines parameters is a clinical decision.

^{ooo} Atezolizumab monotherapy is a treatment option for patients with PS 3, regardless of PD-L1 status.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

**PRINCIPLES OF PATHOLOGIC REVIEW****• Pathologic Evaluation**

- ▶ **The purpose of the pathologic evaluation of NSCLC will vary depending on whether the sample 1) is a biopsy or cytology specimen intended for initial diagnosis in a case of suspected NSCLC; 2) is a resection specimen; or 3) is obtained for molecular evaluation in the setting of an established NSCLC diagnosis.**
 - ◊ **In small biopsies or cytology specimens intended for initial diagnosis, the primary purpose is a) to make an accurate diagnosis using the 2021 WHO classification; and b) to preserve the tissue for molecular studies, especially if the patient has advanced-stage disease.**
 - ◊ **In small biopsies of poorly differentiated carcinomas, the terms "non-small cell carcinoma (NSCC)"¹ or "non-small cell carcinoma not otherwise specified (NSCC-NOS)" should be used as little as possible and only when a more specific diagnosis is not possible by morphology and/or special staining.**
 - ◊ **The following terms are acceptable: "NSCC favor adenocarcinoma" and "NSCC favor squamous cell carcinoma." "NSCC-NOS" should be reserved only for cases in which immunohistochemical testing is uninformative or ambiguous (see section on *Immunohistochemistry*).**
 - ◊ **Preservation of material for molecular testing is critical. Efforts should be undertaken to minimize block reorientation and the number of immunohistochemistry (IHC) stains for cases that cannot be classified on histologic examination alone (see section on *Immunohistochemistry*).**
- ▶ **In resection specimens, the primary purpose is a) to classify the histologic type; and b) to determine all staging parameters, as recommended by the American Joint Committee on Cancer (AJCC), including tumor size, extent of invasion, adequacy of surgical margins, and presence or absence of lymph node metastases.**
 - ◊ **The number of involved lymph node stations should be documented since it has prognostic significance (AJCC 8th ed). Direct extension of the primary tumor into an adjacent lymph node is considered as nodal involvement.**
 - ◊ **All lobectomy specimens should be extensively dissected to search for involved lymph nodes.**
- ▶ **In small biopsies or cytology specimens—obtained for molecular testing in the context of an established diagnosis after progression on targeted therapies—the primary purpose is a) to confirm the original pathologic type with minimal use of tissue for IHC only in suspected small cell carcinoma transformation or a different histology; and b) to preserve material for molecular analysis.**
- **Formalin-fixed paraffin-embedded (FFPE) material is suitable for most molecular analyses, except bone biopsies that were previously treated with acid decalcifying solutions. Non-acid decalcification approaches may be successful for subsequent molecular testing. While many molecular pathology laboratories currently also accept cytopathology specimens such as cell blocks, direct smears, or touch preparations, laboratories that do not currently do so are strongly encouraged to identify approaches to testing on non-FFPE cytopathology specimens.**

¹ Non-small cell carcinomas (NSCC, without the L for lung) that show no clear adenocarcinoma or squamous cell carcinoma morphology or immunohistochemical markers are regarded as NSCC-NOS. In this setting, it is recommended that pathologists use the term NSCC rather than NSCLC, because the lack of pneumocyte marker expression in small biopsies or cytology leaves open the possibility of a metastatic carcinoma and the determination of a lung primary must be established clinically after excluding other primary sites.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

[Continue](#)



PRINCIPLES OF PATHOLOGIC REVIEW

NSCLC Classification

- The types of NSCLC are: adenocarcinoma, squamous cell carcinoma, adenosquamous carcinoma, large cell carcinoma, and sarcomatoid carcinoma.
 - ▶ **Squamous cell carcinoma:** A malignant epithelial tumor that either shows keratinization and/or intercellular bridges, or a morphologically undifferentiated NSCC that expresses immunohistochemical markers of squamous cell differentiation.
 - ▶ **Adenocarcinoma:**
 - ◊ For small (<3 cm), resected lesions, determining extent of invasion is critical.
 - **Adenocarcinoma in situ (AIS; formerly BAC):** A small (≤ 3 cm) localized nodule with lepidic growth, mostly non-mucinous, although mucinous types can occur. Multiple synchronous AIS tumors can also occur.
 - **Minimally invasive adenocarcinoma (MIA):** A small (≤ 3 cm) solitary adenocarcinoma with a predominantly lepidic pattern and ≤ 5 mm invasion in greatest dimension. MIA is usually non-mucinous, but rarely may be mucinous. MIA is, by definition, solitary and discrete.
 - **Invasive adenocarcinoma:** A malignant epithelial tumor with glandular differentiation, mucin production, or pneumocyte marker expression. The tumors show an acinar, papillary, micropapillary, lepidic, or solid growth pattern, with either mucin or pneumocyte marker expression. The invasive adenocarcinoma component should be present in at least one focus measuring >5 mm in greatest dimension.
 - **Invasive adenocarcinoma variants:** invasive mucinous adenocarcinoma, colloid adenocarcinoma, fetal adenocarcinoma, and enteric adenocarcinoma.
 - Refer to College of American Pathologists [Protocols](#) for additional information.
 - ▶ **Adenosquamous carcinoma:** A carcinoma showing components of both squamous cell carcinoma and adenocarcinoma, with each component constituting at least 10% of the tumor. Definitive diagnosis requires a resection specimen, although it may be suggested based on findings in small biopsies, cytology, or excisional biopsies. Presence of any adenocarcinoma component in a biopsy specimen that is otherwise squamous should trigger molecular testing.
 - ▶ **Large cell carcinoma:** Undifferentiated NSCC that lacks the cytologic, architectural, and histochemical features of small cell carcinoma, adenocarcinoma, or squamous cell carcinoma. The diagnosis requires a thoroughly sampled resected tumor with immunohistochemical stains that exclude adenocarcinoma (thyroid transcription factor-1 [TTF-1], napsin A) and squamous cell (p40, p63) carcinoma. This diagnosis cannot be made on non-resection or cytology specimens.
 - ▶ **Sarcomatoid carcinoma** is a general term that includes pleomorphic carcinoma, carcinosarcoma, and pulmonary blastoma. For this reason, it is best to use the specific term for these entities whenever possible rather than the general term.
 - ◊ **Pleomorphic carcinoma** is a poorly differentiated NSCC that contains at least 10% spindle and/or giant cells or a carcinoma consisting only of spindle and giant cells. Spindle cell carcinoma consists of an almost pure population of epithelial spindle cells, while Giant cell carcinoma consists almost entirely of tumor giant cells.
 - ◊ **Carcinosarcoma** is a malignant tumor that consists of a mixture of NSCC and sarcoma-containing heterologous elements (eg, rhabdomyosarcoma, chondrosarcoma, osteosarcoma).
 - ◊ **Pulmonary blastoma** is a biphasic tumor that consists of fetal adenocarcinoma (typically low grade) and primitive mesenchymal stroma.

[Continue](#)

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

**PRINCIPLES OF PATHOLOGIC REVIEW****Immunohistochemistry**

- **Judicious use of IHC is strongly recommended to preserve tissue for molecular testing, most notably in small specimens. When adenocarcinoma or squamous cell carcinomas are poorly differentiated, the defining morphologic criteria that would allow for specific diagnosis may be inconspicuous or absent. In this case, IHC or mucin staining may be necessary to determine a specific diagnosis.**
- **In small specimens, a limited number of immunostains with one lung adenocarcinoma marker (TTF-1, napsin A) and one squamous carcinoma marker (p40, p63) should suffice for most diagnostic problems. Virtually all tumors that lack squamous cell morphology and show co-expression of p63 and TTF-1 are preferably classified as adenocarcinoma. A simple panel of TTF-1 and p40 may be sufficient to classify most NSCC-NOS cases.**
- **Testing for nuclear protein in testis (NUT) expression by IHC should be considered in all poorly differentiated carcinomas that lack glandular differentiation or specific etiology, particularly in patients who do not smoke or in patients presenting at a young age, for consideration of a pulmonary NUT carcinoma.**
- **IHC should be used to differentiate primary lung adenocarcinoma from squamous cell carcinoma, large cell carcinoma, metastatic carcinoma, and primary pleural mesothelioma (particularly for pleural specimens).**
- **Primary pulmonary adenocarcinoma:**
 - ▶ **In patients for whom the primary origin of the carcinoma is uncertain, an appropriate panel of immunohistochemical stains is recommended to assess for metastatic carcinoma to the lung.**
 - ▶ **TTF-1 is a homeodomain-containing nuclear transcription protein of the *NKX2* gene family that is expressed in epithelial cells of the embryonal and mature lung and thyroid. TTF-1 immunoreactivity is seen in primary pulmonary adenocarcinoma in the majority (70%–90%) of non-mucinous adenocarcinoma subtypes. Metastatic adenocarcinoma to the lung is nearly always negative for TTF-1 except in metastatic thyroid malignancies, in which case thyroglobulin and PAX8 are also positive. Rare cases of TTF-1 positivity in tumors of other organs (gynecologic tract, pancreatobiliary) have been noted, and may be dependent on the specific TTF-1 clone utilized, stressing the importance of correlation with clinical and radiologic features.**
 - ▶ **Napsin A—an aspartic proteinase expressed in normal type II pneumocytes and in proximal and distal renal tubules—appears to be expressed in >80% of lung adenocarcinomas and may be a useful adjunct to TTF-1.**
 - ▶ **The panel of TTF-1 (or alternatively napsin A) and p40 (or alternatively p63) may be useful in refining the diagnosis to either adenocarcinoma or squamous cell carcinoma in small biopsy specimens previously classified as NSCC NOS.**

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

[Continue](#)



PRINCIPLES OF PATHOLOGIC REVIEW

Immunohistochemistry

- IHC should be used to confirm neuroendocrine differentiation when there is morphologic evidence of neuroendocrine morphology (eg, speckled chromatin pattern, nuclear molding, peripheral palisading):
 - ▶ NCAM (CD56), chromogranin, synaptophysin, and INSM1 are used to identify neuroendocrine tumors in cases in which morphologic suspicion of neuroendocrine differentiation exists.
 - ▶ A panel of markers is useful, but one positive marker is enough if the staining is unambiguous in more than 10% of the tumor cells.
- Pleural mesothelioma versus pulmonary adenocarcinoma
 - ▶ The distinction between pulmonary adenocarcinoma and pleural mesothelioma (epithelioid type) can be made by correlation of the histology with the clinical impression, imaging studies, and a panel of immunomarkers.
 - ▶ Immunostains sensitive and specific for pleural mesothelioma include WT-1, calretinin, CK5/6, and D2-40 (usually negative in adenocarcinoma).
 - ▶ Immunostains sensitive and specific for adenocarcinoma include pCEA, Claudin-4, TTF-1, and napsin A (negative in pleural mesothelioma). Other potentially useful markers that can be considered include B72.3, Ber-EP4, MOC-31, and CD15, but these generally do not have the sensitivity and specificity of the above markers.
 - ▶ A pancytokeratin such as AE1/AE3 is also useful, as a negative result suggests the possibility of other tumors.
 - ▶ Other markers can be helpful in the differential diagnosis between pleural mesothelioma and metastatic carcinoma, and will also help determine the tumor origin. Examples include markers for lung adenocarcinoma (TTF-1 and napsin A), breast carcinoma (ER α , PR, GCDPF-15, mammaglobin, and GATA-3), renal cell carcinoma (PAX8), papillary serous carcinoma (PAX8, PAX2, and ER), adenocarcinomas of the gastrointestinal tract (CDX2), and prostate cancer (NKX3.1). Additionally, p40 (or p63) is helpful for distinguishing epithelioid pleural mesotheliomas with pseudosquamous morphology from squamous cell carcinomas.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



PRINCIPLES OF SURGICAL THERAPY

Evaluation

- Determination of resectability, surgical staging, and *pulmonary resection should be performed by thoracic surgeons who perform lung cancer surgery as a prominent part of their practice.*
- CT and FDG-PET/CT used for staging should be within 60 days before proceeding with surgical evaluation.
- For medically operable disease, resection is the preferred local treatment modality (other modalities include SABR, thermal ablation such as radiofrequency ablation, and cryotherapy). Thoracic surgical oncology consultation should be part of the evaluation of any patient being considered for curative local therapy. In cases where SABR is considered for high-risk or borderline operable patients, a multidisciplinary evaluation including a radiation oncologist is recommended.
- The overall plan of treatment as well as needed imaging studies should be determined before any non-emergency treatment is initiated.
- Thoracic surgeons should actively participate in multidisciplinary discussions and meetings regarding patients with lung cancer (eg, multidisciplinary clinic and/or tumor board).
- Patients who actively smoke should be provided counseling and smoking cessation support ([NCCN Guidelines for Smoking Cessation](#)). While patients who actively smoke have a mildly increased incidence of postoperative pulmonary complications, these should not be considered a prohibitive risk for surgery. Surgeons should not deny surgery to patients solely due to smoking status, as surgery provides the predominant therapy for patients with early-stage lung cancer.

Resection ([NSCL-B 2 of 6](#))

Margins and Nodal Assessment ([NSCL-B 3 of 6](#))

^a Peripheral is defined as the outer one third of the lung parenchyma.

The Role of Surgery in Patients with N2 NSCLC
([NSCL-B 3 of 6](#) through [NSCL-B 5 of 6](#))

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



PRINCIPLES OF SURGICAL THERAPY

Resection

- Anatomic pulmonary resection is preferred for the majority of patients with NSCLC.
- Sublobar resection - Segmentectomy and wedge resection should be strongly considered for peripheral T1ab, N0 tumors.¹
- Sublobar resection should achieve parenchymal resection margins ≥ 2 cm or \geq the size of the nodule.
- Sublobar resection should also sample appropriate N1 and N2 lymph node stations unless not technically feasible without substantially increasing the surgical risk.
- Segmentectomy (preferred) or wedge resection is appropriate in selected patients with poor pulmonary reserve or other major comorbidity that contraindicates lobectomy.
- Minimally invasive surgery (VATS or robotic-assisted approaches) should be strongly considered for patients with no anatomic or surgical contraindications, as long as there is no compromise of standard oncologic and dissection principles of thoracic surgery. Robotic surgery should only be initiated by surgeons who have completed and maintained proficiency in the technique.
- In high-volume centers with significant VATS experience, VATS lobectomy in selected patients results in improved early outcomes (ie, decreased pain, reduced hospital length of stay, more rapid return to function, fewer complications) without compromise of cancer outcomes.
- Studies of robotic-assisted pulmonary resection show non-inferiority to traditional VATS approaches when performed by experienced robotic surgeons.^{2,3}
- Lung-sparing anatomic resection (sleeve lobectomy) is preferred over pneumonectomy, if anatomically appropriate and margin-negative resection is achieved.
- T3 (invasion) and T4 local extension tumors require en-bloc resection of the involved structure with negative margins. If a surgeon or center is uncertain about potential complete resection, consider obtaining an additional surgical opinion from a high-volume specialized center.

Evaluation ([NSCL-B 1 of 6](#))

Margins and Nodal Assessment ([NSCL-B 3 of 6](#))

The Role of Surgery in Patients with N2 NSCLC
([NSCL-B 3 of 6](#) through [NSCL-B 5 of 6](#))

[References](#)

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

**PRINCIPLES OF SURGICAL THERAPY****Margins and Nodal Assessment**

- Surgical pathologic correlation is critical to assess apparent close or positive margins, as these may not represent true margins or may not truly represent areas of risk for local recurrence (eg, medial surface of mainstem or bronchus intermedius when separate subcarinal lymph node dissection has been performed; pleural margin adjacent to aorta when no attachment to aorta is present).
- N1 and N2 node resection and mapping should be a routine component of lung cancer resections—a minimum of one N1 and three N2 stations sampled or complete lymph node dissection.
- Formal ipsilateral mediastinal lymph node dissection is indicated for patients undergoing resection for N2 disease.
- Complete resection requires free resection margins, systematic node dissection or sampling, and the highest mediastinal node negative for tumor. The resection is defined as incomplete whenever there is involvement of resection margins, unremoved positive lymph nodes, or positive pleural or pericardial effusions. A complete resection is referred to as R0, microscopically positive resection as R1, and macroscopic residual tumor as R2.
- Patients with clinical stage IB or greater, or high-risk factors, should be referred to medical oncology for evaluation.
- Consider referral to a radiation oncologist for N2 disease.

The Role of Surgery in Patients with N2 NSCLC

The role of surgery in patients with pathologically documented N2 disease remains controversial.⁴ Two randomized trials evaluated the role of surgery in this population, but neither showed an overall survival benefit with the use of surgery.^{5,6} However, this population is heterogeneous and the panel believes that these trials did not sufficiently evaluate the nuances present with the heterogeneity of N2 disease and the likely oncologic benefit of surgery in specific clinical situations.

- The presence or absence of N2 disease should be vigorously determined by both radiologic and invasive staging prior to the initiation of therapy since the presence of mediastinal nodal disease has a profound impact on prognosis and treatment decisions. ([NSCL-1](#), [NSCL-2](#), and [NSCL-6](#))
- Patients with occult-positive N2 nodes discovered at the time of pulmonary resection should continue with the planned resection along with formal mediastinal lymph node dissection. If N2 disease is noted in patients undergoing VATS, the surgeon may consider stopping the procedure so that induction therapy can be administered before surgery; however, continuing the procedure is also an option.
- The determination of the role of surgery in a patient with N2-positive lymph nodes should be made prior to the initiation of any therapy by a multidisciplinary team, including a thoracic surgeon who has a major part of his/her practice dedicated to thoracic oncology.⁷
- The presence of N2-positive lymph nodes substantially increases the likelihood of positive N3 lymph nodes. Pathologic evaluation of the mediastinum must include evaluation of the subcarinal station and contralateral lymph nodes. EBUS ± EUS are additional techniques for minimally invasive pathologic mediastinal staging that are complementary to mediastinoscopy. Even when these modalities are employed it is important to have an adequate evaluation of the number of stations involved and biopsy and documentation of negative contralateral lymph node involvement prior to a final treatment decision.

The Role of Surgery in Patients with N2 NSCLC is continued on [NSCL-B 4 of 6](#) through [NSCL-B 5 of 6](#)

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

References

**PRINCIPLES OF SURGICAL THERAPY****The Role of Surgery in Patients with N2 NSCLC**

- Repeat mediastinoscopy, while possible, is technically difficult and has a lower accuracy compared to primary mediastinoscopy. One possible strategy is to perform EBUS (\pm EUS) in the initial pretreatment evaluation and reserve mediastinoscopy for nodal restaging after neoadjuvant therapy.⁸
- Patients with a single lymph node smaller than 3 cm can be considered for a multimodality approach that includes surgical resection.^{4,9,10}
- Restaging after induction therapy is difficult to interpret, but CT \pm FDG-PET/CT should be performed to exclude disease progression or interval development of metastatic disease.
- Patients with negative mediastinum after neoadjuvant therapy have a better prognosis.^{10,11}
- Neoadjuvant chemoradiotherapy is used in one-third of the NCCN Member Institutions, while neoadjuvant chemotherapy is used in the other two-thirds. Overall survival appears similar provided RT is given postoperatively, if not given preoperatively.^{8,12} Neoadjuvant chemoradiotherapy is associated with higher rates of pathologic complete response and negative mediastinal lymph nodes.¹³ However, that is achieved at the expense of higher rates of acute toxicity and increased cost.
- When neoadjuvant chemoradiotherapy is used with doses lower than those used for standard definitive therapy, all efforts should be made to minimize any possible breaks in radiotherapy for surgical evaluation. Treatment breaks of more than 1 week are considered unacceptable.
- When timely surgical evaluation is not available, the strategy of neoadjuvant chemoradiotherapy should not be used. Another option in individual cases, and with the agreement of the thoracic surgeon, is to complete definitive chemoradiotherapy prior to re-evaluation and consideration for surgery.^{14,15} If a surgeon or center is uncertain about the feasibility or safety of resection after definitive doses of radiation, consider obtaining an additional surgical opinion from a high-volume specialized center. These operations may also benefit from additional considerations of soft tissue flap coverage in the radiation field at the time of resection.
- Data from a large multi-institutional trial indicate that pneumonectomy after neoadjuvant chemoradiotherapy has unacceptable morbidity and mortality.⁵ However, it is not clear if this is also true with neoadjuvant chemotherapy alone. Further, many groups have challenged that cooperative group finding with single-institution experiences demonstrating safety of pneumonectomy after induction therapy.¹⁶⁻¹⁹ In addition, there is no evidence that adding RT to induction regimens for patients with operable stage IIIA (N2) disease improves outcomes compared to induction chemotherapy.²⁰

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

References



PRINCIPLES OF SURGICAL THERAPY

The Role of Surgery in Patients with N2 NSCLC

A questionnaire was submitted to the NCCN Member Institutions in 2021 regarding their approach to patients with N2 disease. Their responses indicate the patterns of practice when approaching this difficult clinical problem.

- All NCCN Member Institutions treat select N2 patients with multimodality therapy that includes surgery.
- The majority of NCCN Member Institutions prefer EBUS for initial mediastinal staging, reserving mediastinoscopy for possible restaging.
- The majority of NCCN Member Institutions do not pathologically restage mediastinal lymph nodes after induction therapy and prior to surgery.
- All NCCN Member Institutions consider surgery for single-station non-bulky N2 disease.
- Approximately half of the institutions consider surgery for single-station bulky disease, 39% for multi-station non-bulky disease, and 21% for multi-station bulky disease.
- Two-thirds of NCCN Member Institutions prefer induction chemotherapy; one-third prefer chemoradiation.
- The majority require at least stable disease after induction, but do not require radiologic or pathologic response prior to surgery.
- Roughly a half would consider pneumonectomy after induction chemotherapy, but less than a quarter would consider pneumonectomy after chemoradiation.
- Approximately three-fourths would give adjuvant RT for positive residual N2 disease, but only approximately one-fourth would give RT for N2 pathologic complete response.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

**The Role of Surgery in Patients with N2 NSCLC – References**

- ¹ Altorki N, Wang X, Kozono D, et al. Lobar or sublobar resection for peripheral stage IA non-small-cell lung cancer. *N Engl J Med* 2023;388:489-498.
- ² Lampridis S, Maraschi A, Le Reun C, et al. Robotic versus video-assisted thoracic surgery for lung cancer: short-term outcomes of a propensity matched analysis. *Cancer (Basel)* 2023;15:2391.
- ³ Ma J, Li X, Zhao S, et al. Robot-assisted thoracic surgery versus video-assisted thoracic surgery for lung lobectomy or segmentectomy in patients with non-small cell lung cancer: a meta-analysis. *BMC Cancer* 2021;21:498.
- ⁴ Martins RG, D'Amico TA, Loo BW Jr, et al. The management of patients with stage IIIA non-small cell lung cancer with N2 mediastinal node involvement. *J Natl Compr Canc Netw* 2012;10:599-613.
- ⁵ Albain K, Swann RS, Rusch VW, et al. Radiotherapy plus chemotherapy with or without surgical resection for stage III non-small-cell lung cancer: a phase III randomized controlled trial. *Lancet* 2009;374:379-386.
- ⁶ van Meerbeeck JP, Kramer GW, Van Schil PE, et al. Randomized controlled trial of resection versus radiotherapy after induction chemotherapy in stage IIIA-N2 non-small-cell lung cancer. *J Natl Cancer Inst* 2007;99:442-450.
- ⁷ Farjah F, Flum DR, Varghese TK Jr, et al. Surgeon specialty and long-term survival after pulmonary resection for lung cancer. *Ann Thorac Surg* 2009;87:995-1006.
- ⁸ Thomas M, Rube C, Hoffknecht P, et al. Effect of preoperative chemoradiation in addition to preoperative chemotherapy: a randomised trial in stage III non-small-cell lung cancer. *Lancet Oncol* 2008;9:636-648.
- ⁹ Andre F, Grunenwald D, Pignon J, et al. Survival of patients with resected N2 non-small-cell lung Cancer: Evidence for a subclassification and implications. *J Clin Oncol* 2000;18:2981-2989.
- ¹⁰ Decaluwé H, De Leyn P, Vansteenkiste J, et al. Surgical multimodality treatment for baseline resectable stage IIIA-N2 non-small cell lung cancer. Degree of mediastinal lymph node involvement and impact on survival. *Eur J Cardiothorac Surg* 2009;36:433-439.
- ¹¹ Bueno R, Richards WG, Swanson SJ, et al. Nodal stage after induction therapy for stage IIIA lung cancer determines patient survival. *Ann Thorac Surg* 2000;70:1826-1831.
- ¹² Higgins K, Chino JP, Marks LB, et al. Preoperative chemotherapy versus preoperative chemoradiotherapy for stage III (N2) non-small-cell lung cancer. *Int J Radiat Oncol Biol Phys* 2009;75:1462-1467.
- ¹³ de Cabanyes Candela S, Detterbeck FC. A systematic review of restaging after induction therapy for stage IIIa lung cancer: prediction of pathologic stage. *J Thorac Oncol* 2010;5:389-398.
- ¹⁴ Bauman JE, Mulligan MS, Martins RG, et al. Salvage lung resection after definitive radiation (>59 Gy) for non-small cell lung cancer: surgical and oncologic outcomes. *Ann Thorac Surg* 2008;86:1632-1638.
- ¹⁵ Sonett JR, Suntharalingam M, Edelman MJ, et al. Pulmonary resection after curative intent radiotherapy (>59 Gy) and concurrent chemotherapy in non-small-cell lung cancer. *Ann Thorac Surg* 2004;78:1200-1205.
- ¹⁶ Evans NR 3rd, Li S, Wright CD, et al. The impact of induction therapy on morbidity and operative mortality after resection of primary lung cancer. *J Thorac Cardiovasc Surg* 2010;139:991-996.
- ¹⁷ Gaissert HA, Keum DY, Wright CD, et al. POINT: Operative risk of pneumonectomy—Influence of preoperative induction therapy. *J Thorac Cardiovasc Surg* 2009;138:289-294.
- ¹⁸ Mansour Z, Kochetkova EA, Ducrocq X, et al. Induction chemotherapy does not increase the operative risk of pneumonectomy! *Eur J Cardiothorac Surg* 2007;31:181-185.
- ¹⁹ Weder W, Collaud S, Eberhardt WE, et al. Pneumonectomy is a valuable treatment option after neoadjuvant therapy for stage III non-small-cell lung cancer. *J Thorac Cardiovasc Surg* 2010;139:1424-1430.
- ²⁰ Shah AA, Berry MF, Tzao C, et al. Induction chemoradiotherapy is not superior to induction chemotherapy alone in stage IIIA lung cancer: a systematic review and meta-analysis. *Ann Thorac Surg* 2012;93:1807-1812.

Note: All recommendations are category 2A unless otherwise indicated.**Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.**

**PRINCIPLES OF RADIATION THERAPY****I. General Principles** ([Table 1. Commonly Used Abbreviations in Radiation Therapy](#))

- Determination of the appropriateness of radiation therapy (RT) should be made by radiation oncologists who perform lung cancer RT as a prominent part of their practice.
- RT has a potential role in all stages of NSCLC, as either definitive/consolidative or palliative therapy. Radiation oncology input as part of a multidisciplinary evaluation or discussion should be provided for all patients with stage III NSCLC, with early-stage disease who are medically inoperable, who refuse surgery, or who are high-risk surgical candidates, and with stage IV disease that may benefit from local therapy.
- The critical goals of modern RT are to maximize tumor control and to minimize treatment toxicity. A minimum technologic standard is CT-planned 3D-CRT.¹
- More advanced technologies are appropriate when needed to deliver curative RT safely. These technologies include (but are not limited to) 4D-CT, FDG-PET/CT and/or MRI simulation, IMRT/VMAT, IGRT, motion management, and proton therapy (<https://www.astro.org/Daily-Practice/Reimbursement/Model-Policies/Model-Policies>). Nonrandomized comparisons of using advanced technologies demonstrate reduced toxicity and improved survival versus older techniques.²⁻⁴ In a prospective trial of definitive/consolidative chemotherapy/RT for patients with stage III NSCLC (RTOG 0617), IMRT was associated with a nearly 60% decrease (from 7.9% to 3.5%) in high-grade radiation pneumonitis as well as similar survival and tumor control outcomes despite a higher proportion of stage IIIB and larger treatment volumes compared to 3D-CRT;⁵ as such, IMRT is preferred over 3D-CRT in this setting. In retrospective studies, intensity-modulated proton therapy (IMPT) has also been shown to reduce the toxicities as compared with 3D-based passive scattering proton therapy in stage III NSCLC.³
- Highly conformal RT, such as IMRT or proton therapy, should be used in the setting of prior RT, potentially with hyperfractionation, to reduce risk of toxicity.
- Centers using advanced technologies should implement and document modality-specific quality assurance measures. The ideal is external credentialing of both treatment planning and delivery such as required for participation in RTOG clinical trials employing advanced technologies. Useful references include the ACR Practice Parameters and Technical Standards (<https://www.acr.org/~media/ACR/Documents/PGTS/toc.pdf>).
- The interaction of strong vascular endothelial growth factor (VEGF) inhibitors with prior or subsequent dose-intensive RT (SABR or definitive dose accelerated fractionation) involving the proximal bronchial tree, hilar vessels, or esophagus can lead to serious toxicity. Careful coordination of medical and radiation oncology on the therapeutic strategy is important, including the choice and sequencing of systemic agents with strong VEGF inhibitors and the dose and fractionation of radiation, especially for patients with metastatic disease.

II. Radiation Therapy Simulation, Planning, and Delivery

- Simulation should be performed using CT scans obtained in the RT treatment position with appropriate immobilization devices. IV contrast with or without oral contrast is recommended for better target/organ delineation whenever possible in patients with central tumors or nodal disease. Because IV contrast can affect tissue heterogeneity correction calculations, density masking or use of a pre-contrast scan may be needed when intense enhancement is present.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

[Continued](#)**NSCL-C**
1 OF 11

**PRINCIPLES OF RADIATION THERAPY****II. Radiation Therapy Simulation, Planning, and Delivery (continued)**

- FDG-PET/CT significantly improves targeting accuracy,⁶ especially for patients with significant atelectasis and when IV CT contrast is contraindicated. A randomized trial of FDG-PET/CT versus CT-only RT planning demonstrated improved preemption of futile radical RT, decreased recurrences, and a trend toward improved overall survival with FDG-PET/CT RT planning.⁷ Given the potential for rapid progression of NSCLC,^{8,9} FDG-PET/CT should be obtained preferably within 4 weeks before treatment. It is ideal to obtain FDG-PET/CT in the treatment position.
- Tumor and organ motion, especially owing to breathing, should be assessed or accounted for at simulation. Options include fluoroscopy, inhale/exhale or slow scan CT, or, ideally, 4D-CT.
- Photon beam energy should be individualized based on the anatomic location of the tumors and beam paths. In general, photon energies between 4 to 10 MV are recommended for beams passing through low-density lung tissue before entering the tumor. When there is no air gap before the beam enters the tumor (such as for some large mediastinal tumors or tumors attached to the chest wall), higher energies may improve the dose distribution, especially when using a smaller number of fixed beam angles.
- Tissue heterogeneity correction and accurate dose calculation algorithms are recommended that account for buildup and lateral electron scatter effects in heterogeneous density tissues. Heterogeneity correction with simple pencil beam algorithms is not recommended.¹⁰
- Respiratory motion should be managed when motion is excessive. This includes (but is not limited to) forced shallow breathing with abdominal compression, accelerator beam gating with the respiratory cycle, dynamic tumor tracking, active breathing control (ABC), or coaching/biofeedback techniques. If motion is minimal or the ITV is small, motion-encompassing targeting is appropriate. A useful resource for implementation of respiratory motion management is the report of AAPM Task Group 76.¹¹
- IGRT—including (but not limited to) orthogonal pair planar imaging and/or volumetric imaging (such as CBCT, CT on rails, or MRI)—is recommended when using SABR, 3D-CRT/IMRT, and proton therapy with steep dose gradients around the target, when OARs are in close proximity to high-dose regions, and when using complex motion management techniques.

III. Target Volumes, Prescription Doses, and Normal Tissue Dose Constraints (See Tables 2–5 on [NSCL-C 7 of 10](#) and [NSCL-C 8 of 10](#))

- ICRU Reports 62 and 83 detail the current definitions of target volumes for 3D-RT and IMRT. GTV comprises the known extent of disease (primary and nodal) on imaging and pathologic assessment, CTV includes regions of presumed microscopic extent or dissemination, and PTV comprises the ITV (which includes margin for target motion) plus a setup margin for positioning and mechanical variability.
<https://www.nrgoncology.org/ciro-lung>
- PTV margin can be decreased by immobilization, motion management, and IGRT techniques.
- Consistent delineation of normal structures is critical for evaluating plans for safety. The RTOG consensus lung-contouring atlas is a useful resource. <https://www.nrgoncology.org/ciro-lung>
- Commonly used prescription doses and normal tissue dose constraints are summarized in Tables 2 through 5. These are based on published experience, ongoing trials, historical data, modeling, and empirical judgment.^{12,13} Useful references include the recent reviews of normal organ dose responses from the QUANTEC project.¹⁴⁻¹⁸ Because risk of normal organ toxicity increases with dose, doses to normal organs should be kept as low as reasonably achievable rather than simply meeting nominal constraints. This is generally facilitated by more advanced techniques to achieve better dose conformity.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

[Continued](#)

**PRINCIPLES OF RADIATION THERAPY****IV. General Treatment Information****Early-Stage NSCLC (Stage I, selected node-negative Stage IIA)**

- SABR (also known as SBRT)¹⁹ has achieved good primary tumor control rates and overall survival, higher than conventionally fractionated radiotherapy. Although SABR is not proven equivalent to lobectomy, some prospective series have demonstrated similar overall and cancer-specific survival with reduced acute toxicity.²⁰⁻³¹
- SABR is also an appropriate option for patients with high surgical risk (able to tolerate sublobar resection but not lobectomy [eg, age ≥75 years, poor lung or cardiac function]).
- More modestly hypofractionated or dose-intensified conventionally fractionated highly conformal radiation (IMRT with IGRT preferred) are less preferred alternatives and may be considered if referral for SABR is not feasible.³²⁻³⁴
- In patients treated with surgery, postoperative radiotherapy (PORT) is not recommended unless there are positive margins (see *Locally Advanced NSCLC* in this section for patients upstaged to N2).
- Close follow-up and therapy for isolated local and/or locoregional recurrence after SABR have been shown to improve overall survival in a large retrospective study.³⁵

SABR for Node-Negative Early-Stage NSCLC

- The high-dose intensity and conformity of SABR require minimizing the PTV.
- Dosing regimen
 - For SABR, intensive regimens of BED ≥100 Gy are associated with significantly better local control and survival than less intensive regimens.^{36,37} In the United States, only regimens of ≤5 fractions meet the arbitrary billing code definition of SBRT, but slightly more protracted regimens are appropriate as well.^{36,38} For centrally located tumors (defined variably as within 2 cm of the proximal bronchial tree and/or abutting mediastinal pleura) and even ultra-central tumors (defined as abutting the proximal bronchial tree or, in some definitions, other critical mediastinal structures as well), 4 to 10 fraction risk-adapted SABR regimens appear to be effective and safe,³⁹⁻⁴² while 54 to 60 Gy in 3 fractions is unsafe and should be avoided.⁴³ However, particular attention should be paid to tumors abutting the bronchial tree and esophagus to avoid severe toxicity. RTOG 0813 evaluated the toxicity of 5-fraction regimens and found no high-grade toxicities at 50 Gy in 5 fractions.⁴⁴
- SABR is most commonly used for tumors up to 5 cm in size, though selected larger isolated tumors can be treated safely if normal tissue constraints are respected.^{44,45}
- Prescription doses incompletely describe the actual delivered doses, which also strongly depend on how the dose is prescribed (to the isocenter vs. an isodose volume covering a proportion of the PTV), the degree of dose heterogeneity, whether tissue density heterogeneity corrections are used, and the type of dose calculation algorithm.^{10,46,47} All of these must be considered when interpreting or emulating regimens from prior studies.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

[Continued](#)**NSCL-C**
3 OF 11

**PRINCIPLES OF RADIATION THERAPY****Locally Advanced NSCLC (Stage II–III)**

- Concurrent chemotherapy/RT is recommended for patients with inoperable stage II (node-positive) and stage III NSCLC.⁴⁸⁻⁵¹
- RT interruptions and dose reductions for manageable acute toxicities should be avoided by employing supportive care.
- Sequential chemotherapy/RT or RT alone is appropriate for frail patients unable to tolerate concurrent therapy.^{52,53}
Accelerated RT regimens may be beneficial, particularly if concurrent chemotherapy would not be tolerated (ie, in a sequential or RT-only approach).^{54,55}
- Preoperative systemic therapy and postoperative RT is an option for patients with resectable N2 NSCLC (minimal N2 and treatable with lobectomy).^{56,57}
- Preoperative concurrent chemotherapy/RT is an alternative option for patients with resectable N2 NSCLC and is recommended for resectable superior sulcus tumors.^{58,59} RT should be planned up front such that it continues to a definitive dose without interruption if the patient does not proceed to surgery as initially planned.
- The optimal timing of RT in trimodality therapy (preoperative with chemotherapy or postoperative) is not established and is controversial.^{60,61}
- The determination of resectability in trimodality therapy should be made prior to initiation of all treatment. Upfront multidisciplinary consultation is particularly important when considering surgical treatment of patients with stage III NSCLC.
- In patients with clinical stage I/II upstaged surgically to N2 with completely resected disease, two randomized studies did not show an overall survival benefit of PORT, although locoregional control was significantly improved.^{62,63} PORT (generally following postoperative chemotherapy) may be considered for selected patients with high-risk N2 disease, such as extracapsular extension, multi-station involvement, inadequate lymph node dissection/sampling, and/or refusal or intolerance of adjuvant systemic therapy. To minimize potential lung and heart toxicities, highly conformal RT techniques such as IMRT or proton therapy are preferred.⁶⁴⁻⁶⁷
- In patients with completely resected pN1 receiving adjuvant systemic therapy, PORT is not recommended. PORT may be considered for these patients if they are unable to receive adjuvant systemic therapy.⁶¹

Conventionally Fractionated RT for Locally Advanced NSCLC

- IFI omitting ENI allows tumor dose escalation and is associated with a low risk of isolated nodal relapse, particularly in a patient staged with FDG-PET/CT.⁶⁸⁻⁷² Three randomized trials found improved survival for IFI versus ENI, possibly because it enabled dose escalation.⁷³⁻⁷⁵ IFI is reasonable in order to optimize definitive dosing to the tumor and/or decrease normal tissue toxicity.^{74,75}
- Dosing Regimens
 - ▶ The most commonly prescribed doses for definitive RT are 60 to 70 Gy in 2 Gy fractions. Doses of at least 60 Gy should be given.⁷⁶ Dose escalation is associated with better survival in non-randomized comparisons in RT alone,⁷⁷ sequential chemotherapy/RT,⁷⁸ or concurrent chemotherapy/RT.⁷⁹ While optimal RT dose intensification remains a valid question, a high dose of 74 Gy is not currently recommended for routine use.⁸⁰⁻⁸⁵
A meta-analysis demonstrated improved survival with accelerated fractionation RT regimens,⁸⁶ and RTOG 1106 found that PET-based individualized accelerated RT dose intensification potentially improved local control but not overall survival.⁸⁷

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

[Continued](#)

**PRINCIPLES OF RADIATION THERAPY****Conventionally Fractionated RT for Locally Advanced NSCLC (continued)****• Dosing Regimens**

- ▶ Doses of 45 to 54 Gy in 1.8 to 2 Gy fractions are standard preoperative doses.⁸⁸ Definitive RT doses delivered as preoperative chemotherapy/RT can safely be administered and achieve promising nodal clearance and survival rates,⁸⁹⁻⁹² but require experience in thoracic surgical techniques to minimize the risk of surgical complications after high-dose RT.
- ▶ In PORT, the CTV includes the bronchial stump and high-risk draining lymph node stations.⁹³ Standard doses after complete resection are 50 to 54 Gy in 1.8 to 2 Gy fractions, but a boost may be administered to high-risk regions including areas of nodal extracapsular extension or microscopic positive margins.⁹⁴⁻⁹⁶ Lung dose constraints should be more conservative, because tolerance appears to be reduced after surgery. The LungART and PORT-C trials provide useful guidelines for PORT technique.⁹⁷ Highly conformal techniques to minimize lung and heart dose are preferred.

Advanced/Metastatic NSCLC (Stage IV)

- RT is recommended for local palliation or prevention of symptoms (such as pain, bleeding, or obstruction).
- Definitive/consolidative local therapy to isolated or limited metastatic sites (oligometastases) (including but not limited to brain, lung, and adrenal gland) achieves prolonged survival in a small proportion of well-selected patients with good performance status who have also received radical therapy to the intrathoracic disease.⁹⁸ Definitive RT to oligometastases (limited number is not universally defined but clinical trials have included 3–5 metastases), particularly SABR, is an appropriate option in such cases if it can be delivered safely to the involved sites.^{99,100} In two randomized phase II trials, significantly improved progression-free survival and overall survival in one trial^{101,102} were found for local consolidative therapy (RT or surgery) to oligometastatic lesions versus maintenance systemic therapy or observation for patients not progressing on systemic therapy.¹⁰¹⁻¹⁰³
- In the setting of progression at a limited number of sites on a given line of systemic therapy (oligoprogression), local ablative therapy to the oligoprogressive sites may extend the duration of benefit of the current line of systemic therapy.
- When treating oligometastatic/oligoprogressive lesions, if SABR is not feasible, other dose-intensive accelerated/hypofractionated CRT regimens may be used.
- See the [NCCN Guidelines for Central Nervous System Cancers](#) regarding RT for brain metastases.
- A pooled analysis of two randomized trials indicated that adding radiotherapy to a certain immune checkpoint inhibitor (anti-PD-1) significantly increased responses and clinical outcomes in patients with metastatic NSCLC.¹⁰⁴ Larger phase III randomized studies are ongoing.

Palliative RT for Advanced/Metastatic NSCLC

- The dose and fractionation of palliative RT should be individualized based on goals of care, symptoms, performance status, and logistical considerations. Shorter courses of RT are preferred for patients with poor performance status and/or shorter life expectancy because they provide similar pain relief as longer courses, although there is a higher potential need for retreatment.¹⁰⁵⁻¹⁰⁸ For palliation of thoracic symptoms, higher dose/longer-course thoracic RT (eg, ≥30 Gy in 10 fractions) is associated with modestly improved survival and symptoms, particularly in patients with good performance status.^{109,110} When higher doses (>30 Gy) are warranted, technologies to reduce normal tissue irradiation (at least 3D-CRT and including IMRT or proton therapy as appropriate) may be used.
- Single-fraction SRT of 12–16 Gy produced better control of pain response and local control of non-spine bone metastases compared to standard 30 Gy in 10 fractions in a randomized phase II trial, and may be promising for patients with longer expected survival.¹¹¹ SABR/SRS has been found in randomized clinical trials to produce better pain and tumor control of spine and non-spine bone metastases than conventionally fractionated palliative RT, and is appropriate especially for patients with longer expected survival.¹¹²

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

[Continued](#)**NSCL-C**
5 OF 11



PRINCIPLES OF RADIATION THERAPY

Table 1. Commonly Used Abbreviations in Radiation Therapy

RT	Radiation Therapy or Radiotherapy	ICRU	International Commission on Radiation Units and Measurements
2D-RT	2-Dimensional RT	IFI	Involved Field Irradiation
3D-CRT	3-Dimensional Conformal RT	IGRT	Image-Guided RT
4D-CT	4-Dimensional Computed Tomography	IMRT	Intensity-Modulated RT
AAPM	American Association of Physicists in Medicine	ITV*	Internal Target Volume
ABC	Active Breathing Control	OAR	Organ at Risk
ACR	American College of Radiology	OBI	On-Board Imaging
ASTRO	American Society for Radiation Oncology	PORT	Postoperative RT
BED	Biologically Effective Dose	PTV*	Planning Target Volume
CBCT	Cone-Beam CT	QUANTEC	Quantitative Analysis of Normal Tissue Effects in the Clinic
CTV*	Clinical Target Volume	RTOG	Radiation Therapy Oncology Group now part of NRG Oncology
ENI	Elective Nodal Irradiation	SABR	Stereotactic Ablative RT, also known as Stereotactic Body RT (SBRT)
GTV*	Gross Tumor Volume	VMAT	Volumetric Modulated Arc Therapy

***Refer to ICRU Report 83 for detailed definitions.**

Note: All recommendations are category 2A unless otherwise indicated.
Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

PRINCIPLES OF RADIATION THERAPY

Please note: Tables 2–5 provide doses and constraints used commonly or in past clinical trials as useful references rather than specific recommendations.

Table 2. Commonly Used Doses for SABR

Total Dose	# Fractions	Example Indications
25–34 Gy	1	Peripheral, small
45–60 Gy	3	Peripheral tumors
48–50 Gy	4	Central or peripheral tumors <4–5 cm
50–55 Gy	5	Central tumors
50–60 Gy	5	Peripheral tumors
60–70 Gy	8–10	Central tumors

Table 3. Maximum Dose Constraints for SABR*

OAR/Regimen	1 Fraction	3 Fractions	4 Fractions	5 Fractions
Spinal cord	14 Gy	18 Gy (6 Gy/fx)	26 Gy (6.5 Gy/fx)	30 Gy (6 Gy/fx)
Esophagus	15.4 Gy	27 Gy (9 Gy/fx)	30 Gy (7.5 Gy/fx)	105% of PTV prescription^
Brachial plexus	17.5 Gy	24 Gy (8 Gy/fx)	27.2 Gy (6.8 Gy/fx)	32 Gy (6.4 Gy/fx)
Heart/ pericardium	22 Gy	30 Gy (10 Gy/fx)	34 Gy (8.5 Gy/fx)	105% of PTV prescription^
Great vessels	37 Gy	NS	49 Gy (12.25 Gy/fx)	105% of PTV prescription^
Trachea & proximal bronchi	20.2 Gy	30 Gy (10 Gy/fx)	34.8 Gy (8.7 Gy/fx)	105% of PTV prescription^
Rib	30 Gy	30 Gy (10 Gy/fx)	40 Gy (10 Gy/fx)	NS
Skin	26 Gy	24 Gy (8 Gy/fx)	36 Gy (9 Gy/fx)	32 Gy (6.4 Gy/fx)
Stomach	12.4 Gy	NS	27.2 Gy (6.8 Gy/fx)	NS

*Based on constraints used in recent RTOG SABR trials (RTOG 0618, 0813, & 0915).

^For central tumor location. NS = not specified.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

[Continued](#)

NSCL-C
7 OF 11



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

PRINCIPLES OF RADIATION THERAPY

Please note: Tables 2–5 provide doses and constraints used commonly or in past clinical trials as useful references rather than specific recommendations.

Table 4. Commonly Used Doses for Conventionally Fractionated and Palliative RT

Treatment Type	Total Dose	Fraction Size	Treatment Duration
Definitive RT with or without chemotherapy	60–70 Gy	2 Gy	6–7 weeks
Preoperative RT	45–54 Gy	1.8–2 Gy	5 weeks
Postoperative RT • Negative margins • Extracapsular nodal extension or microscopic positive margins • Gross residual tumor	50–54 Gy	1.8–2 Gy	5–6 weeks
	54–60 Gy	1.8–2 Gy	6 weeks
	60–70 Gy	2 Gy	6–7 weeks
Palliative RT • Obstructive disease (SVC syndrome or obstructive pneumonia) • Bone metastases with soft tissue mass • Bone metastases without soft tissue mass • Brain metastases • Symptomatic chest disease in patients with poor PS • Any metastasis in patients with poor PS	30–45 Gy	3 Gy	2–3 weeks
	20–30 Gy	4–3 Gy	1–2 weeks
	8–30 Gy	8–3 Gy	1 day–2 weeks
	CNS GLs* 17 Gy**	CNS GLs* 8.5 Gy**	CNS GLs* 1–2 weeks**
	8–20 Gy	8–4 Gy	1 day–1 week

* [NCCN Guidelines for Central Nervous System Cancers](#).

** This regimen includes one dose per week, as the phase 3 study included day 1 & 8 treatments.

Table 5. Normal Tissue Dose-Volume Constraints for Conventionally Fractionated RT with Concurrent Chemotherapy^{†,‡}

OAR	Constraints in 30–35 fractions
Spinal cord	Max ≤50 Gy
Lung	V20 ≤35%–40%; [§] MLD ≤20 Gy
Heart	V50 ≤25%; Mean ≤20 Gy
Esophagus	Mean ≤34 Gy; Max ≤105% of prescription dose; V60 ≤17%; contralateral sparing is desirable
Brachial plexus	Median dose ≤69 Gy

Vxx = % of the whole OAR receiving ≥xx Gy.

[†]These constraints represent doses that generally should not be exceeded, based on a consensus survey of NCCN Member Institutions. Because the risk of toxicity increases progressively with dose to normal tissues, a key principle of radiation treatment planning is to keep normal tissue doses "as low as reasonably achievable" while adequately covering the target. The doses to any given OAR should typically be lower than these constraints, approaching them only when there is close proximity to the target volume.

[‡]Speirs CK, et al. J Thorac Oncol 2017;12:293-301; Wang K, et al. J Clin Oncol 2017;35:1387-1394; Amini A, et al. Int J Radiat Oncol Biol Phys 2012;82:e391-398; Graham MV, et al. Int J Radiat Oncol Biol Phys 1999;45:323-329; Palma DA, et al. Int J Radiat Oncol Biol Phys 2013;85:444-450; Kamran SC, et al. JAMA Oncol 2021;7:910-914.

[§] Use V20 <35%, especially for the following: patients ≥70 years, taxane chemotherapy, and poor PFTs (such as FEV1 or DLCO <50% normal). Use more conservative limits with a diagnosis or radiologic evidence of idiopathic pulmonary fibrosis (IPF)/usual interstitial pneumonia (UIP) (the tolerance of these patients is lower though not well characterized).

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

References

NSCL-C
8 OF 11



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

- 1 Chen AB, et al. Survival outcomes after radiation therapy for stage III non-small-cell lung cancer after adoption of computed tomography-based simulation. *J Clin Oncol* 2011;29:2305-2311.
- 2 Liao ZX, et al. Influence of technologic advances on outcomes in patients with unresectable, locally advanced non-small-cell lung cancer receiving concomitant chemoradiotherapy. *Int J Radiat Oncol Biol Phys* 2010;76:775-781.
- 3 Gijshi O, et al. Toxicity and survival after intensity-modulated proton therapy versus passive scattering proton therapy for NSCLC. *J Thorac Oncol* 2021;16:269-277.
- 4 Chang JY, et al. Proton beam radiotherapy and concurrent chemotherapy for unresectable stage III non-small cell lung cancer: final results of a phase 2 study. *JAMA Oncol* 2017;3:e172032.
- 5 Chun SG, et al. Impact of intensity-modulated radiation therapy technique for locally advanced non-small cell lung cancer: a secondary analysis of the NRG Oncology RTOG 0617 randomized clinical trial. *J Clin Oncol* 2017;35:56-62.
- 6 MacManus M, et al. Use of PET and PET/CT for radiation therapy planning: IAEA expert report 2006-2007. *Radiother Oncol* 2009;91:85-94.
- 7 Ung YC, et al. An Ontario Clinical Oncology Group (OCOG) randomized trial (PET START) of FDG PET/CT in patients with stage 3 non-small cell lung cancer (NSCLC): impact of PET on radiation treatment volumes [abstract]. *J Thorac Oncol* 2011;6: S428.
- 8 Everitt S, et al. High rates of tumor growth and disease progression detected on serial pretreatment positrooxyglucose-positron emission tomography/computed tomography scans in radical radiotherapy candidates with nonsmall cell lung cancer. *Cancer* 2010;116:5030-5037.
- 9 Mohammed N, et al. Rapid disease progression with delay in treatment of non-small-cell lung cancer. *Int J Radiat Oncol Biol Phys* 2011;79:466-472.
- 10 Liu MB, et al. Clinical impact of dose overestimation by effective path length calculation in stereotactic ablative radiation therapy of lung tumors. *Pract Radiat Oncol* 2013;3:294-300.
- 11 Keall PJ, et al. The management of respiratory motion in radiation oncology report of AAPM Task Group 76. *Med Phys* 2006;33:3874-3900.
- 12 Kong FM, et al. Physical models and simpler dosimetric descriptors of radiation late toxicity. *Semin Radiat Oncol* 2007;17:108-120.
- 13 Timmerman RD. An overview of hypofractionation and introduction to this issue of seminars in radiation oncology. *Semin Radiat Oncol* 2008;18:215-222.
- 14 Marks LB, et al. Use of normal tissue complication probability models in the clinic. *Int J Radiat Oncol Biol Phys* 2010;76:S10-19.
- 15 Marks LB, et al. Radiation dose-volume effects in the lung. *Int J Radiat Oncol Biol Phys* 2010;76:S70-76.
- 16 Werner-Wasik M, et al. Radiation dose-volume effects in the esophagus. *Int J Radiat Oncol Biol Phys* 2010;76:S86-93.
- 17 Gagliardi G, et al. Radiation dose-volume effects in the heart. *Int J Radiat Oncol Biol Phys* 2010;76:S77-85.
- 18 Kirkpatrick JP, et al. Radiation dose-volume effects in the spinal cord. *Int J Radiat Oncol Biol Phys* 2010;76:S42-49.
- 19 Videtec GMM, et al. Stereotactic body radiation therapy for early-stage non-small cell lung cancer: executive summary of an ASTRO evidence-based guideline. *Pract Radiat Oncol* 2017;7:295-301.
- 20 Timmerman R, et al. Stereotactic body radiation therapy for inoperable early stage lung cancer. *JAMA* 2010;303:1070-1076.
- 21 Baumann P, et al. Outcome in a prospective phase II trial of medically inoperable stage I non-small-cell lung cancer patients treated with stereotactic body radiotherapy. *J Clin Oncol* 2009;27:3290-3296.
- 22 Onishi H, et al. Stereotactic body radiotherapy (SBRT) for operable stage I non-small-cell lung cancer: can SBRT be comparable to surgery? *Int J Radiat Oncol Biol Phys* 2011;81:1352-1358.
- 23 Grutters JPC, et al. Comparison of the effectiveness of radiotherapy with photons, protons and carbon-ions for non-small cell lung cancer: a meta-analysis. *Radiother Oncol* 2010;95:32-40.
- 24 Palma D, et al. Impact of introducing stereotactic lung radiotherapy for elderly patients with stage I non-small-cell lung cancer: a population-based time-trend analysis. *J Clin Oncol* 2010;28:5153-5159.
- 25 Shirvani SM, et al. Comparative effectiveness of 5 treatment strategies for early-stage non-small cell lung cancer in the elderly. *Int J Radiat Oncol Biol Phys* 2012;84:1060-1070.
- 26 Sun B, et al. 7-year follow-up after stereotactic ablative radiotherapy for patients with stage I non-small cell lung cancer: Results of a phase 2 clinical trial. *Cancer* 2017;123:3031-3039.
- 27 Grills IS, et al. Outcomes after stereotactic lung radiotherapy or wedge resection for stage I non-small-cell lung cancer. *J Clin Oncol* 2010;28:928-935.
- 28 Crabtree TD, et al. Stereotactic body radiation therapy versus surgical resection for stage I non-small cell lung cancer. *J Thorac Cardiovasc Surg* 2010;140:377-386.
- 29 Chang JY, et al. Stereotactic ablative radiotherapy versus lobectomy for operable stage I non-small-cell lung cancer: a pooled analysis of two randomised trials. *Lancet Oncol* 2015;16:630-637.
- 30 Chang JY, et al. Stereotactic ablative radiotherapy for operable stage I non-small-cell lung cancer (revised STARS): long-term results of a single-arm, prospective trial with prespecified comparison to surgery. *Lancet Oncol* 2021;22:1448-1457.
- 31 Ball D, et al. Stereotactic ablative radiotherapy versus standard radiotherapy in stage 1 non-small-cell lung cancer (TROG 09.02 CHISEL): a phase 3, open-label, randomised controlled trial. *Lancet Oncol* 2019;20:494-503.
- 32 Bogart JA, et al. Phase I study of accelerated conformal radiotherapy for stage I non-small-cell lung cancer in patients with pulmonary dysfunction: CALGB 39904. *J Clin Oncol* 2010;28:202-206.
- 33 Zhao L, et al. High radiation dose may reduce the negative effect of large gross tumor volume in patients with medically inoperable early-stage non-small cell lung cancer. *Int J Radiat Oncol Biol Phys* 2007;68:103-110.
- 34 Cheung P, et al. Phase II study of accelerated hypofractionated three-dimensional conformal radiotherapy for stage T1-3 N0 M0 non-small cell lung cancer: NCIC CTG BR.25. *J Natl Cancer Inst* 2014;106:1-8.
- 35 Brooks ED, et al. Association of long-term outcomes and survival with multidisciplinary salvage treatment for local and regional recurrence after stereotactic ablative radiotherapy for early-stage lung cancer. *JAMA Netw Open* 2018;1:e181390.
- 36 Onishi H, et al. Hypofractionated stereotactic radiotherapy (HypoFXSRT) for stage I non-small cell lung cancer: updated results of 257 patients in a Japanese multi-institutional study. *J Thorac Oncol* 2007;2:S94-100.
- 37 Moreno AC, et al. Biologically effective dose in stereotactic body radiotherapy and survival for patients with early-stage NSCLC. *J Thorac Oncol* 2020;15:101-109.
- 38 Lagerwaard FJ, et al. Outcomes of risk-adapted fractionated stereotactic radiotherapy for stage I non-small-cell lung cancer. *Int J Radiat Oncol Biol Phys* 2008;70:685-692.
- 39 Chang JY, et al. Stereotactic body radiation therapy for centrally located early stage or isolated parenchymal recurrences of non-small-cell lung cancer: how to fly in a "no fly zone". *Int J Radiat Oncol Biol Phys* 2014;88:1120-1128.
- 40 Timmerman R, et al. Excessive toxicity when treating central tumors in a phase II study of stereotactic body radiation therapy for medically inoperable early-stage lung cancer. *J Clin Oncol* 2006; 24:4833-4839.
- 41 Chaudhuri AA, et al. Stereotactic ablative radiotherapy for treatment of central and ultra-central lung tumors. *Lung Cancer* 2015;89:50-56.

[Continued](#)**Note:** All recommendations are category 2A unless otherwise indicated.**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

- ⁴² Haseltine JM, et al. Fatal complications after stereotactic body radiation therapy for central lung tumors abutting the proximal bronchial tree. *Pract Radiat Oncol* 2016;6:e27-33.
- ⁴³ Woody NM, et al. Stereotactic body radiation therapy for non-small cell lung cancer tumors greater than 5 cm: safety and efficacy. *Int J Radiat Oncol Biol Phys* 2015;92:325-331.
- ⁴⁴ Bezjak A, et al. Safety and efficacy of a five-fraction stereotactic body radiotherapy schedule for centrally located non-small-cell lung cancer: NRG Oncology/RTOG 0813 Trial. *J Clin Oncol* 2019;37:1316-1325.
- ⁴⁵ Fakiris AJ, et al. Stereotactic body radiation therapy for early-stage non-small-cell lung carcinoma: four-year results of a prospective phase II study. *Int J Radiat Oncol Biol Phys* 2009;75:677-682.
- ⁴⁶ Xiao Y, et al. Dosimetric evaluation of heterogeneity corrections for RTOG 0236: stereotactic body radiotherapy of inoperable stage I-II non-small-cell lung cancer. *Int J Radiat Oncol Biol Phys* 2009;73:1235-1242.
- ⁴⁷ Zhao L, et al. Planning target volume D95 and mean dose should be considered for optimal local control for stereotactic ablative radiation therapy. *Int J Radiat Oncol Biol Phys* 2016;95:1226-35.
- ⁴⁸ Aupérin A, et al. Meta-analysis of concomitant versus sequential radiochemotherapy in locally advanced non-small-cell lung cancer. *J Clin Oncol* 2010;28:2181-2190.
- ⁴⁹ O'Rourke N, et al. Concurrent chemoradiotherapy in non-small cell lung cancer. *Cochrane Database Syst Rev* 2010;CD002140.
- ⁵⁰ Curran WJ Jr, et al. Sequential vs. concurrent chemoradiation for stage III non-small cell lung cancer: randomized phase III trial RTOG 9410. *J Natl Cancer Inst* 2011;103:1452-1460.
- ⁵¹ Sause W, et al. Final results of phase III trial in regionally advanced unresectable non-small cell lung cancer: Radiation Therapy Oncology Group, Eastern Cooperative Oncology Group, and Southwest Oncology Group. *Chest* 2000;117:358-364.
- ⁵² Dillman RO, et al. Improved survival in stage III non-small-cell lung cancer: seven-year follow-up of cancer and leukemia group B (CALGB) 8433 trial. *J Natl Cancer Inst* 1996;88:1210-1215.
- ⁵³ Baumann M, et al. Final results of the randomized phase III CHARTWEL-trial (ARO 97-1) comparing hyperfractionated-accelerated versus conventionally fractionated radiotherapy in non-small cell lung cancer (NSCLC). *Radiother Oncol* 2011;100:76-85.
- ⁵⁴ Mauguén A, et al. Hyperfractionated or accelerated radiotherapy in lung cancer: an individual patient data meta-analysis. *J Clin Oncol* 2012;30:2788-2797.
- ⁵⁵ Albain KS, et al. Radiotherapy plus chemotherapy with or without surgical resection for stage III non-small-cell lung cancer: a phase III randomised controlled trial. *Lancet* 2009;374:379-386.
- ⁵⁶ Higgins K, et al. Preoperative chemotherapy versus preoperative chemoradiotherapy for stage III (N2) non-small-cell lung cancer. *Int J Radiat Oncol Biol Phys* 2009;75:1462-1467.
- ⁵⁷ Sher DJ, et al. Comparative effectiveness of neoadjuvant chemoradiotherapy versus chemotherapy alone followed by surgery for patients with stage IIIA non-small cell lung cancer. *Lancet Oncol* 2015;88:267-274.
- ⁵⁸ Rusch VW, et al. Induction chemoradiation and surgical resection for superior sulcus non-small-cell lung carcinomas: long-term results of Southwest Oncology Group Trial 9416 (Intergroup Trial 0160). *J Clin Oncol* 2007;25:313-318.
- ⁵⁹ Thomas M, et al. Effect of preoperative chemoradiation in addition to preoperative chemotherapy: a randomized trial in stage III non-small-cell lung cancer. *Lancet Oncol* 2008;9:607-608.
- ⁶⁰ Shah AA, et al. Induction chemoradiation is not superior to induction chemotherapy alone in stage IIIA lung cancer. *Ann Thorac Surg* 2012;93:1807-1812.
- ⁶¹ Douillard J-Y, et al. Impact of postoperative radiation therapy on survival in patients with complete resection and stage I, II, or IIIA non-small-cell lung cancer treated with adjuvant chemotherapy: the adjuvant Navelbine International Trialist Association (ANITA) Randomized Trial. *Int J Radiat Oncol Biol Phys* 2008;72:695-701.
- ⁶² Le Pechoux C, et al. Postoperative radiotherapy versus no postoperative radiotherapy in patients with completely resected non-small-cell lung cancer and proven mediastinal N2 involvement (Lung ART): an open-label, randomised, phase 3 trial. *Lancet Oncol* 2022;23:104-114.
- ⁶³ Hui Z, et al. Effect of postoperative radiotherapy for patients with pIIIA-N2 non-small cell lung cancer after complete resection and adjuvant chemotherapy: the phase 3 PORT-C randomized clinical trial. *JAMA Oncol* 2021;7:1178-1185.
- ⁶⁴ Bradley JD, et al. Phase II trial of postoperative adjuvant paclitaxel/carboplatin and thoracic radiotherapy in resected stage II and IIIA non-small-cell lung cancer: promising long-term results of the Radiation Therapy Oncology Group–RTOG 9705. *J Clin Oncol* 2005;23:3480-3487.
- ⁶⁵ Keller SM, et al. A randomized trial of postoperative adjuvant therapy in patients with completely resected stage II or IIIA non-small-cell lung cancer. *Eastern Cooperative Oncology Group. N Engl J Med* 2000; 343:1217-1222.
- ⁶⁶ Hancock JG, et al. Impact of adjuvant treatment for microscopic residual disease after non-small cell lung cancer surgery. *Ann Thorac Surg* 2015;99:406-416.
- ⁶⁷ Burdett S, Stewart L, Group PM-a. Postoperative radiotherapy in non-small-cell lung cancer: update of an individual patient data meta-analysis. *Lung Cancer* 2005;47:81-83.
- ⁶⁸ Belderbos JS, et al. Report from the International Atomic Energy Agency (IAEA) consultants' meeting on elective nodal irradiation in lung cancer: non-small cell lung cancer (NSCLC). *Int J Radiat Oncol Biol Phys* 2008;72:335-342.
- ⁶⁹ Bradley J, et al. A phase II comparative study of gross tumor volume definition with or without PET/CT fusion in dosimetric planning for non-small-cell lung cancer (NSCLC): primary analysis of radiation therapy oncology group (RTOG) 0515. *Int J Radiat Oncol Biol Phys* 2012;82:435-441.
- ⁷⁰ Sanuki-Fujimoto N, et al. Relation between elective nodal failure and irradiated volume in non-small-cell lung cancer (NSCLC) treated with radiotherapy using conventional fields and doses. *Radiother Oncol* 2009; 91:433-437.
- ⁷¹ Sulman EP, et al. Exclusion of elective nodal irradiation is associated with minimal elective nodal failure in non-small cell lung cancer. *Radiat Oncol* 2009;4:5-11.
- ⁷² Rosenzweig KE, Sura S, Jackson A, Yorke E. Involved-field radiation therapy for inoperable non small-cell lung cancer. *J Clin Oncol* 2007;25:5557-5561.
- ⁷³ Yuan S, et al. A randomized study of involved-field irradiation versus elective nodal irradiation in combination with concurrent chemotherapy for inoperable stage III non-small cell lung cancer. *Am J Clin Oncol* 2007;30:239-244.
- ⁷⁴ Chen M, et al. Involved-field radiotherapy versus elective nodal irradiation in combination with concurrent chemotherapy for locally advanced non-small cell lung cancer: a prospective randomized study. *Biomed Res Int* 2013;37:11819.
- ⁷⁵ Nestle U, et al. Imaging-based target volume reduction in chemoradiotherapy for locally advanced non-small-cell lung cancer (PET-Plan): a multicentre, open-label, randomised, controlled trial. *Lancet Oncol* 2020;21:581-592.
- ⁷⁶ Perez CA, et al. Long-term observations of the patterns of failure in patients with unresectable non-oat cell carcinoma of the lung treated with definitive radiotherapy. Report by the Radiation Therapy Oncology Group. *Cancer* 1987;59:1874-1881.
- ⁷⁷ Kong FM, et al. High-dose radiation improved local tumor control and overall survival in patients with inoperable/unresectable non-small-cell lung cancer: long-term results of a radiation dose escalation study. *Int J Radiat Oncol Biol Phys* 2005;63:324-333.

Note: All recommendations are category 2A unless otherwise indicated.**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.[Continued](#)**NSCL-C
10 OF 11**



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

- ⁷⁸ Rengan R, et al. Improved local control with higher doses of radiation in large-volume stage III non-small-cell lung cancer. *Int J Radiat Oncol Biol Phys* 2004;60:741-747.
- ⁷⁹ Machtay M, et al. Higher biologically effective dose of radiotherapy is associated with improved outcomes for locally advanced non-small cell lung carcinoma treated with chemoradiation: an analysis of the radiation therapy oncology group. *Int J Radiat Oncol Biol Phys* 2012;82:425-434.
- ⁸⁰ Schild SE, et al. Results of a phase I trial of concurrent chemotherapy and escalating doses of radiation for unresectable non-small-cell lung cancer. *Int J Radiat Oncol Biol Phys* 2006;65:1106-1111.
- ⁸¹ Socinski MA, et al. Randomized phase II trial of induction chemotherapy followed by concurrent chemotherapy and dose-escalated thoracic conformal radiotherapy (74 Gy) in stage III non-small-cell lung cancer: CALGB 30105. *J Clin Oncol* 2008;26:2457-2463.
- ⁸² Stinchcombe TE, et al. Long-term follow-up of a phase I/II trial of dose escalating three-dimensional conformal thoracic radiation therapy with induction and concurrent carboplatin and paclitaxel in unresectable stage IIIA/B non-small cell lung cancer. *J Thorac Oncol* 2008;3:1279-1285.
- ⁸³ Bradley JD, et al. Primary analysis of the phase II component of a phase I/II dose intensification study using three-dimensional conformal radiation therapy and concurrent chemotherapy for patients with inoperable non-small-cell lung cancer: RTOG 0117. *J Clin Oncol* 2010;28:2475-2480.
- ⁸⁴ Bradley JD, et al. Standard-dose versus high-dose conformal radiotherapy with concurrent and consolidation carboplatin plus paclitaxel with or without cetuximab for patients with stage IIIA or IIIB non-small-cell lung cancer (RTOG 0617): a randomised, two-by-two factorial phase 3 study. *Lancet Oncol* 2015;16:187-199.
- ⁸⁵ Schild SE, et al. Exploring radiotherapy targeting strategy and dose: a pooled analysis of cooperative group trials of combined modality therapy for stage III NSCLC. *J Thorac Oncol* 2018;13:1171-1182.
- ⁸⁶ Maugen A, et al. Hyperfractionated or accelerated radiotherapy in lung cancer: an individual patient data meta-analysis. *J Clin Oncol* 2012;30:2788-2797.
- ⁸⁷ Kong F-M S, et al. NRG-TOG 1106/ACRIN 6697: A phase IIR trial of standard versus adaptive (mid-treatment PET-based) chemoradiotherapy for stage III NSCLC—Results and comparison to NRG-TOG 0617 (non-personalized RT dose escalation). *J Clin Oncol* 2021;39:8548-8548.
- ⁸⁸ Sher DJ, et al. Relationship between radiation therapy dose and outcome in patients treated with neoadjuvant chemoradiation therapy and surgery for stage IIIA non-small cell lung cancer: a population-based, comparative effectiveness analysis. *Int J Radiat Oncol Biol Phys* 2015;92:307-316.
- ⁸⁹ Cerfolio RJ, et al. Pulmonary resection after concurrent chemotherapy and high dose (60Gy) radiation for non-small cell lung cancer is safe and may provide increased survival. *Eur J Cardiothorac Surg* 2009;35:718-723; discussion 723.
- ⁹⁰ Kwong KF, et al. High-dose radiotherapy in trimodality treatment of Pancoast tumors results in high pathologic complete response rates and excellent long-term survival. *J Thorac Cardiovasc Surg* 2005;129:1250-1257.
- ⁹¹ Sonett JR, et al. Pulmonary resection after curative intent radiotherapy (>59 Gy) and concurrent chemotherapy in non-small-cell lung cancer. *Ann Thorac Surg* 2004;78:1200-1205.
- ⁹² Suntharalingam M, et al. Radiation therapy oncology group protocol 02-29: a phase II trial of neoadjuvant therapy with concurrent chemotherapy and full-dose radiation therapy followed by surgical resection and consolidative therapy for locally advanced non-small cell carcinoma of the lung. *Int J Radiat Oncol Biol Phys* 2012;84:456-463.
- ⁹³ Kelsey CR, Light KL, Marks LB. Patterns of failure after resection of non-small-cell lung cancer: implications for postoperative radiation therapy volumes. *Int J Radiat Oncol Biol Phys* 2006;65:1097-1105.
- ⁹⁴ Lally BE, et al. Postoperative radiotherapy for stage II or III non-small-cell lung cancer using the surveillance, epidemiology, and end results database. *J Clin Oncol* 2006;24:2998-3006.
- ⁹⁵ Feigenberg SJ, et al. A phase II study of concurrent carboplatin and paclitaxel and thoracic radiotherapy for completely resected stage II and IIIA non-small cell lung cancer. *J Thorac Oncol* 2007;2:287-292.
- ⁹⁶ Corso CD, et al. Re-evaluation of the role of postoperative radiotherapy and the impact of radiation dose for non-small-cell lung cancer using the National Cancer Database. *J Thorac Oncol* 2015;10:148-155.
- ⁹⁷ Spoelstra FOB, et al. Variations in target volume definition for postoperative radiotherapy in stage III non-small-cell lung cancer: analysis of an international contouring study. *Int J Radiat Oncol Biol Phys* 2010;76:1106-1113.
- ⁹⁸ Ashworth AB, et al. An individual patient data metaanalysis of outcomes and prognostic factors after treatment of oligometastatic non-small-cell lung cancer. *Clin Lung Cancer* 2014;15:346-355.
- ⁹⁹ Milano MT, Katz AW, Okunieff P. Patterns of recurrence after curative-intent radiation for oligometastases confined to one organ. *Am J Clin Oncol* 2010;33:157-163.
- ¹⁰⁰ Salama JK, et al. An initial report of a radiation dose-escalation trial in patients with one to five sites of metastatic disease. *Clin Cancer Res* 2008;14:5255-5259.
- ¹⁰¹ Gomez DR, et al. Local consolidative therapy versus maintenance therapy or observation for patients with oligometastatic non-small-cell lung cancer without progression after first-line systemic therapy: a multicentre, randomised, controlled, phase 2 study. *Lancet Oncol* 2016;17:1672-1682.
- ¹⁰² Gomez DR, et al. Local consolidative therapy vs. maintenance therapy or observation for patients with oligometastatic non-small-cell lung cancer: long-term results of a multi-institutional, phase II, randomized study. *J Clin Oncol* 2019;37:1558-1565.
- ¹⁰³ Iyengar P, et al. Consolidative radiotherapy for limited metastatic non-small-cell lung cancer: A phase 2 randomized clinical trial. *JAMA Oncol* 2018;4:e173501.
- ¹⁰⁴ Theelen WSME, et al. Pembrolizumab with or without radiotherapy for metastatic non-small-cell lung cancer: a pooled analysis of two randomised trials. *Lancet Respir Med* 2021;9:467-475.
- ¹⁰⁵ Chow E, et al. Palliative radiotherapy trials for bone metastases: a systematic review. *J Clin Oncol* 2007;25:1423-1436.
- ¹⁰⁶ Lutz S, et al. Palliative radiotherapy for bone metastases: an ASTRO evidence-based guideline. *Int J Radiat Oncol Biol Phys* 2011;79:965-976.
- ¹⁰⁷ Cross CK, et al. Prospective study of palliative hypofractionated radiotherapy (8.5 Gy x 2) for patients with symptomatic non-small-cell lung cancer. *Int J Radiat Oncol Biol Phys* 2004;58:1098-1105.
- ¹⁰⁸ Medical Research Council Lung Cancer Working Party. A Medical Research Council (MRC) randomised trial of palliative radiotherapy with two fractions or a single fraction in patients with inoperable non-small-cell lung cancer (NSCLC) and poor performance status. *Medical Research Council Lung Cancer Working Party. Br J Cancer* 1992;65:934-941.
- ¹⁰⁹ Rodrigues G, et al. Palliative thoracic radiotherapy in lung cancer: An American Society for Radiation Oncology evidence-based clinical practice guideline. *Pract Radiat Oncol* 2011;1:60-71.
- ¹¹⁰ Koshy M, et al. Comparative effectiveness of aggressive thoracic radiation therapy and concurrent chemoradiation therapy in metastatic lung cancer. *Pract Radiat Oncol* 2015;5:374-382.
- ¹¹¹ Nguyen QN, et al. Single-fraction stereotactic vs conventional multifraction radiotherapy for pain relief in patients with predominantly nonspine bone metastases: a randomized phase 2 trial. *JAMA Oncol* 2019;5:872-878.
- ¹¹² Sahgal A, et al. Stereotactic body radiotherapy versus conventional external beam radiotherapy in patients with painful spinal metastases: an open-label, multicentre, randomised, controlled, phase 2/3 trial. *Lancet Oncol* 2021;22:1023-1033.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

**PRINCIPLES OF IMAGE-GUIDED THERMAL ABLATION THERAPY****General Principles**

- **Interventional radiologists should actively participate in multidisciplinary discussions and meetings regarding patients with NSCLC (eg, multidisciplinary clinic and/or tumor board).**
- **Decisions about whether ablation is feasible should be performed by interventional radiologists who perform IGTA as a prominent part of their practice.**
- **IGTA includes radiofrequency ablation, microwave ablation, and cryoablation. IGTA is a form of “local therapy” or “local ablative therapy.”¹**
- **IGTA is a lung parenchymal sparing technique with at most a temporary decrement in FEV1 and DLCO, which is statistically indistinguishable from baseline after recovery.²⁻⁶**

Evaluation

- **IGTA may be considered for those patients who are deemed “high risk”—those with tumors that are for the most part surgically resectable but rendered medically inoperable due to comorbidities. In cases where IGTA is considered for high-risk or borderline operable patients, a multidisciplinary evaluation is recommended.**
- **IGTA has been successfully accomplished in patients considered “high risk,” objectively defined with a single major and/or two or more minor criteria. Major criteria included an FEV1 or DLCO ≤50%, and minor criteria included a less depressed FEV1 or DLCO between 51%–60%, age ≥75 years, pulmonary hypertension, LVEF ≤40%, resting or exercise PaO2 <55 mmHg, and pCO2 >45 mmHg.⁴**
- **If an interventional radiologist or center is uncertain about the feasibility or safety of IGTA or the use of IGTA for radiation failure, consider obtaining an additional interventional radiology opinion from a high-volume specialized center.**

Ablation

- **Each energy modality has advantages and disadvantages. Determination of energy modality to be used for ablation should take into consideration the size and location of the target tumor, risk of complication, as well as local expertise and/or operator familiarity.⁷**

Ablation for NSCLC

- **IGTA is an option for the management of NSCLC lesions <3 cm. Ablation for NSCLC lesions >3 cm may be associated with higher rates of local recurrence and complications.^{8,9}**
- **There is evidence on the use of IGTA for selected patients with stage 1A NSCLC, those who present with multiple lung cancers, or those who present with locoregional recurrence of symptomatic local thoracic disease.**
- **Like surgery, pneumothorax may occur after IGTA, particularly if multiple lesions are treated in a single session. Pneumothorax has been reported in 18.7%–45.7% of IGTA cases. Self-limited pneumothorax, not requiring chest tube placement, is an expected event and not considered a complication unless escalation of care is required. In 20.7% of IGTA cases, chest tube insertion may be required.¹⁰**

¹ Lam A, Yoshida EJ, Bui K, et al. Patient and facility demographics related outcomes in early-stage non-small cell lung cancer treated with radiofrequency ablation: a National Cancer Database analysis. *J Vasc Interv Radiol* 2018;29:1535-1541.

² Dupuy DE, DiPetrillo T, Gandhi S, et al. Radiofrequency ablation followed by conventional radiotherapy for medically inoperable stage I non-small cell lung cancer. *Chest* 2006;129:738-745.

³ Lencioni R, Crocetti L, Cioni R, et al. Response to radiofrequency ablation of pulmonary tumours: a prospective, intention-to-treat, multicentre clinical trial (the RAPTURE study). *Lancet Oncol* 2008;9:621-628.

⁴ Dupuy DE, Fernando HC, Hillman S, et al. Radiofrequency ablation of stage IA non-small cell lung cancer in medically inoperable patients: Results from the American College of Surgeons Oncology Group Z4033 (Alliance) trial. *Cancer* 2015;121:3491-3498.

⁵ de Baere T, Tselikas L, Woodrum D, et al. Evaluating cryoablation of metastatic lung tumors in patients—safety and efficacy: The ECLIPSE Trial—interim analysis at 1 year. *J Thorac Oncol* 2015;10:1468-1474.

⁶ Tada A, Hiraki T, Iguchi T, et al. Influence of radiofrequency ablation of lung cancer on pulmonary function. *Cardiovasc Intervent Radiol* 2012;35:860-867.

⁷ Abtin F, De Baere T, Dupuy DE, et al. Updates on current role and practice of lung ablation. *J Thorac Imaging* 2019;34:266-277.

⁸ Lee JM, Jin GY, Goldberg SN, et al. Percutaneous radiofrequency ablation for inoperable non-small cell lung cancer and metastases: preliminary report. *Radiology* 2004;230:125-134.

⁹ Akeboshi M, Yamakado K, Nakatsuka A, et al. Percutaneous radiofrequency ablation of lung neoplasms: initial therapeutic response. *J Vasc Interv Radiol* 2004;15:463-470.

¹⁰ Genshaft SJ, Suh RD, Abtin F, et al. Society of Interventional Radiology Quality Improvement Standards on Percutaneous Ablation of Non-Small Cell Lung Cancer and Metastatic Disease to the Lungs. *J Vasc Interv Radiol* 2021;32:1242.e1-1242.e10.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

**PERIOPERATIVE SYSTEMIC THERAPY**

- **Neoadjuvant Systemic Therapy in Patients Who Are Candidates for Immune Checkpoint Inhibitors, see below.**
- [Neoadjuvant Systemic Therapy for Patients Who Are Not Candidates for Immune Checkpoint Inhibitors](#)
- [Adjuvant Chemotherapy](#)
- [Systemic Therapy Following Previous Neoadjuvant or Adjuvant Systemic Therapy](#)

Neoadjuvant Systemic Therapy

- All patients should be evaluated for preoperative therapy, with strong consideration for nivolumab or pembrolizumab + chemotherapy for those patients with tumors ≥ 4 cm or node positive and no contraindications to immune checkpoint inhibitors.^a Otherwise refer to the [Neoadjuvant Systemic Therapy for Patients Who Are Not Candidates for Immune Checkpoint Inhibitors](#).
- Test for PD-L1 status, *EGFR* mutations, and *ALK* rearrangements (stages IB–IIIA, IIIB [T3,N2]). PD-L1 status can be incorporated with other clinical factors to determine patients who may benefit from induction chemotherapy and immunotherapy. [Principles of Molecular and Biomarker Analysis \(NSCL-H\)](#).
- Clinical trials for neoadjuvant nivolumab + chemotherapy excluded patients harboring *EGFR* mutations and *ALK* rearrangements. Thus, exclusion of these biomarkers, at a minimum, is recommended prior to consideration for neoadjuvant nivolumab + chemotherapy.
- After surgical evaluation, patients likely to receive adjuvant chemotherapy may be treated with induction systemic therapy as an alternative.

Neoadjuvant Systemic Therapy in Patients Who Are Candidates for Immune Checkpoint Inhibitors

- Nivolumab 360 mg and platinum-doublet chemotherapy every 3 weeks for 3 cycles¹
 - ▶ Platinum-doublet chemotherapy options include:
 - ◊ Carboplatin AUC 5 or AUC 6 day 1, paclitaxel 175 mg/m² or 200 mg/m² day 1 (any histology)
 - ◊ Cisplatin 75 mg/m² day 1, pemetrexed 500 mg/m² day 1 (nonsquamous histology)
 - ◊ Cisplatin 75 mg/m² day 1, gemcitabine 1000 mg/m² or 1250 mg/m² days 1 and 8 (squamous histology)
 - ◊ Cisplatin 75 mg/m² day 1, paclitaxel 175 mg/m² or 200 mg/m² day 1 (any histology)
 - ▶ Chemotherapy regimens for patients who are not candidates for cisplatin-based therapy
 - ◊ Carboplatin AUC 5 or AUC 6 day 1, pemetrexed 500 mg/m² day 1 (nonsquamous histology)
 - ◊ Carboplatin AUC 5 or AUC 6 day 1, gemcitabine 1000 mg/m² or 1250 mg/m² days 1 and 8 (squamous histology)
- Pembrolizumab 200 mg and cisplatin-based doublet therapy every 3 weeks for 4 cycles and then continued as single-agent pembrolizumab as adjuvant treatment after surgery (category 1); [Systemic Therapy Following Previous Neoadjuvant or Adjuvant Systemic Therapy](#)²
 - ▶ Cisplatin 75 mg/m² day 1, gemcitabine 1000 mg/m² days 1 and 8 (squamous histology)
 - ▶ Cisplatin 75 mg/m² day 1, pemetrexed 500 mg/m² day 1 (nonsquamous histology)

^a Contraindications for treatment with PD-1/PD-L1 inhibitors may include active or previously documented autoimmune disease and/or current use of immunosuppressive agents; some oncogenic drivers (ie, *EGFR* exon 19 deletion or exon 21 L858R, *ALK* rearrangements) have been shown to be associated with less benefit from PD-1/PD-L1 inhibitors.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

[References](#)



PERIOPERATIVE SYSTEMIC THERAPY

Neoadjuvant Systemic Therapy for Patients Who Are Not Candidates for Immune Checkpoint Inhibitors

Preferred (nonsquamous)

- Cisplatin 75 mg/m² day 1, pemetrexed 500 mg/m² day 1 every 21 days for 4 cycles³

Preferred (squamous)

- Cisplatin 75 mg/m² day 1, gemcitabine 1250 mg/m² days 1 and 8, every 21 days for 4 cycles⁴
- Cisplatin 75 mg/m² day 1, docetaxel 75 mg/m² day 1 every 21 days for 4 cycles⁵

Other Recommended

- Cisplatin 50 mg/m² days 1 and 8; vinorelbine 25 mg/m² days 1, 8, 15, and 22, every 28 days for 4 cycles⁶
- Cisplatin 100 mg/m² day 1, vinorelbine 30 mg/m² days 1, 8, 15, and 22, every 28 days for 4 cycles^{7,8}
- Cisplatin 75–80 mg/m² day 1, vinorelbine 25–30 mg/m² days 1 and 8, every 21 days for 4 cycles
- Cisplatin 100 mg/m² day 1, etoposide 100 mg/m² days 1–3, every 28 days for 4 cycles⁷

Useful in Certain Circumstances

- Chemotherapy regimens for patients who are not candidates for cisplatin-based therapy
 - Carboplatin AUC 6 day 1, paclitaxel 200 mg/m² day 1, every 21 days for 4 cycles⁹
 - Carboplatin AUC 5 day 1, gemcitabine 1000 mg/m² days 1 and 8, every 21 days for 4 cycles¹⁰ (squamous histology)
 - Carboplatin AUC 5 day 1, pemetrexed 500 mg/m² day 1 every 21 days for 4 cycles¹¹ (nonsquamous histology)

All chemotherapy regimens listed above can be used for sequential chemotherapy/RT.

Neoadjuvant Systemic Therapy in Patients Who Are Candidates for Immune Checkpoint Inhibitors

Adjuvant Chemotherapy

Systemic Therapy Following Previous Neoadjuvant or Adjuvant Systemic Therapy

References

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



PERIOPERATIVE SYSTEMIC THERAPY

Adjuvant Chemotherapy

- For stage IIA (T2b, N0) and negative margins, adjuvant chemotherapy is recommended for high-risk features.^b
- For stage IIB (T1 abc–T2a, N1), stage IIB (T3, N0; T2b, N1), stage IIIA (T1–2, N2; T3, N1), stage IIIB (T3, N2) and negative margins (R0), adjuvant chemotherapy is recommended as a category 1.
- Test for PD-L1 status, *EGFR* mutations, and *ALK* rearrangements (stages II–IIIA, IIIB [T3,N2]).

[Principles of Molecular and Biomarker Analysis \(NSCL-H\).](#)

Preferred (nonsquamous)

- Cisplatin 75 mg/m² day 1, pemetrexed 500 mg/m² day 1 every 21 days for 4 cycles³

Preferred (squamous)

- Cisplatin 75 mg/m² day 1, gemcitabine 1250 mg/m² days 1 and 8, every 21 days for 4 cycles⁴
- Cisplatin 75 mg/m² day 1, docetaxel 75 mg/m² day 1 every 21 days for 4 cycles⁵

Other Recommended

- Cisplatin 50 mg/m² days 1 and 8; vinorelbine 25 mg/m² days 1, 8, 15, and 22, every 28 days for 4 cycles⁶
- Cisplatin 100 mg/m² day 1, vinorelbine 30 mg/m² days 1, 8, 15, and 22, every 28 days for 4 cycles^{7,8}
- Cisplatin 75–80 mg/m² day 1, vinorelbine 25–30 mg/m² days 1 and 8, every 21 days for 4 cycles
- Cisplatin 100 mg/m² day 1, etoposide 100 mg/m² days 1–3, every 28 days for 4 cycles⁷

Useful in Certain Circumstances

- Chemotherapy regimens for patients who are not candidates for cisplatin-based therapy
 - ▶ Carboplatin AUC 6 day 1, paclitaxel 200 mg/m² day 1, every 21 days for 4 cycles⁸
 - ▶ Carboplatin AUC 5 day 1, gemcitabine 1000 mg/m² days 1 and 8, every 21 days for 4 cycles¹⁰
 - ▶ Carboplatin AUC 5 day 1, pemetrexed 500 mg/m² day 1 every 21 days for 4 cycles¹¹ (nonsquamous histology)

All chemotherapy regimens listed above can be used for sequential chemotherapy/RT.

[Neoadjuvant Systemic Therapy in Patients Who Are Candidates for Immune Checkpoint Inhibitors](#)

[Neoadjuvant Systemic Therapy for Patients Who Are Not Candidates for Immune Checkpoint Inhibitors](#)

[Systemic Therapy Following Previous Neoadjuvant or Adjuvant Systemic Therapy](#)

^b Examples of high-risk features may include poorly differentiated tumors (including lung neuroendocrine tumors [excluding well-differentiated neuroendocrine tumors]), vascular invasion, wedge resection, visceral pleural involvement, and unknown lymph node status (Nx). These factors independently may not be an indication and may be considered when determining treatment with adjuvant chemotherapy.

[References](#)

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



PERIOPERATIVE SYSTEMIC THERAPY

Systemic Therapy Following Previous Neoadjuvant or Adjuvant Systemic Therapy^c

- Test for PD-L1 status, *EGFR* mutations, and *ALK* rearrangements (stages IB–IIIA, IIIB [T3,N2]).

[Principles of Molecular and Biomarker Analysis \(NSCL-H\).](#)

- Osimertinib 80 mg daily¹²
 - ▶ For patients with completely resected stage IB–IIIA or stage IIIB (T3, N2) NSCLC and positive for *EGFR* (exon 19 deletion, exon 21 L858R) mutations who received previous adjuvant chemotherapy or are ineligible to receive platinum-based chemotherapy.
- Atezolizumab 840 mg every 2 weeks, 1200 mg every 3 weeks, or 1680 mg every 4 weeks for up to 1 year¹³
 - ▶ For patients with completely resected stage IIB–IIIA, stage IIIB (T3, N2), or high-risk stage IIA NSCLC with PD-L1 $\geq 1\%$ and negative for *EGFR* exon 19 deletion or exon 21 L858R mutations or *ALK* rearrangements who received previous adjuvant chemotherapy and with no contraindications to immune checkpoint inhibitors.^a
- Pembrolizumab 200 mg every 3 weeks or 400 mg every 6 weeks for up to 1 year
 - ▶ For patients with completely resected stage IIB–IIIA, stage IIIB (T3, N2), or high-risk stage IIA NSCLC and negative for *EGFR* exon 19 deletion or exon 21 L858R mutations or *ALK* rearrangements who received previous adjuvant chemotherapy and with no contraindications to immune checkpoint inhibitors.^{a,14} The benefit for patients with PD-L1 $<1\%$ is unclear.
 - ▶ For patients with completely resected stage II–IIIA or stage IIIB (T3, N2) NSCLC who received previous neoadjuvant pembrolizumab + chemotherapy (category 1).²

[Neoadjuvant Systemic Therapy in Patients Who Are Candidates for Immune Checkpoint Inhibitors](#)

[Neoadjuvant Systemic Therapy for Patients Who Are Not Candidates for Immune Checkpoint Inhibitors](#)

[Adjuvant Chemotherapy](#)

^a Contraindications for treatment with PD-1/PD-L1 inhibitors may include active or previously documented autoimmune disease and/or current use of immunosuppressive agents; some oncogenic drivers (ie, *EGFR* exon 19 deletion or exon 21 L858R, *ALK* rearrangements) have been shown to be associated with less benefit from PD-1/PD-L1 inhibitors.

^c In general, perioperative therapy should be given as a single regimen and change of immunotherapy is not recommended.

[References](#)

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

**PERIOPERATIVE SYSTEMIC THERAPY – REFERENCES**

- ¹ Forde PM, Spicer J, Lu S, et al. Neoadjuvant nivolumab plus chemotherapy in resectable lung cancer. *N Engl J Med* 2022;386:1973-1985.
- ² Wakelee H, Liberman M, Kato T, et al. Perioperative pembrolizumab for early-stage non-small-cell lung cancer. *N Engl J Med* 2023;389:491-503.
- ³ Kreuter M, Vansteenkiste J, Fishcer JR, et al. Randomized phase 2 trial on refinement of early-stage NSCLC adjuvant chemotherapy with cisplatin and pemetrexed versus cisplatin and vinorelbine: the TREAT study. *Ann Oncol* 2013;24:986-992.
- ⁴ Pérol M, Chouaid C, Pérol D, et al. Randomized, phase III study of gemcitabine or erlotinib maintenance therapy versus observation, with predefined second-line treatment, after cisplatin-gemcitabine induction chemotherapy in advanced non-small-cell lung cancer. *J Clin Oncol* 2012;30:3516-3524.
- ⁵ Fossella F, Pereira JR, von Pawel J, et al. Randomized, multinational, phase III study of docetaxel plus platinum combinations versus vinorelbine plus cisplatin for advanced non-small-cell lung cancer: the TAX 326 study group. *J Clin Oncol* 2003;21:3016-3024.
- ⁶ Winton T, Livingston R, Johnson D, et al. Vinorelbine plus cisplatin vs. observation in resected non-small-lung cancer. *N Engl J Med* 2005;352:2589-2597.
- ⁷ Arriagada R, Bergman B, Dunant A, et al. The International Adjuvant Lung Cancer Trial Collaborative Group. Cisplatin-based adjuvant chemotherapy in patients with completely resected non-small cell lung cancer. *N Engl J Med* 2004;350:351-360.
- ⁸ Douillard JY, Rosell R, De Lena M, et al. Adjuvant vinorelbine plus cisplatin versus observation in patients with completely resected stage IB-IIIa non-small-cell lung cancer (Adjuvant Navelbine International Trialist Association [ANITA]): a randomised controlled trial. *Lancet Oncol* 2006;7:719-727.
- ⁹ Strauss GM, Herndon III JE, Maddaus MA, et al. Adjuvant paclitaxel plus carboplatin compared with observation in stage IB non-small cell lung cancer: CALGB 9633 with the Cancer and Leukemia Group B, Radiation Therapy Oncology Group, and North Central Cancer Treatment Group Study Groups. *J Clin Oncol* 2008;26:5043-5051.
- ¹⁰ Usami N, Yokoi K, Hasegawa Y, et al. Phase II study of carboplatin and gemcitabine as adjuvant chemotherapy in patients with completely resected non-small cell lung cancer: a report from the Central Japan Lung Study Group, CJLSG 0503 trial. *Int J Clin Oncol* 2010;15:583-587.
- ¹¹ Zhang L, Ou W, Liu Q, et al. Pemetrexed plus carboplatin as adjuvant chemotherapy in patients with curative resected non-squamous non-small cell lung cancer. *Thorac Cancer* 2014;5:50-56.
- ¹² Wu Y-L, Tsuboi M, He J, et al. Osimertinib in resected EGFR-mutated non-small-cell lung cancer. *N Engl J Med* 2020;383:1711-1723.
- ¹³ Felip E, Altorki N, Zhou C, et al. Adjuvant atezolizumab after adjuvant chemotherapy in resected stage IB–IIIA non-small-cell lung cancer (IMpower010): a randomised, multicentre, open-label, phase 3 trial. *Lancet* 2021;398:1344-1357.
- ¹⁴ O'Brien M, Paz-Ares L, Marreaud S, et al. Pembrolizumab versus placebo as adjuvant therapy for completely resected stage IB-IIIa non-small-cell lung cancer (PEARLS/KEYNOTE-091): an interim analysis of a randomised, triple-blind, phase 3 trial. *Lancet Oncol* 2022;23:1274-1286.

Note: All recommendations are category 2A unless otherwise indicated.**Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.**



CONCURRENT CHEMORADIATION REGIMENS

Concurrent Chemoradiation Regimens^a

Preferred (nonsquamous)

- Carboplatin AUC 5 on day 1, pemetrexed 500 mg/m² on day 1 every 21 days for 4 cycles; concurrent thoracic RT^{1,b,c,d}
- Cisplatin 75 mg/m² on day 1, pemetrexed 500 mg/m² on day 1 every 21 days for 3 cycles; concurrent thoracic RT^{2,3,b,c,d,e}
- Paclitaxel 45–50 mg/m² weekly; carboplatin AUC 2, concurrent thoracic RT^{4,b,c,d,f}
- Cisplatin 50 mg/m² on days 1, 8, 29, and 36; etoposide 50 mg/m² days 1–5 and 29–33; concurrent thoracic RT^{5,6,b,c,d}

Preferred (squamous)

- Paclitaxel 45–50 mg/m² weekly; carboplatin AUC 2, concurrent thoracic RT^{6,b,c,d,f}
- Cisplatin 50 mg/m² on days 1, 8, 29, and 36; etoposide 50 mg/m² days 1–5 and 29–33; concurrent thoracic RT^{5,6,b,c,d}

Consolidation Immunotherapy for Patients with Unresectable Stage II/III NSCLC, PS 0–1, and No Disease Progression After Definitive Concurrent Chemoradiation

Durvalumab 10 mg/kg IV every 2 weeks or 1500 mg every 4 weeks for up to 12 months (patients with a body weight of ≥30 kg)^{7,8,g,h}
(category 1 for stage III; category 2A for stage II)

^a For patients with superior sulcus tumors, the recommendation is for 2 cycles concurrent with RT and 2 more cycles after surgery. Rusch VW, Giroux DJ, Kraut MJ, et al. Induction chemoradiation and surgical resection for superior sulcus non-small-cell lung carcinomas: long-term results of Southwest Oncology Group Trial 9416 (Intergroup Trial 0160). J Clin Oncol 2007;25:313-318.

^b Regimens can be used as preoperative/adjuvant chemotherapy/RT.

^c Regimens can be used as definitive concurrent chemotherapy/RT.

^d For eligible patients, durvalumab may be used after noted concurrent chemotherapy/RT regimens.

^e If using durvalumab, additional chemotherapy after radiation is not recommended. If not using durvalumab, an additional 4 cycles of pemetrexed 500 mg/m² may be used.

^f If using durvalumab, additional chemotherapy after radiation is not recommended. If not using durvalumab, an additional 2 cycles every 21 days of paclitaxel 200 mg/m² and carboplatin AUC 6 may be used.

^g In patients with tumors that are positive for *EGFR* exon 19 deletion or exon 21 L858R mutations there is risk of toxicity when TKI is administered in temporal proximity to immuno-oncology (IO) therapy.

^h For patients who have received sequential chemoradiation, durvalumab can be considered as consolidation immunotherapy.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



CONCURRENT CHEMORADIATION REGIMENS – REFERENCES

- ¹ Govindan R, Bogart J, Stinchcombe T, et al. Randomized phase II study of pemetrexed, carboplatin, and thoracic radiation with or without cetuximab in patients with locally advanced unresectable non-small-cell lung cancer: Cancer and Leukemia Group B trial 30407. *J Clin Oncol* 2011;29:3120-3125.
- ² Choy H, Gerber DE, Bradley JD, et al. Concurrent pemetrexed and radiation therapy in the treatment of patients with inoperable stage III non-small cell lung cancer: a systematic review of completed and ongoing studies. *Lung Cancer* 2015;87:232-240.
- ³ Senan S, Brade A, Wang LH, et al. PROCLAIM: randomized phase III trial of pemetrexed-cisplatin or etoposide-cisplatin plus thoracic radiation therapy followed by consolidation chemotherapy in locally advanced nonsquamous non-small-cell lung cancer. *J Clin Oncol* 2016;34:953-962.
- ⁴ Bradley JD, Paulus R, Komaki R, et al. Standard-dose versus high-dose conformal radiotherapy with concurrent and consolidation carboplatin plus paclitaxel with or without cetuximab for patients with stage IIIA or IIIB non-small-cell lung cancer (RTOG 0617): a randomised, two-by-two factorial phase 3 study. *Lancet Oncol* 2015;16:187-199.
- ⁵ Albain KS, Crowley JJ, Turrisi AT III, et al. Concurrent cisplatin, etoposide, and chest radiotherapy in pathologic stage IIIB non-small-cell lung cancer: A Southwest Oncology Group Phase II Study, SWOG 9019. *J Clin Oncol* 2002;20:3454-3460.
- ⁶ Rusch VW, Giroux DJ, Kraut MJ, et al. Induction chemoradiation and surgical resection for superior sulcus non-small-cell lung carcinomas: long-term results of Southwest Oncology Group Trial 9416 (Intergroup Trial 0160). *J Clin Oncol* 2007;25:313-318.
- ⁷ Baverel PG, Dubois VFS, Jin CY, et al. Population pharmacokinetics of durvalumab in cancer patients and association with longitudinal biomarkers of disease status. *Clin Pharmacol Ther* 2018;103:631-642.
- ⁸ Faivre-Finn C, Vicente D, Kurata T, et al. Four-year survival with durvalumab after chemoradiotherapy in stage III NSCLC-an update from the PACIFIC trial. *J Thorac Oncol* 2021;16:860-867.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



CANCER SURVIVORSHIP CARE

NSCLC Long-Term Follow-up Care

- Cancer Surveillance ([NSCL-17](#))
- Immunizations
 - ▶ Annual influenza vaccination
 - ▶ Herpes zoster vaccine
 - ▶ Pneumococcal vaccination with revaccination as appropriate
 - ▶ COVID vaccination as per the guidance of the Centers for Disease Control and Prevention (CDC)
 - ▶ Hepatitis vaccination

• [NCCN Guidelines for Survivorship](#)

Counseling Regarding Health Promotion and Wellness¹

- Maintain a healthy weight
- Adopt a physically active lifestyle (regular physical activity: 30 minutes of moderate-intensity physical activity on most days of the week)
- Consume a healthy diet with emphasis on plant sources
- Limit consumption of alcohol if one consumes alcoholic beverages

Additional Health Monitoring

- Routine blood pressure, cholesterol, and glucose monitoring
- Bone health: Bone density testing as appropriate
- Dental health: Routine dental examinations
- Routine sun protection

Resources

- National Cancer Institute Facing Forward: Life After Cancer Treatment
<https://www.cancer.gov/publications/patient-education/facing-forward>

Cancer Screening Recommendations^{2,3}

These recommendations are for individuals at average risk for cancer; patients at high risk should be individualized.

- Colorectal Cancer:
[NCCN Guidelines for Colorectal Cancer Screening](#)
- Prostate Cancer:
[NCCN Guidelines for Prostate Cancer Early Detection](#)
- Breast Cancer:
[NCCN Guidelines for Breast Cancer Screening and Diagnosis](#)

¹ American Cancer Society Guideline for Diet and Physical Activity for Cancer Prevention: <https://www.cancer.org/cancer/risk-prevention/diet-physical-activity/acs-guidelines-nutrition-physical-activity-cancer-prevention.html>.

² Memorial Sloan Kettering Cancer Center Screening Guidelines: <https://www.mskcc.org/cancer-care/risk-assessment-screening/screening-guidelines>.

³ American Cancer Society Guidelines for the Early Detection of Cancer:
<http://www.cancer.org/healthy/findcancerearly/cancerscreeningguidelines/american-cancer-society-guidelines-for-the-early-detection-of-cancer?sitearea=PED>.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

**PRINCIPLES OF MOLECULAR AND BIOMARKER ANALYSIS****Molecular Diagnostic Studies in NSCLC**

- Numerous gene alterations have been identified that impact therapy selection. Testing of lung cancer specimens for these alterations is important for identification of potentially efficacious targeted therapies, as well as avoidance of therapies unlikely to provide clinical benefit.
- Some selection approaches for targeted therapy include predictive immunohistochemical analyses, which are distinct from immunohistochemical studies utilized to identify tumor type and lineage.
- Major elements of molecular testing that are critical for utilization and interpretation of molecular results include:
 - ▶ Use of a laboratory that is properly accredited, with a minimum of Clinical Laboratory Improvement Amendments (CLIA) accreditation
 - ▶ Understanding the methodologies that are utilized and the major limitations of those methodologies
 - ▶ Understanding the spectrum of alterations tested (and those not tested) by a specific assay
 - ▶ Knowledge of whether a tumor sample is subjected to pathologic review and tumor enrichment (ie, microdissection, macrodissection) prior to testing
 - ▶ The types of samples accepted by the testing laboratory
 - ▶ Understanding what constitutes an informative versus a null result. A positive finding for a known oncogenic driver is considered an informative result. Absence of a known oncogenic driver, including results that show only "passenger" alterations, might be considered null depending on the context of the specimen limitations and clinical situation. Specifically, technical issues related to tumor cellularity in tissue testing or burden of disease in circulating tumor DNA (ctDNA) testing are considerations in interpretation of findings. Additional considerations are elaborated in sections below.
- Tissue Specimen Acquisition and Management:
 - ▶ Although tumor testing has been primarily focused on use of FFPE tissues, increasingly, laboratories accept other specimen types, notably cytopathology preparations not processed by FFPE methods. Although testing on cell blocks is not included in the FDA approval for multiple companion diagnostic assays, testing on these specimen types is highly recommended when it is the only or best material.
 - ▶ A major limitation in obtaining tissue molecular testing results for NSCLC occurs when minimally invasive techniques are used to obtain samples; the yield may be insufficient for molecular, biomarker, and histologic testing. Therefore, bronchoscopists and interventional radiologists should procure sufficient tissue to enable all appropriate testing.
 - ▶ When tissue is minimal, laboratories should deploy techniques to maximize tissue for molecular and ancillary testing, including dedicated histology protocols for small biopsies, including "up-front" slide sectioning for diagnostic and predictive testing. Peripheral blood (plasma ctDNA) can be a surrogate sample ([NSCL-H 8 of 8](#)).
- Testing Methodologies
 - ▶ Appropriate possible testing methodologies are indicated below for each analyte separately; however, several methodologies are generally considerations for use:
 - ◊ Next-generation sequencing (NGS) is used in clinical laboratories. Not all types of alterations are detected by individual NGS assays and it is important to be familiar with the types of alterations identifiable in individual assays or combination(s) of assays.
 - Broad-based genomic testing approaches that efficiently utilize limited biopsy tissue while maximizing diagnostic genomic information are most commonly NGS-based. NCCN acknowledges that many currently available NGS-based assays used to fully genotype NSCLC are larger than the 50-gene limit threshold utilized by Current Procedural Terminology (CPT) coding convention; as a result, utilization of panels greater than 50 genes may be practical to follow these recommendations. Either a single assay or a combination of a limited number of assays may be appropriate.
 - Although broad-based genomic testing approaches are preferred, in some clinical situations rapid testing may be warranted and should be followed up with broad-based genomic testing.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

[Continued](#)

**PRINCIPLES OF MOLECULAR AND BIOMARKER ANALYSIS****• Testing Methodologies (continued)**

- ◊ It is recommended at this time that when feasible, testing be performed via a broad, panel-based approach, most typically performed by NGS. For patients who, in broad panel testing don't have identifiable driver oncogenes, consider RNA-based NGS if not already performed, to maximize detection of fusion events.
 - Broad molecular profiling is defined as molecular testing that identifies all biomarkers identified in [NSCL-20](#) in either a single assay or a combination of a limited number of assays, and optimally also identifies emerging biomarkers ([NSCL-I](#)). Tiered approaches based on low prevalence of co-occurring biomarkers are acceptable.
- ◊ Real-time polymerase chain reaction (PCR) can be used in a highly targeted fashion (specific mutations targeted). When this technology is deployed, only those specific alterations that are targeted by the assay are assessed and the potential for mutations outside of the testing scope of the specific assay must be considered.
- ◊ Sanger sequencing requires the greatest degree of tumor enrichment. Unmodified Sanger sequencing is not appropriate for detection of mutations in tumor samples with less than 25% to 30% tumor after enrichment and is not appropriate for assays in which identification of subclonal events (eg, resistance mutations) is important. If Sanger sequencing is utilized, tumor enrichment methodologies are nearly always recommended.
- ◊ Any method that interrogates sequences other than a subset of highly specific alterations (eg, NGS, Sanger) has the potential to identify variants of uncertain significance (VUS). Any variant classified as a VUS, even if in a gene in which other variants are clinically actionable, should not be considered as a basis for targeted therapy selection.
- ◊ Other methodologies may be utilized, including multiplex approaches not listed above.
- ◊ Fluorescence in situ hybridization (FISH) analysis is utilized for many assays examining copy number, amplification, and structural alterations such as gene rearrangements. FISH may have better sensitivity for gene amplification events in some circumstances.

• Specimen Selection

- ▶ Testing platforms using tissue and peripheral blood are available.
- ▶ Testing using tissue sample requires acquisition of a suitable sample.
 - ◊ Alterations detected using NSCLC tumor tissue can be most directly attributed to the tumor.
 - ◊ If an assay has a technical failure related to an insufficient quantity, consideration of an alternate testing modality or procurement of additional tissue is recommended to achieve broad molecular profiling.
- ▶ Testing using peripheral blood (most commonly plasma-based testing of circulating tumor ctDNA) can be utilized in conjunction with tissue-based testing to achieve genotyping for recommended biomarkers.
 - ◊ Many, but not all, ctDNA tests use NGS-based technology. The technical advantages and limitations of ctDNA testing are further elaborated below.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

[Continued](#)**NSCL-H**
2 OF 8

**PRINCIPLES OF MOLECULAR AND BIOMARKER ANALYSIS****• Molecular Targets for Analysis**

- ▶ In general, the mutations/alterations described below are seen in a non-overlapping fashion, although between 1%–3% of NSCLC may harbor concurrent alterations.
- ▶ ***EGFR* (Epidermal Growth Factor Receptor) Gene Mutations:** *EGFR* is a receptor tyrosine kinase normally found on the surface of epithelial cells and is often overexpressed in a variety of human malignancies.
 - ◊ The most commonly described mutations in *EGFR* (exon 19 deletions, p.L858R point mutation in exon 21) are associated with responsiveness to oral *EGFR* tyrosine kinase inhibitor (TKI) therapy; most recent data indicate that tumors that do not harbor a sensitizing *EGFR* mutation should not be treated with *EGFR* TKI in any line of therapy.
 - ◊ Molecular testing for *EGFR* mutations should be performed when adjuvant TKI therapy is a consideration for NSCLC stage IB–IIIA and stage IIIB (T3, N2). While the testing process may be technically easier on a resection specimen, initial diagnostic biopsy specimens are also acceptable for testing for this indication.
 - ◊ Many of the less commonly observed alterations in *EGFR*, which cumulatively account for ~10% of *EGFR*-mutation positive NSCLC (ie, exon 21 p.L861Q, exon 18 p.G719X, and exon 20 p.S768I mutations), are also associated with responsiveness to certain *EGFR* TKIs, such as osimertinib and afatinib, and should be considered on a mutation-specific basis, when possible.
 - ◊ ***EGFR* p.T790M** is most commonly observed as a mutation that arises in response to and as a mechanism of resistance to first- and second-generation *EGFR* TKI. In patients with progression on first- or second-generation TKI with p.T790M as the primary mechanism of resistance, third-generation TKIs are typically efficacious.
 - If *EGFR* p.T790M is identified in the absence of prior *EGFR* TKI therapy, genetic counseling and possible germline genetic testing are warranted. Identification of germline *EGFR* p.T790M confers a high risk for lung cancer regardless of smoking status.
 - ◊ ***EGFR* exon 20 (*EGFR*ex20) mutations** (other than *EGFR* p.T790M) are a heterogeneous group, some of which are responsive to targeted therapy and that require detailed knowledge of the specific alteration.
 - Most *EGFR*ex20 alterations are a diverse group of in-frame duplication or insertion mutations.
 - These are generally associated with lack of response to first-, second-, and third-generation *EGFR* TKI therapy, with select exceptions: p.A763_Y764insFQEA is associated with sensitivity to TKI therapy and p.A763_Y764insLQEA may be associated with sensitivity to first- and third-generation TKI therapy.
 - *EGFR*ex20 insertions/duplications are associated with responsiveness to specific targeted subsequent therapy agents designed for *EGFR*ex20 events. The most commonly represented *EGFR*ex20 insertions/duplications in the clinical studies have been insASV, insSVD, and insNPH, although a wide spectrum of other alterations were included. There is currently no evidence that the specific alteration type impacts the probability of responsiveness to this class of targeted therapy.
 - Because some *EGFR*ex20 mutations are or may be sensitive to first- and third-generation inhibitors, the specific sequence of *EGFR*ex20 insertion mutations remains important. Some assays will identify the presence of an *EGFR*ex20 insertion without specifying the sequence, and additional testing to further clarify the *EGFR*ex20 insertion may be indicated for therapy selection.
 - Targeted PCR-based approaches for detection of *EGFR* variants may under-detect *EGFR*ex20 insertion events; therefore, NGS-based strategies are preferred.
 - ◊ Some clinicopathologic features—such as smoking status, ethnicity, and histology—are associated with the presence of an *EGFR* mutation; however, these features should not be utilized in selecting patients for testing.
 - ◊ **Testing Methodologies:** Real-time PCR, Sanger sequencing (ideally paired with tumor enrichment), and NGS are the most commonly deployed methodologies for examining *EGFR* mutation status.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

[Continued](#)

**PRINCIPLES OF MOLECULAR AND BIOMARKER ANALYSIS**• **Molecular Targets for Analysis (continued)**

- ▶ **ALK (anaplastic lymphoma kinase) Gene Rearrangements:** ALK is a receptor tyrosine kinase that can be rearranged in NSCLC, resulting in dysregulation and inappropriate signaling through the ALK kinase domain.
 - ◊ The most common fusion partner seen with ALK is echinoderm microtubule-associated protein-like 4 (EML4), although a variety of other fusion partners have been identified.
 - ◊ The presence of an *ALK* rearrangement is associated with responsiveness to oral ALK TKIs.
 - ◊ Some clinicopathologic features—such as smoking status and histology—have been associated with the presence of an *ALK* rearrangement; however, these features should not be utilized in selecting patients for testing.
 - ◊ **Testing Methodologies:** FISH break-apart probe methodology was the first methodology deployed widely. IHC can be deployed as an effective screening strategy. FDA-approved IHC can be utilized as a stand-alone test, not requiring confirmation by FISH. Numerous NGS methodologies can detect *ALK* fusions. Targeted real-time PCR assays are used in some settings, although it is unlikely to detect fusions with novel partners.
- ▶ **ROS1 (ROS proto-oncogene 1) Gene Rearrangements:** ROS1 is a receptor tyrosine kinase that can be rearranged in NSCLC, resulting in dysregulation and inappropriate signaling through the ROS1 kinase domain.
 - ◊ Numerous fusion partners are seen with *ROS1*, and common fusion partners include: *CD74*, *SLC34A2*, *CCDC6*, and *GOPC* (*FIG*).
 - ◊ The presence of a *ROS1* rearrangement is associated with responsiveness to oral ROS1 TKIs.
 - ◊ Some clinicopathologic features—such as smoking status and histology—have been associated with the presence of a *ROS1* rearrangement; however, these features should not be utilized in selecting patients for testing.
 - ◊ **Testing Methodologies:** FISH break-apart probe methodology can be deployed; however, it may under-detect the *FIG*-*ROS1* variant. IHC approaches can be deployed; however, IHC for *ROS1* fusions has low specificity, and follow-up confirmatory testing is a necessary component of utilizing ROS1 IHC as a screening modality. Numerous NGS methodologies can detect *ROS1* fusions, although DNA-based NGS may under-detect *ROS1* fusions. Targeted real-time PCR assays are utilized in some settings, although they are unlikely to detect fusions with novel partners.
- ▶ **BRAF (B-Raf proto-oncogene) Point Mutations:** BRAF is a serine/threonine kinase that is part of the canonical MAP/ERK signaling pathway. Activating mutations in BRAF result in unregulated signaling through the MAP/ERK pathway.
 - ◊ Mutations in *BRAF* can be seen in NSCLC. The presence of a specific mutation resulting in a change in amino acid position 600 (most commonly p.V600E, but rarely p.V600K, p.V600D, or other changes) has been associated with responsiveness to combined therapy with oral inhibitors of BRAF and MEK.
 - ◊ Note that other mutations in *BRAF* are observed in NSCLC, and the impact of those mutations on therapy selection is not well understood at this time.
 - ◊ **Testing Methodologies:** Real-time PCR, Sanger sequencing (ideally paired with tumor enrichment), and NGS are the most commonly deployed methodologies for examining *BRAF* mutation status. While an anti-BRAF p.V600E-specific monoclonal antibody is commercially available, and some studies have examined utilizing this approach, it should only be deployed after extensive validation.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

[Continued](#)**NSCL-H**
4 OF 8

**PRINCIPLES OF MOLECULAR AND BIOMARKER ANALYSIS**• **Molecular Targets for Analysis (continued)**

- ▶ **KRAS (KRAS proto-oncogene) point mutations:** KRAS is a G-protein with intrinsic GTPase activity, and activating mutations result in unregulated signaling through the MAP/ERK pathway.
 - ◊ Mutations in *KRAS* are most commonly seen at codon 12, although other mutations can be seen in NSCLC.
 - ◊ The presence of a *KRAS* mutation is prognostic of poor survival when compared to patients with tumors without *KRAS* mutation.
 - ◊ Owing to the low probability of overlapping targetable alterations, the presence of a known activating mutation in *KRAS* identifies patients who are unlikely to benefit from further molecular testing.
 - ◊ The presence of *KRAS* p.G12C is associated with responsiveness to an oral *KRAS* G12C inhibitor used for subsequent therapy, which was designed specifically for this mutation. Responsiveness to this class of inhibitor has not been prospectively evaluated with mutations other than *KRAS* p.G12C.
 - ◊ Testing methodologies: NGS, real-time PCR, and Sanger sequencing (ideally paired with tumor enrichment) are the most commonly deployed methodologies for examining *KRAS* mutation status.
- ▶ **MET (mesenchymal-epithelial transition) exon 14 (*METex14*) skipping variants:** MET is a receptor tyrosine kinase. A mutation that results in loss of exon 14 can occur in NSCLC. Loss of *METex14* leads to dysregulation and inappropriate signaling.
 - ◊ The presence of *METex14* skipping mutation is associated with responsiveness to oral MET TKIs.
 - ◊ A broad range of molecular alterations lead to *METex14* skipping.
 - ◊ Testing Methodologies: NGS-based testing is the primary method for detection of *METex14* skipping events; RNA-based NGS may have improved detection. IHC is not a method for detection of *METex14* skipping.
- ▶ **RET (rearranged during transfection) Gene Rearrangements:** RET is a receptor tyrosine kinase that can be rearranged in NSCLC, resulting in dysregulation and inappropriate signaling through the RET kinase domain.
 - ◊ Common fusion partners are *KIF5B*, *NCOA4*, and *CCDC6*; however, numerous other fusion partners have been identified.
 - ◊ The presence of a *RET* rearrangement is associated with responsiveness to oral RET TKIs regardless of fusion partner.
 - ◊ Testing Methodologies: FISH break-apart probe methodology can be deployed; however, it may under-detect some fusions. Targeted real-time reverse-transcriptase PCR assays are utilized in some settings, although they are unlikely to detect fusions with novel partners. NGS-based methodology has a high specificity, and RNA-based NGS is preferable to DNA-based NGS for fusion detection.
- ▶ **ERBB2 (*Erb-B2* Receptor Tyrosine Kinase 2)/HER2 Gene Mutations:** *ERBB2* encodes for HER2, a receptor tyrosine kinase found on the surface of normal epithelial cells that is often overexpressed or mutated in a variety of human malignancies.
 - ◊ A spectrum of genomic alterations involving *ERBB2* have been identified in multiple tumor types, including mutations and amplifications, and less commonly, other alterations.
 - ◊ Mutations in *ERBB2* are most commonly insertion/duplication events in exon 20, and less commonly, other activating mutations are observed.
 - ◊ Activating mutations in *ERBB2* are associated with responsiveness to anti-HER2 targeted therapy agents.
 - While some mutations are known to be activating, including mutations in the extracellular domain (eg, p.S310X) and exon 20 insertion/duplication mutations, not all single-nucleotide or double-nucleotide changes in *ERBB2* are activating and not all potentially activating mutations were studied in pivotal trials.
 - ◊ Some clinicopathologic features, such as smoking status and histology, are associated with the presence of *ERBB2* activating mutations; however, these features should not be utilized in selecting patients for testing.
 - ◊ Testing methodologies: While Sanger sequencing and targeted PCR techniques can be utilized to examine *ERBB2* mutations, the diversity and spectrum of mutations are best surveyed using NGS-based approaches.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

[Continued](#)

**PRINCIPLES OF MOLECULAR AND BIOMARKER ANALYSIS**• **Molecular Targets for Analysis (continued)**▶ ***NTRK1/2/3* (neurotrophic tyrosine receptor kinase) gene fusions**

- ◊ The presence of *NTRK1/2/3* gene fusions is associated with responsiveness to oral TRK inhibitors.
- ◊ *NTRK1/2/3* are tyrosine receptor kinases that are rarely rearranged in NSCLC as well as in other tumor types, resulting in dysregulation and inappropriate signaling.
- ◊ Numerous fusion partners have been identified.
- ◊ To date, no specific clinicopathologic features, other than absence of other driver alterations, have been identified in association with these fusions.
- ◊ Point mutations in *NTRK1/2/3* are generally non-activating and have not been studied in association with targeted therapy.
- ◊ Testing Methodologies: Various methodologies can be used to detect *NTRK1/2/3* gene fusions, including: FISH, IHC, PCR, and NGS; false negatives may occur. IHC methods are complicated by baseline expression in some tissues. FISH testing may require at least 3 probe sets for full analysis. NGS testing can detect a broad range of alterations. DNA-based NGS may under-detect *NTRK1* and *NTRK3* fusions.

• **PD-L1 (programmed death ligand 1):** PD-L1 is a co-regulatory molecule that can be expressed on tumor cells and inhibit T-cell-mediated cell death. T-cells express PD-1, a negative regulator, which binds to ligands including PD-L1 (CD274) or PD-L2 (CD273). In the presence of PD-L1, T-cell activity is suppressed.▶ **Checkpoint inhibitor antibodies block the PD-1 and PD-L1 interaction, thereby improving the antitumor effects of endogenous T cells.**▶ **IHC for PD-L1 can be utilized to identify disease most likely to respond to first-line anti PD-1/PD-L1.**

- ◊ Various antibody clones have been developed for IHC analysis of PD-L1 expression, and while several are comparable regarding intensity and proportion of cells stained, some are not.
 - The definition of positive and negative testing is dependent on the individual antibody, clone, and platform deployed, which may be unique to each checkpoint inhibitor therapy. The approval of multiple different assays for PD-L1 has raised concern among both pathologists and oncologists.
 - While some clones for PD-L1 IHC are FDA-approved for specific indications, use of multiple IHC tests is not routinely recommended, provided any individual IHC test has been internally validated for comparability for categorical results against the FDA-approved clone.
 - Interpretation of PD-L1 IHC in NSCLC is typically focused on the proportion of tumor cells expressing membranous staining at any level and therefore is a linear variable; scoring systems may be different in other tumor types.
- ◊ Although PD-L1 expression can be elevated in patients with an oncogenic driver, targeted therapy for the oncogenic driver should take precedence over treatment with an immune checkpoint inhibitor.

Note: All recommendations are category 2A unless otherwise indicated.**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.[Continued](#)**NSCL-H**
6 OF 8

**PRINCIPLES OF MOLECULAR AND BIOMARKER ANALYSIS**

- **Testing in the setting of a limited number of pulmonary nodules can aid in distinguishing separate primary lung carcinoma versus intrapulmonary metastatic disease.**
 - ▶ **Studies to explore tumor relatedness by testing tissue from separately sampled lesions using a broad gene coverage NGS approach suggest it may be superior to histopathologic assessment.**
 - ▶ **Tumor pairs exhibiting entirely non-overlapping, unique mutations are considered clonally unrelated separate primary lung cancers, even if histologically similar. Tumors that share multiple (≥2) mutations are more likely to be clonally related; however, this may depend on the extent to which any individual mutation is extremely common in NSCLC and whether identified alterations are driver or passenger alterations. Results in which no mutations or only one mutation are identified are not informative for this evaluation.**
- **Testing in the Setting of Progression on Targeted Therapy:**
 - ▶ **For any patient with progression on targeted therapy, histologic transformation (such as small cell) is a possible mechanism of resistance. Tissue biopsy of a progressing lesion should be considered to evaluate morphology and biomarker analysis.**
 - ▶ **For many of the above listed analytes, there is growing recognition of the molecular mechanisms of resistance to therapy. Re-testing of a sample from a tumor that is actively progressing while exposed to targeted therapy can shed light on appropriate next therapeutic steps:**
 - ◊ **For patients with an underlying *EGFR* sensitizing mutation who have been treated with EGFR TKI, minimum appropriate testing includes high-sensitivity evaluation for *p.T790M*; testing for alternate mechanisms of resistance (*MET* amplification, *ERBB2* amplification) may be used to direct patients for additional therapies. The presence of *p.T790M* can direct patients to third-generation EGFR TKI therapy.**
 - **Assays for the detection of *EGFR p.T790M* should be designed to have an analytic sensitivity of a minimum of 5% allelic fraction. The original sensitizing mutation can be utilized as an internal control in many assays to determine whether a *p.T790M* is within the range of detection if present as a subclonal event.**
 - ◊ **For patients with underlying *ALK* rearrangement who have been treated with ALK TKI, it is unclear whether identification of specific tyrosine kinase domain mutation can identify appropriate next steps in therapy, although some preliminary data suggest that specific kinase domain mutations may impact the next line of therapy.**
 - ◊ **Broad genomic profiling may be the most informative approach to examining potential mechanisms of resistance, which may require more than one instance of such profiling over the course of an individual patient's therapy. Assay methodology selection can impact the ability to identify subclonal events in this setting.**

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

[Continued](#)

NSCL-H
7 OF 8



PRINCIPLES OF MOLECULAR AND BIOMARKER ANALYSIS

- **Circulating Tumor DNA (ctDNA) Testing:**
 - ▶ **ctDNA testing should not be used in lieu of a histologic tissue diagnosis.**
 - ▶ **ctDNA is not routinely recommended in settings other than advanced/metastatic disease. For stages I–III, tissue-based testing is preferred. Metastatic disease confined to the thorax may have a higher yield with tissue-based testing.**
 - ▶ **Some laboratories offer testing for molecular alterations examining nucleic acids in peripheral circulation, most commonly in processed plasma.**
 - ▶ **Studies have demonstrated ctDNA and tissue testing to have very high specificity. Both ctDNA and tissue testing have appreciable false-negative rates, supporting the complementarity of these approaches, and data support complementary testing to reduce turnaround time and increase yield of targetable alteration detection.**
 - ◊ **Limitations of ctDNA testing that can impact interpretation include:**
 - **Low tumor fraction/ctDNA; some assays include a measure of ctDNA fraction, which can aid in identification of situations in which low ctDNA fraction might suggest compromised sensitivity**
 - **The presence of mutations from sites other than the target lesion, most commonly clonal hematopoiesis of indeterminate potential (CHIP) or post-chemotherapy marrow clones. *KRAS* and *TP53* can be seen in either of these circumstances**
 - **The inherent ability of the assay to detect fusions or other genomic variation of relevance**
 - ◊ **Limitations of tissue testing that can impact interpretation include:**
 - **Low tumor percent in a sample not sufficiently mitigated by tumor enrichment or high analytic sensitivity methods**
 - **The inherent ability of the assay to detect fusions or other genomic variation of relevance**

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

EMERGING BIOMARKERS TO IDENTIFY NOVEL THERAPIES FOR PATIENTS WITH METASTATIC NSCLC

Genetic Alteration (ie, Driver event)	Available Targeted Agents with Activity Against Driver Event in Lung Cancer
High-level <i>MET</i> amplification ^{a,b}	Capmatinib ¹ Tepotinib ² Crizotinib ^{3,4}

^a The definition of high-level *MET* amplification is evolving and may differ according to the assay used for testing. For NGS-based results, a copy number greater than 10 is consistent with high-level *MET* amplification.

^b In patients with *EGFR*-mutant NSCLC who develop high-level *MET* amplifications, administration of these agents with continuation of osimertinib is acceptable.

¹ Wolf J, Seto T, Han JY, et al; GEOMETRY mono-1 Investigators. Capmatinib in MET exon 14-mutated or MET-amplified non-small-cell lung cancer. *N Engl J Med* 2020;383:944-957.

² Le X, Paz-Ares LG, Van Meerbeeck J, et al. Tepotinib in patients with advanced non-small cell lung cancer (NSCLC) with *MET* amplification (*METamp*) [abstract]. *J Clin Oncol* 2021;39(Suppl):Abstract 9021.

³ Ou SH, Kwak EL, Siwak-Tapp C, et al. Activity of crizotinib (PF02341066), a dual mesenchymal-epithelial transition (MET) and anaplastic lymphoma kinase (ALK) inhibitor, in a non-small cell lung cancer patient with de novo MET amplification. *J Thorac Oncol* 2011;6:942-946.

⁴ Camidge DR, Otterson GA, Clark JW, et al. Crizotinib in patients with MET-amplified NSCLC. *J Thorac Oncol* 2021;16:1017-1029.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

MOLECULAR AND BIOMARKER-DIRECTED THERAPY FOR ADVANCED OR METASTATIC DISEASE^{a,b}

EGFR Exon 19 Deletion or Exon 21 L858R

- First-line therapy
 - Afatinib¹
 - Erlotinib²
 - Dacomitinib³
 - Gefitinib^{4,5}
 - Osimertinib⁶
 - Osimertinib + pemetrexed + (cisplatin or carboplatin) (nonsquamous)⁷
 - Erlotinib + ramucirumab⁸
 - Erlotinib + bevacizumab^c (nonsquamous)⁹
- Subsequent therapy
 - Osimertinib¹⁰
 - Amivantamab-vmjw + carboplatin + pemetrexed (nonsquamous)¹¹

EGFR S768I, L861Q, and/or G719X

- First-line therapy
 - Afatinib^{1,12}
 - Erlotinib²
 - Dacomitinib³
 - Gefitinib^{4,5}
 - Osimertinib^{6,13}
- Subsequent therapy
 - Osimertinib¹⁰
 - Amivantamab-vmjw + carboplatin + pemetrexed (nonsquamous)¹¹

EGFR Exon 20 Insertion Mutation

- First-line therapy
 - Amivantamab-vmjw + carboplatin + pemetrexed (nonsquamous)¹⁴
- Subsequent therapy
 - Amivantamab-vmjw¹⁵

KRAS G12C Mutation^d

- Subsequent therapy
 - Sotorasib¹⁶
 - Adagrasib¹⁷

ALK Rearrangement

- First-line therapy
 - Alectinib^{18,19}
 - Brigatinib²⁰
 - Ceritinib²¹
 - Crizotinib^{18,22}
 - Lorlatinib²³
- Subsequent therapy
 - Alectinib^{24,25}
 - Brigatinib²⁶
 - Ceritinib²⁷
 - Lorlatinib²⁸

ROS1 Rearrangement

- First-line therapy
 - Ceritinib²⁹
 - Crizotinib³⁰
 - Entrectinib³¹
 - Repotrectinib³²
- Subsequent therapy
 - Lorlatinib³³
 - Entrectinib³¹
 - Repotrectinib³²

BRAF V600E Mutation

- First-line therapy
 - Dabrafenib/trametinib³⁴
 - Encorafenib/binimetinib³⁵
 - Dabrafenib³⁶
 - Vemurafenib
- Subsequent therapy
 - Dabrafenib/trametinib^{36,37}
 - Encorafenib/binimetinib³⁵

NTRK1/2/3 Gene Fusion

- First-line/Subsequent therapy
 - Larotrectinib³⁸
 - Entrectinib³⁹

MET Exon 14 Skipping Mutation^d

- First-line therapy/Subsequent therapy
 - Capmatinib⁴⁰
 - Crizotinib⁴¹
 - Tepotinib⁴²

RET Rearrangement^d

- First-line therapy/Subsequent therapy
 - Selpercatinib⁴³
 - Pralsetinib⁴⁴
 - Cabozantinib^{45,46}

ERBB2 (HER2) Mutation^d

- Subsequent therapy
 - Fam-trastuzumab deruxtecan-rxki⁴⁷
 - Ado-trastuzumab emtansine⁴⁸

[PD-L1 ≥50% First-line Therapy](#)

[PD-L1 ≥1%–49% First-line Therapy](#)

^a Monitoring During Initial Therapy: Response assessment after 2 cycles, then every 2–4 cycles with CT of known or high-risk sites of disease with or without contrast or when clinically indicated. Timing of CT scans within Guidelines parameters is a clinical decision.

^b Monitoring During Subsequent Therapy or Maintenance Therapy: Response assessment with CT of known or high-risk sites of disease with or without contrast every 6–12 weeks. Timing of CT scans within Guidelines parameters is a clinical decision.

^c An FDA-approved biosimilar is an appropriate substitute for bevacizumab.

^d For agents with a similar mechanism of action, it is not recommended to switch between these drugs at the time of progression.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

References



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

MOLECULAR AND BIOMARKER-DIRECTED THERAPY FOR ADVANCED OR METASTATIC DISEASE^{a,b}

PD-L1 ≥50% FIRST-LINE THERAPY (PS 0–2)	
Adenocarcinoma, Large Cell, NSCLC NOS	Squamous Cell Carcinoma
<u>Preferred</u> <ul style="list-style-type: none"> • Pembrolizumab (category 1)^{49,50} • (Carboplatin or cisplatin) + pemetrexed + pembrolizumab (category 1)^{51,52} • Atezolizumab (category 1)⁵³ • Cemiplimab-rwlc (category 1)⁵⁴ • Cemiplimab-rwlc + pemetrexed + (carboplatin or cisplatin) (category 1)⁵⁵ <u>Other Recommended</u> <ul style="list-style-type: none"> • Carboplatin + paclitaxel + bevacizumab^{c,e} + atezolizumab (category 1)⁵⁶ • Carboplatin + albumin-bound paclitaxel + atezolizumab⁵⁷ • Nivolumab + ipilimumab + pemetrexed + (carboplatin or cisplatin) (category 1)⁵⁸ • Cemiplimab-rwlc + paclitaxel + (carboplatin or cisplatin) (category 1)⁵⁵ • Tremelimumab-actl + durvalumab + carboplatin + albumin-bound paclitaxel (category 2B)⁵⁹ • Tremelimumab-actl + durvalumab + (carboplatin or cisplatin) + pemetrexed (category 2B)⁵⁹ <u>Useful in Certain Circumstances</u> <ul style="list-style-type: none"> • Nivolumab + ipilimumab (category 1)⁶⁰ 	<u>Preferred</u> <ul style="list-style-type: none"> • Pembrolizumab (category 1)^{49,50} • Carboplatin + (paclitaxel or albumin-bound paclitaxel) + pembrolizumab (category 1)⁶¹ • Atezolizumab (category 1)⁵³ • Cemiplimab-rwlc (category 1)⁵⁴ • Cemiplimab-rwlc + paclitaxel + (carboplatin or cisplatin) (category 1)⁵⁵ <u>Other Recommended</u> <ul style="list-style-type: none"> • Nivolumab + ipilimumab + paclitaxel + carboplatin (category 1)⁵⁸ • Tremelimumab-actl + durvalumab + carboplatin + albumin-bound paclitaxel (category 2B)⁵⁹ • Tremelimumab-actl + durvalumab + (carboplatin or cisplatin) + gemcitabine (category 2B)⁵⁹ <u>Useful in Certain Circumstances</u> <ul style="list-style-type: none"> • Nivolumab + ipilimumab (category 1)⁶⁰
PD-L1 ≥50% FIRST-LINE THERAPY (PS 3–4) ^f	
Best supportive care (NCCN Guidelines for Palliative Care)	

PD-L1 ≥1%–49% First-line Therapy Continuation Maintenance

^a Monitoring During Initial Therapy: Response assessment after 2 cycles, then every 2–4 cycles with CT of known or high-risk sites of disease with or without contrast or when clinically indicated. Timing of CT scans within Guidelines parameters is a clinical decision.

^b Monitoring During Subsequent Therapy or Maintenance Therapy: Response assessment with CT of known or high-risk sites of disease with or without contrast every 6–12 weeks. Timing of CT scans within Guidelines parameters is a clinical decision.

^c An FDA-approved biosimilar is an appropriate substitute for bevacizumab.

^e Criteria for treatment with bevacizumab: nonsquamous NSCLC, and no recent history of hemoptysis.

^f Atezolizumab monotherapy is a treatment option for patients with PS 3, regardless of PD-L1 status.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

References



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

MOLECULAR AND BIOMARKER-DIRECTED THERAPY FOR ADVANCED OR METASTATIC DISEASE^{a,b}

PD-L1 ≥1%–49% FIRST-LINE THERAPY (PS 0–2)	
Adenocarcinoma, Large Cell, NSCLC NOS	Squamous Cell Carcinoma
<u>Preferred</u> <ul style="list-style-type: none"> • (Carboplatin or cisplatin) + pemetrexed + pembrolizumab (category 1)^{51,52} • Cemiplimab-rwlc + pemetrexed + (carboplatin or cisplatin) (category 1)⁵⁵ <u>Other Recommended</u> <ul style="list-style-type: none"> • Carboplatin + paclitaxel + bevacizumab^{c,e} + atezolizumab (category 1)⁵⁶ • Carboplatin + albumin-bound paclitaxel + atezolizumab⁵⁷ • Nivolumab + ipilimumab + pemetrexed + (carboplatin or cisplatin) (category 1)⁵⁸ • Nivolumab + ipilimumab (category 1)⁶⁰ • Cemiplimab-rwlc + paclitaxel + (carboplatin or cisplatin) (category 1)⁵⁵ • Tremelimumab-actl + durvalumab + carboplatin + albumin-bound paclitaxel (category 1)⁵⁹ • Tremelimumab-actl + durvalumab + (carboplatin or cisplatin) + pemetrexed (category 1)⁵⁹ <u>Useful in Certain Circumstances</u> <ul style="list-style-type: none"> • Pembrolizumab (category 2B)^{9,49,50} 	<u>Preferred</u> <ul style="list-style-type: none"> • Carboplatin + (paclitaxel or albumin-bound paclitaxel) + pembrolizumab (category 1)⁶¹ • Cemiplimab-rwlc + paclitaxel + (carboplatin or cisplatin) (category 1)⁵⁵ <u>Other Recommended</u> <ul style="list-style-type: none"> • Nivolumab + ipilimumab + paclitaxel + carboplatin (category 1)⁵⁸ • Nivolumab + ipilimumab (category 1)⁶⁰ • Tremelimumab-actl + durvalumab + carboplatin + albumin-bound paclitaxel⁵⁹ • Tremelimumab-actl + durvalumab + (carboplatin or cisplatin) + gemcitabine⁵⁹ <u>Useful in Certain Circumstances</u> <ul style="list-style-type: none"> • Pembrolizumab (category 2B)^{9,49,50}
PD-L1 ≥1%–49% FIRST-LINE THERAPY (PS 3–4) ^f	
Best supportive care (NCCN Guidelines for Palliative Care)	

[PD-L1 ≥50% First-line Therapy](#)

[Continuation Maintenance](#)

^a Monitoring During Initial Therapy: Response assessment after 2 cycles, then every 2–4 cycles with CT of known or high-risk sites of disease with or without contrast or when clinically indicated. Timing of CT scans within Guidelines parameters is a clinical decision.

^b Monitoring During Subsequent Therapy or Maintenance Therapy: Response assessment with CT of known or high-risk sites of disease with or without contrast every 6–12 weeks. Timing of CT scans within Guidelines parameters is a clinical decision.

^c An FDA-approved biosimilar is an appropriate substitute for bevacizumab.

^e Criteria for treatment with bevacizumab: nonsquamous NSCLC, and no recent history of hemoptysis.

^f Atezolizumab monotherapy is a treatment option for patients with PS 3, regardless of PD-L1 status.

^g Pembrolizumab monotherapy can be considered in PD-L1 1%–49%, when there are contraindications to combination chemotherapy.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

[References](#)

**NCCN Guidelines Version 3.2024**
Non-Small Cell Lung Cancer**MOLECULAR AND BIOMARKER-DIRECTED THERAPY FOR ADVANCED OR METASTATIC DISEASE^b**

PD-L1 ≥50% CONTINUATION MAINTENANCE	
Adenocarcinoma, Large Cell, NSCLC NOS	Squamous Cell Carcinoma
<ul style="list-style-type: none"> • Pembrolizumab (category 1)^h • Pembrolizumab + pemetrexed (category 1)ⁱ • Atezolizumab and bevacizumab^{c,e} (category 1)^j • Atezolizumab^k • Nivolumab + ipilimumab (category 1)^l • Cemiplimab-rwlc (category 1)^m • Cemiplimab-rwlcⁿ ± pemetrexed^o (category 1) • Durvalumab^p ± pemetrexed^q 	<ul style="list-style-type: none"> • Pembrolizumab (category 1)^{h,r} • Atezolizumab^k • Nivolumab + ipilimumab (category 1)^l • Cemiplimab-rwlc (category 1)^{m,n} • Durvalumab^p
PD-L1 ≥1%–49% CONTINUATION MAINTENANCE	
Adenocarcinoma, Large Cell, NSCLC NOS	Squamous Cell Carcinoma
<ul style="list-style-type: none"> • Pembrolizumab (category 2B)^h • Pembrolizumab + pemetrexed (category 1)ⁱ • Atezolizumab and bevacizumab^{c,e} (category 1)^j • Atezolizumab^k • Nivolumab + ipilimumab (category 1)^l • Cemiplimab-rwlcⁿ ± pemetrexed^o (category 1) • Durvalumab^p ± pemetrexed^q 	<ul style="list-style-type: none"> • Pembrolizumab^{h,r} • Nivolumab + ipilimumab (category 1)^l • Cemiplimab-rwlc (category 1)ⁿ • Durvalumab^p

[References](#)

^b Monitoring During Subsequent Therapy or Maintenance Therapy: Response assessment with CT of known or high-risk sites of disease with or without contrast every 6–12 weeks. Timing of CT scans within Guidelines parameters is a clinical decision.

^c An FDA-approved biosimilar is an appropriate substitute for bevacizumab.

^e Criteria for treatment with bevacizumab: nonsquamous NSCLC, and no recent history of hemoptysis.

^h If pembrolizumab monotherapy given.

ⁱ If pembrolizumab/carboplatin/pemetrexed or pembrolizumab/cisplatin/pemetrexed given.

^j If atezolizumab/carboplatin/paclitaxel/bevacizumab given.

^k If atezolizumab/carboplatin/albumin-bound paclitaxel given (category 1 following atezolizumab alone).

^l If nivolumab + ipilimumab ± chemotherapy given.

^m If cemiplimab-rwlc monotherapy given.

ⁿ If cemiplimab-rwlc combination therapy given.

^o If cemiplimab-rwlc + pemetrexed + (carboplatin or cisplatin) given.

^p If tremelimumab-actl combination therapy given. Refer to categories of evidence for first-line regimen.

^q If tremelimumab-actl + durvalumab + (carboplatin or cisplatin) + pemetrexed given. Refer to categories of evidence for first-line regimen.

^r If pembrolizumab/carboplatin/(paclitaxel or albumin-bound paclitaxel) given.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

**MOLECULAR AND BIOMARKER-DIRECTED THERAPY FOR ADVANCED OR METASTATIC DISEASE – REFERENCES**

- 1 Yang JC, et al. Afatinib versus cisplatin based chemotherapy for EGFR mutation positive lung adenocarcinoma (LUX Lung 3 and LUX Lung 6): analysis of overall survival data from two randomised, phase 3 trials. *Lancet Oncol* 2015;16:141-151.
- 2 Rosell R, et al. Erlotinib versus standard chemotherapy as first line treatment for European patients with advanced EGFR mutation positive non small cell lung cancer (EORTAC): a multicentre, open label, randomised phase 3 trial. *Lancet Oncol* 2012;13:239-246.
- 3 Wu Y-L, et al. Dacomitinib versus gefitinib as first-line treatment for patients with EGFR-mutation-positive non-small-cell lung cancer (ARCHER 1050): a randomized, open-label, phase 3 trial. *Lancet Oncol* 2017;18:1454-1466.
- 4 Mok TS, et al. Gefitinib or carboplatin paclitaxel in pulmonary adenocarcinoma. *N Engl J Med* 2009;361:947-957.
- 5 Douillard JY, et al. First line gefitinib in Caucasian EGFR mutation positive NSCLC patients: a phase IV, open label, single arm study. *Br J Cancer* 2014;110:55-62.
- 6 Soria JC, et al. Osimertinib in untreated EGFR-mutated advanced non-small-cell lung cancer. *N Engl J Med* 2018;378:113-125.
- 7 Planchard D, et al. Osimertinib with or without chemotherapy in EGFR-mutated advanced NSCLC. *N Engl J Med* 2023;389:1935-1948.
- 8 Nakagawa K, et al. Ramucirumab plus erlotinib in patients with untreated, EGFR-mutated, advanced non-small-cell lung cancer (RELAY): a randomised, double-blind, placebo-controlled, phase 3 trial. *Lancet Oncol* 2019;20:1655-1669.
- 9 Kawashima Y, et al. Bevacizumab plus erlotinib versus erlotinib alone in Japanese patients with advanced, metastatic, EGFR-mutant non-small-cell lung cancer (NEJ026): overall survival analysis of an open-label, randomised, multicentre, phase 3 trial. *Lancet Respir Med* 2022;10:72-82.
- 10 Mok TS, et al. Osimertinib or platinum pemetrexed in EGFR T790M positive lung cancer. *N Engl J Med* 2017;376:629-640.
- 11 Passaro A, et al. Amivantamab plus chemotherapy with and without lazertinib in EGFR-mutant advanced NSCLC after disease progression on osimertinib: primary results from the phase III MARIPOSA-2 study. *Ann Oncol* 2023;S0923-7534:04281-04283.
- 12 Yang JC, et al. Clinical activity of afatinib in patients with advanced non-small-cell lung cancer harbouring uncommon EGFR mutations: a combined post-hoc analysis of LUX-Lung 2, LUX-Lung 3, and LUX-Lung 6. *Lancet Oncol* 2015;16:830-838.
- 13 Cho JH, et al. Osimertinib for patients with non-small-cell lung cancer harboring uncommon EGFR mutations: a multicenter, open-label, phase II trial (KCSG-LU15-09). *J Clin Oncol* 2020;38:488-495.
- 14 Zhou C, et al. Amivantamab plus chemotherapy in NSCLC with EGFR Exon 20 insertions. *N Engl J Med* 2023;389:2039-2051.
- 15 Park K, et al. Amivantamab in EGFR exon 20 insertion-mutated non-small cell lung cancer progressing on platinum chemotherapy: initial results from the CHRYSALIS phase I study. *J Clin Oncol* 2021;39:3391-3402.
- 16 Skoulidis F, et al. Sotorasib for lung cancers with KRAS p.G12C mutation. *N Engl J Med* 2021;384:2371-2381.
- 17 Janne PA, et al. Adagrasib in non-small-cell lung cancer harboring a KRAS G12C mutation. *N Engl J Med* 2022;387:120-131.
- 18 Peters S, et al. Alectinib versus crizotinib in untreated ALK-positive non-small-cell lung cancer. *N Engl J Med* 2017;377:829-838.
- 19 Hida T, et al. Alectinib versus crizotinib in patients with ALK-positive non-small-cell lung cancer (J-ALEX): an open-label, randomised phase 3 trial. *Lancet* 2017;390:29-39.
- 20 Camidge DR, et al. Brigatinib versus crizotinib in ALK-positive non-small-cell lung cancer. *N Engl J Med* 2018;379:2027-2039.
- 21 Soria JC, et al. First line ceritinib versus platinum based chemotherapy in advanced ALK rearranged non small cell lung cancer (ASCEND 4): a randomised, open label, phase 3 study. *Lancet* 2017;389:917-929.
- 22 Solomon BJ, et al. First line crizotinib versus chemotherapy in ALK positive lung cancer. *N Engl J Med* 2014;371:2167-2177.
- 23 Shaw AT, et al. First-line lorlatinib or crizotinib in advanced ALK-positive lung cancer. *N Engl J Med* 2020;383:2018-2029.
- 24 Ou SI, et al. Alectinib in crizotinib refractory ALK rearranged non small cell lung cancer: a phase II global study. *J Clin Oncol* 2016;34:661-668.
- 25 Shaw AT, et al. Alectinib in ALK positive, crizotinib resistant, non small cell lung cancer: a single group, multicentre, phase 2 trial. *Lancet Oncol* 2016;17:234-242.
- 26 Kim DW, et al. Brigatinib in patients with crizotinib refractory anaplastic lymphoma kinase positive non small cell lung cancer: a randomized, multicenter phase II trial. *J Clin Oncol* 2017;35:2490-2498.
- 27 Shaw AT, et al. Ceritinib versus chemotherapy in patients with ALK-rearranged non-small-cell lung cancer previously given chemotherapy and crizotinib (ASCEND-5): a randomised, controlled, open-label, phase 3 trial. *Lancet Oncol* 2017;18:874-86.
- 28 Solomon BJ, et al. Lorlatinib in patients with ALK-positive non-small-cell lung cancer: results from a global phase 2 study. *Lancet Oncol* 2018;19:1654-1667.
- 29 Lim SM, et al. Open-label, multicenter, phase II study of ceritinib in patients with non-small-cell lung cancer harboring ROS1 rearrangement. *J Clin Oncol* 2017;35:2613-2618.
- 30 Shaw AT, et al. Crizotinib in ROS1-rearranged non-small-cell lung cancer. *N Engl J Med* 2014;371:1963-1971.

[Continued](#)**Note: All recommendations are category 2A unless otherwise indicated.****Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.**

**MOLECULAR AND BIOMARKER-DIRECTED THERAPY FOR ADVANCED OR METASTATIC DISEASE – REFERENCES**

- ³¹ Drilon A, et al. Entrectinib in ROS1 fusion-positive non-small-cell lung cancer: integrated analysis of three phase 1-2 trials. *Lancet Oncol* 2020;21:261-270.
- ³² Drilon A, et al. Repotrectinib in ROS1 fusion-positive non-small-cell lung cancer. *N Engl J Med* 2024;390:118-131.
- ³³ Shaw AT, et al. Lorlatinib in advanced ROS1-positive non-small-cell lung cancer: a multicentre, open-label, single-arm, phase 1-2 trial. *Lancet Oncol* 2019;20:1691-1701.
- ³⁴ Planchard D, et al. Dabrafenib plus trametinib in patients with previously untreated BRAF(V600E)-mutant metastatic non-small-cell lung cancer: an open-label, phase 2 trial. *Lancet Oncol* 2017;18:1307-1316.
- ³⁵ Riely GJ, Smit EF, Ahn M-J, et al. Phase II, open-label study of encorafenib plus binimetinib in patients with BRAF V600-mutant metastatic non-small-cell lung cancer. *J Clin Oncol* 2022;41:3700-3711.
- ³⁶ Planchard D, et al. Phase 2 study of dabrafenib plus trametinib in patients with BRAF V600E-mutant metastatic NSCLC: updated 5-year survival rates and genomic analysis. *J Thorac Oncol* 2022;17:103-115.
- ³⁷ Planchard D, et al. Dabrafenib plus trametinib in patients with previously treated BRAF(V600E) mutant metastatic non small cell lung cancer: an open label, multicentre phase 2 trial. *Lancet Oncol* 2016;17:984-993.
- ³⁸ Drilon A, et al. Efficacy of larotrectinib in TRK fusion-positive cancers in adults and children. *N Engl J Med* 2018;378:731-739.
- ³⁹ Doebele RC, et al. Entrectinib in patients with advanced or metastatic NTRK fusion-positive solid tumours: integrated analysis of three phase 1-2 trials. *Lancet Oncol* 2020;21:271-282.
- ⁴⁰ Wolf J, et al; GEOMETRY mono-1 Investigators. Capmatinib in MET exon 14-mutated or MET-amplified non-small-cell lung cancer. *N Engl J Med* 2020;383:944-957.
- ⁴¹ Drilon A, et al. Antitumor activity of crizotinib in lung cancers harboring a MET exon 14 alteration. *Nat Med* 2020;26:47-51.
- ⁴² Paik PK, et al. Tepotinib in non-small-cell lung cancer with MET Exon 14 skipping mutations. *N Engl J Med* 2020;383:931-943.
- ⁴³ Drilon A, et al. Efficacy of selpercatinib in RET fusion-positive non-small-cell lung cancer. *N Engl J Med* 2020;383:813-824.
- ⁴⁴ Gainor JF, et al. Pralsetinib for RET fusion-positive non-small cell lung cancer (ARROW): a multi-cohort, open-label, phase 1/2 study. *Lancet Oncol* 2021;22:959-969.
- ⁴⁵ Drilon A, et al. Response to cabozantinib in patients with RET fusion-positive lung adenocarcinomas. *Cancer Discov* 2013;3:630-635.
- ⁴⁶ Drilon A, et al. Cabozantinib in patients with advanced RET-rearranged non-small-cell lung cancer: an open-label, single-centre, phase 2, single-arm trial. *Lancet Oncol* 2016;17:1653-1660.
- ⁴⁷ Li BT, et al. Trastuzumab deruxtecan in HER2-mutant non-small-cell lung cancer. *N Engl J Med* 2022;386:241-251.
- ⁴⁸ Li BT, et al. Ado-trastuzumab emtansine in patients with HER2 mutant lung cancers: Results from a phase II basket trial. *J Clin Oncol* 2018;36:2532-2537.
- ⁴⁹ Reck M, et al. Pembrolizumab versus chemotherapy for PD L1 positive non small cell lung cancer. *N Engl J Med* 2016;375:1823-1833.
- ⁵⁰ Mok TSK, et al. Pembrolizumab versus chemotherapy for previously untreated, PD-L1-expressing, locally advanced or metastatic non-small-cell lung cancer (Keynote-042): a randomized open-label, controlled, phase 3 trial. *Lancet* 2019;393:1819-1830.
- ⁵¹ Gandhi L, et al. Pembrolizumab plus chemotherapy in metastatic non-small-cell lung cancer. *N Engl J Med* 2018;378:2078-2092.
- ⁵² Langer CJ, et al. Carboplatin and pemetrexed with or without pembrolizumab for advanced, non squamous non small cell lung cancer: a randomised, phase 2 cohort of the open label KEYNOTE 021 study. *Lancet Oncol* 2016;17:1497-1508.
- ⁵³ Herbst RS, et al. Atezolizumab for first-line treatment of PD-L1-selected patients with NSCLC. *N Engl J Med* 2020;383:1328-1339.
- ⁵⁴ Sezer A, et al. Cemiplimab monotherapy for first-line treatment of advanced non-small-cell lung cancer with PD-L1 of at least 50%: a multicentre, open-label, global, phase 3, randomised, controlled trial. *Lancet* 2021;397:592-604.
- ⁵⁵ Gogishvili M, et al. Cemiplimab plus chemotherapy versus chemotherapy alone in non-small cell lung cancer: a randomized, controlled, double-blind phase 3 trial. *Nat Med* 2022;28:2374-2380.
- ⁵⁶ Socinski M, et al. Atezolizumab for first-line treatment of metastatic nonsquamous NSCLC. *N Engl J Med* 2018;378:2288-2301.
- ⁵⁷ West H, et al. Atezolizumab in combination with carboplatin plus nab-paclitaxel chemotherapy compared with chemotherapy alone as first-line treatment for metastatic non-squamous non-small-cell lung cancer (IMpower130): a multicentre, randomised, open-label, phase 3 trial. *Lancet Oncol* 2019;20:924-937.
- ⁵⁸ Paz-Ares L, et al. First-line nivolumab plus ipilimumab combined with two cycles of chemotherapy in patients with non-small-cell lung cancer (CheckMate 9LA): an international, randomised, open-label, phase 3 trial. *Lancet Oncol* 2021;22:198-211.
- ⁵⁹ Johnson ML, et al. Durvalumab with or without tremelimumab in combination with chemotherapy as first-line therapy for metastatic non-small-cell lung cancer: the phase III POSEIDON study. *J Clin Oncol* 2023;41:1213-1227.
- ⁶⁰ Hellmann MD, et al. Nivolumab plus ipilimumab in advanced non-small-cell lung cancer. *N Engl J Med* 2019;381:2020-2031.
- ⁶¹ Paz-Ares L, et al. Pembrolizumab plus chemotherapy for squamous non-small-cell lung cancer. *N Engl J Med* 2018;379:2040-2051.

Note: All recommendations are category 2A unless otherwise indicated.**Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.**



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

SYSTEMIC THERAPY FOR ADVANCED OR METASTATIC DISEASE^{a,b,c}

ADENOCARCINOMA, LARGE CELL, NSCLC NOS (PS 0–1)	
No contraindications to PD-1 or PD-L1 inhibitors ^d	Contraindications to PD-1 or PD-L1 inhibitors ^d
Preferred <ul style="list-style-type: none"> Pembrolizumab/carboplatin/pemetrexed (category 1)^{1,2,e} Pembrolizumab/cisplatin/pemetrexed (category 1)^{2,e} Cemiplimab-rwlc/pemetrexed/(carboplatin or cisplatin) (category 1)^{7,e} Other Recommended <ul style="list-style-type: none"> Atezolizumab/carboplatin/paclitaxel/bevacizumab^e (category 1)^{3,f,g,h,i} Atezolizumab/carboplatin/albumin-bound paclitaxel^{4,e} Nivolumab/ipilimumab^{5,e} Nivolumab/ipilimumab/pemetrexed/(carboplatin or cisplatin) (category 1)^{6,e} Cemiplimab-rwlc/paclitaxel/(carboplatin or cisplatin) (category 1)^{7,e} Tremelimumab-actl/durvalumab/carboplatin/albumin-bound paclitaxel^{8,e} (category 1) Tremelimumab-actl/durvalumab/(carboplatin or cisplatin)/pemetrexed^{8,e} (category 1) 	Useful in Certain Circumstances <ul style="list-style-type: none"> Bevacizumab^f/carboplatin/paclitaxel (category 1)^{9,g,h,i} Bevacizumab^f/carboplatin/pemetrexed^{9,10,g,h,i} Bevacizumab^f/cisplatin/pemetrexed^{11,g,h,i} Carboplatin-combination therapy (category 1) <ul style="list-style-type: none"> Combination options include: albumin-bound paclitaxel¹² docetaxel,¹³ etoposide,^{14,15} gemcitabine,¹⁶ paclitaxel,¹⁷ or pemetrexed¹⁸ Cisplatin-combination therapy (category 1) <ul style="list-style-type: none"> Combinations options include: docetaxel,¹³ etoposide,¹⁹ gemcitabine,^{17,20} paclitaxel,²¹ or pemetrexed²⁰ Gemcitabine/docetaxel (category 1)²² Gemcitabine/vinorelbine (category 1)²³
ADENOCARCINOMA, LARGE CELL, NSCLC NOS (PS 2)	
Preferred <ul style="list-style-type: none"> Carboplatin/pemetrexed¹⁸ Other Recommended <ul style="list-style-type: none"> Carboplatin/albumin-bound paclitaxel^{25,26} Carboplatin/docetaxel¹³ Carboplatin/etoposide^{14,15} Carboplatin/gemcitabine¹⁶ Carboplatin/paclitaxel¹⁷ 	Useful in Certain Circumstances <ul style="list-style-type: none"> Albumin-bound paclitaxel²⁴ Docetaxel^{27,28} Gemcitabine²⁹⁻³¹ Gemcitabine/docetaxel²² Gemcitabine/vinorelbine²³ Paclitaxel³²⁻³⁴ Pemetrexed³⁵
ADENOCARCINOMA, LARGE CELL, NSCLC NOS (PS 3–4) ^j	
Best supportive care (NCCN Guidelines for Palliative Care)	

^a Albumin-bound paclitaxel may be substituted for either paclitaxel or docetaxel in patients who have experienced hypersensitivity reactions after receiving paclitaxel or docetaxel despite premedication, or for patients where the standard premedications (ie, dexamethasone, H2 blockers, H1 blockers) are contraindicated.

^b Carboplatin-based regimens are often used for patients with comorbidities or those who cannot tolerate cisplatin.

^c If first-line systemic therapy is completed before treatment for an actionable mutation, and disease has progressed, see Subsequent Therapy ([NSCL-K 4 of 5](#)).

^d Contraindications for treatment with PD-1/PD-L1 inhibitors may include active or previously documented autoimmune disease and/or current use of immunosuppressive agents; some oncogenic drivers (ie, *EGFR* exon 19 deletion or L858R, *ALK* rearrangements) have been shown to be associated with less benefit from PD-1/PD-L1 inhibitors.

^e If progression on PD-1/PD-L1 inhibitor, using a PD-1/PD-L1 inhibitor is not recommended.

^f An FDA-approved biosimilar is an appropriate substitute for bevacizumab.

^g Bevacizumab should be given until progression.

^h Any regimen with a high risk of thrombocytopenia and the potential risk of bleeding should be used with caution in combination with bevacizumab.

ⁱ Criteria for treatment with bevacizumab: nonsquamous NSCLC, and no recent history of hemoptysis. Bevacizumab should not be given as a single agent, unless as maintenance if initially used with chemotherapy.

^j Atezolizumab monotherapy is a treatment option for patients with PS 3, regardless of PD-L1 status.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

[Continued](#)

[Maintenance Therapy NSCL-K \(3 of 5\)](#)

[Subsequent Therapy NSCL-K \(4 of 5\)](#)

[References](#)

NSCL-K
1 OF 5



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

SYSTEMIC THERAPY FOR ADVANCED OR METASTATIC DISEASE^{a,b,c}

SQUAMOUS CELL CARCINOMA (PS 0–1)	
No contraindications to PD-1 or PD-L1 inhibitors ^d	Contraindications to PD-1 or PD-L1 inhibitors ^d
Preferred <ul style="list-style-type: none"> Pembrolizumab/carboplatin/paclitaxel (category 1)^{36,e} Pembrolizumab/carboplatin/albumin-bound paclitaxel (category 1)^{36,e} Cemiplimab-rwlc/paclitaxel/(carboplatin or cisplatin) (category 1)^{7,e} Other Recommended <ul style="list-style-type: none"> Nivolumab/ipilimumab^{5,e} Nivolumab/ipilimumab/paclitaxel/carboplatin (category 1)^{6,e} Tremelimumab-actl/durvalumab/carboplatin/albumin-bound paclitaxel^{8,e} (category 1) Tremelimumab-actl/durvalumab/(carboplatin or cisplatin)/gemcitabine^{8,e} (category 1) 	Useful in Certain Circumstances <ul style="list-style-type: none"> Carboplatin/albumin-bound paclitaxel (category 1)¹⁰ Carboplatin/docetaxel (category 1)¹³ Carboplatin/gemcitabine (category 1)¹⁶ Carboplatin/paclitaxel (category 1)¹⁷ Cisplatin/docetaxel (category 1)¹³ Cisplatin/etoposide (category 1)¹⁹ Cisplatin/gemcitabine (category 1)^{19,20} Cisplatin/paclitaxel (category 1)²¹ Gemcitabine/docetaxel (category 1)²² Gemcitabine/vinorelbine (category 1)²³
SQUAMOUS CELL CARCINOMA (PS 2)	
Preferred <ul style="list-style-type: none"> Carboplatin/albumin-bound paclitaxel^{25,26} Carboplatin/gemcitabine¹⁶ Carboplatin/paclitaxel¹⁷ Other Recommended <ul style="list-style-type: none"> Carboplatin/docetaxel¹³ Carboplatin/etoposide^{14,15} 	Useful in Certain Circumstances <ul style="list-style-type: none"> Albumin-bound paclitaxel²⁴ Docetaxel^{27,28} Gemcitabine²⁹⁻³¹ Gemcitabine/docetaxel²² Gemcitabine/vinorelbine²³ Paclitaxel³²⁻³⁴
SQUAMOUS CELL CARCINOMA (PS 3–4) ^j	
Best supportive care (NCCN Guidelines for Palliative Care)	

^a Albumin-bound paclitaxel may be substituted for either paclitaxel or docetaxel in patients who have experienced hypersensitivity reactions after receiving paclitaxel or docetaxel despite premedication, or for patients where the standard premedications (ie, dexamethasone, H2 blockers, H1 blockers) are contraindicated.

^b Carboplatin-based regimens are often used for patients with comorbidities or those who cannot tolerate cisplatin.

^c If first-line systemic therapy completed before treatment for an actionable mutation, and disease has progressed, see Subsequent Therapy ([NSCL-K 4 of 5](#)).

^d Contraindications for treatment with PD-1/PD-L1 inhibitors may include active or previously documented autoimmune disease and/or current use of immunosuppressive agents; some oncogenic drivers (ie, *EGFR* exon 19 deletion or L858R, *ALK* rearrangements) have been shown to be associated with less benefit from PD-1/PD-L1 inhibitors.

^e If progression on PD-1/PD-L1 inhibitor, using a PD-1/PD-L1 inhibitor is not recommended.

^j Atezolizumab monotherapy is a treatment option for patients with PS 3, regardless of PD-L1 status.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

[Continued](#)

[Maintenance Therapy NSCL-K \(3 of 5\)](#)

[Subsequent Therapy NSCL-K \(4 of 5\)](#)

[References](#)

NSCL-K
2 OF 5



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

SYSTEMIC THERAPY FOR ADVANCED OR METASTATIC DISEASE – MAINTENANCE^k

Maintenance Therapy

- Continuation maintenance refers to the use of at least one of the agents given in first line, beyond 4–6 cycles, in the absence of disease progression. Switch maintenance refers to the initiation of a different agent, not included as part of the first-line regimen, in the absence of disease progression, after 4–6 cycles of initial therapy.
- Patients should receive maintenance therapy for 2 years if they received front-line immunotherapy.
- Patients should receive maintenance therapy until progression if they received second-line immunotherapy.

ADENOCARCINOMA, LARGE CELL, NSCLC NOS (PS 0–2)	SQUAMOUS CELL CARCINOMA (PS 0–2)
Continuation maintenance <ul style="list-style-type: none"> • Bevacizumab (category 1) • Pemetrexed (category 1) • Bevacizumab/pemetrexed^l • Pembrolizumab/pemetrexed (category 1)^m • Atezolizumab/bevacizumab (category 1)ⁿ • Nivolumab/ipilimumab^o • Atezolizumab^p • Gemcitabine (category 2B) • Cemiplimab-rwlc^q ± pemetrexed^r (category 1) • Durvalumab^s ± pemetrexed^t Switch maintenance <ul style="list-style-type: none"> • Pemetrexed 	Continuation maintenance <ul style="list-style-type: none"> • Pembrolizumab^u • Nivolumab/ipilimumab^o • Gemcitabine (category 2B) • Cemiplimab-rwlc^q (category 1) • Durvalumab^s
ADENOCARCINOMA, LARGE CELL, NSCLC NOS, SQUAMOUS CELL CARCINOMA (PS 3–4)	
Best supportive care (NCCN Guidelines for Palliative Care)	

[Continued](#)[Subsequent Therapy NSCL-K \(4 of 5\)](#)

^k Monitoring During Subsequent Therapy or Maintenance Therapy: Response assessment with CT of known or high-risk sites of disease with or without contrast every 6–12 weeks. Timing of CT scans within Guidelines parameters is a clinical decision.

^l If bevacizumab was used with a first-line pemetrexed/platinum chemotherapy regimen.

^m If pembrolizumab/carboplatin/pemetrexed or pembrolizumab/cisplatin/pemetrexed given.

ⁿ If atezolizumab/carboplatin/paclitaxel/bevacizumab given.

^o If nivolumab + ipilimumab ± chemotherapy given.

^p If atezolizumab/carboplatin/albumin-bound paclitaxel given.

^q If cemiplimab-rwlc combination therapy given.

^r If cemiplimab-rwlc + pemetrexed + (carboplatin or cisplatin) given.

^s If tremelimumab-actl combination therapy given.

^t If tremelimumab-actl + durvalumab + (carboplatin or cisplatin) + pemetrexed given.

^u If pembrolizumab/carboplatin/(paclitaxel or albumin-bound paclitaxel) given.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

[References](#)

NSCL-K
3 OF 5



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

SYSTEMIC THERAPY FOR ADVANCED OR METASTATIC DISEASE – SUBSEQUENT^{d,k}

ADENOCARCINOMA, LARGE CELL, NSCLC NOS (PS 0–2)	SQUAMOUS CELL CARCINOMA (PS 0–2)
<p>Preferred (no previous IO): Systemic immune checkpoint inhibitors^e</p> <ul style="list-style-type: none"> • Nivolumab (category 1) • Pembrolizumab (category 1)^v • Atezolizumab (category 1) <p>Other Recommended (no previous IO or previous IO):^w</p> <ul style="list-style-type: none"> • Docetaxel • Pemetrexed • Gemcitabine • Ramucirumab/docetaxel • Albumin-bound paclitaxel 	<p>Preferred (no previous IO): Systemic immune checkpoint inhibitors^e</p> <ul style="list-style-type: none"> • Nivolumab (category 1) • Pembrolizumab (category 1)^v • Atezolizumab (category 1) <p>Other Recommended (no previous IO or previous IO):^w</p> <ul style="list-style-type: none"> • Docetaxel • Gemcitabine • Ramucirumab/docetaxel • Albumin-bound paclitaxel
ADENOCARCINOMA, LARGE CELL, NSCLC NOS, SQUAMOUS CELL CARCINOMA (PS 3–4)	
Best supportive care (NCCN Guidelines for Palliative Care)	

SYSTEMIC THERAPY FOR ADVANCED OR METASTATIC DISEASE – PROGRESSION^{d,k}

ADENOCARCINOMA, LARGE CELL, NSCLC NOS ^{e,w}	SQUAMOUS CELL CARCINOMA ^{e,w}
<ul style="list-style-type: none"> • PS 0–2: nivolumab, pembrolizumab, or atezolizumab; docetaxel (category 2B); pemetrexed (category 2B); gemcitabine (category 2B); ramucirumab/docetaxel (category 2B); or albumin-bound paclitaxel (category 2B) • PS 3–4: Best supportive care • Options for further progression are best supportive care or clinical trial. 	<ul style="list-style-type: none"> • PS 0–2: nivolumab, pembrolizumab, or atezolizumab; docetaxel (category 2B); gemcitabine (category 2B); ramucirumab/docetaxel (category 2B); or albumin-bound paclitaxel (category 2B) • PS 3–4: Best supportive care • Options for further progression are best supportive care or clinical trial.

^d Contraindications for treatment with PD-1/PD-L1 inhibitors may include active or previously documented autoimmune disease and/or current use of immunosuppressive agents; some oncogenic drivers (ie, *EGFR* exon 19 deletion or L858R, *ALK* rearrangements) have been shown to be associated with less benefit from PD-1/PD-L1 inhibitors.

^e If progression on PD-1/PD-L1 inhibitor, using a PD-1/PD-L1 inhibitor is not recommended.

^k Monitoring During Subsequent Therapy or Maintenance Therapy: Response assessment with CT of known or high-risk sites of disease with or without contrast every 6–12 weeks. Timing of CT scans within Guidelines parameters is a clinical decision.

^v Pembrolizumab is approved for patients with NSCLC tumors with PD-L1 expression levels $\geq 1\%$, as determined by an FDA-approved test.

^w If not previously given.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

References



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

SYSTEMIC THERAPY FOR ADVANCED OR METASTATIC DISEASE – REFERENCES

- ¹ Langer CJ, et al. Carboplatin and pemetrexed with or without pembrolizumab for advanced, non-squamous non-small-cell lung cancer: a randomised, phase 2 cohort of the open-label KEYNOTE-021 study. *Lancet Oncol* 2016;17:1497-1508.
- ² Gandhi L, Rodriguez-Abreu D, Gadgeel S, et al. Pembrolizumab plus chemotherapy in metastatic non-small-cell lung cancer. *N Engl J Med* 2018;378:2078-2092.
- ³ Socinski M, Jotte R, Cappuzzo F, et al. Atezolizumab for first-line treatment of metastatic nonsquamous NSCLC. *N Engl J Med* 2018;378:2288-2301.
- ⁴ West H, McCleod M, Hussein M, et al. Atezolizumab in combination with carboplatin plus nab-paclitaxel chemotherapy compared with chemotherapy alone as first-line treatment for metastatic non-squamous non-small-cell lung cancer (IMPower130): a multicentre, randomised, open-label, phase 3 trial. *Lancet Oncol* 2019;20:924-937.
- ⁵ Hellmann MD, Paz-Ares L, Bernabe Caro, R, et al. Nivolumab plus ipilimumab in advanced non-small-cell lung cancer. *N Engl J Med* 2019;381:2020-2031.
- ⁶ Paz-Ares L, Ciuleanu TE, Cobo M, et al. First-line nivolumab plus ipilimumab combined with two cycles of chemotherapy in patients with non-small-cell lung cancer (CheckMate 9LA): an international, randomised, open-label, phase 3 trial. *Lancet Oncol* 2021;22:198-211.
- ⁷ Gogishvili M, Melkadze T, Makharadze T, et al. Cemiplimab plus chemotherapy versus chemotherapy alone in non-small cell lung cancer: a randomized, controlled, double-blind phase 3 trial. *Nat Med* 2022;28:2374-2380.
- ⁸ Johnson ML, Cho BC, Luft A, et al. Durvalumab with or without tremelimumab in combination with chemotherapy as first-line therapy for metastatic non-small-cell lung cancer: the phase III POSEIDON study. *J Clin Oncol* 2023;41:1213-1227.
- ⁹ Sandler A, Gray R, Perry MC, et al. Paclitaxel-carboplatin alone or with bevacizumab for non-small cell lung cancer. *N Engl J Med* 2006;355:2542-2550.
- ¹⁰ Patel JD, Socinski MA, Garon EB, et al. Pointbreak: a randomized phase III study of pemetrexed plus carboplatin and bevacizumab followed by maintenance pemetrexed and bevacizumab versus paclitaxel plus carboplatin and bevacizumab followed by maintenance bevacizumab in patients with stage IIIB or IV nonsquamous non-small cell lung cancer. *J Clin Oncol* 2013;31:4349-4357.
- ¹¹ Barlesi F, Scherpereel A, Rittmeyer A, et al. Randomized phase III trial of maintenance bevacizumab with or without pemetrexed after first-line induction with bevacizumab, cisplatin, and pemetrexed in advanced nonsquamous non-small cell lung cancer: AVAPERL. *J Clin Oncol* 2013;31:3004-3011.
- ¹² Socinski MA, Bondarenko I, Karaseva NA, et al. Weekly nab-paclitaxel in combination with carboplatin versus solvent-based paclitaxel plus carboplatin as first-line therapy in patients with advanced non-small cell lung cancer: final results of a phase III trial. *J Clin Oncol* 2012;30:2055-2062.
- ¹³ Fossella F, Periera JR, von Pawel J, et al. Randomized, multinational, phase III study of docetaxel plus platinum combinations versus vinorelbine plus cisplatin for advanced non-small-cell lung cancer: the TAX 326 study group. *J Clin Oncol* 2003;21:3016-3024.
- ¹⁴ Klastersky J, Sculier JP, Lacroix H, et al. A randomized study comparing cisplatin or carboplatin with etoposide in patients with advanced non-small cell lung cancer: European Organization for Research and Treatment of Cancer Protocol 07861. *J Clin Oncol* 1990;8:1556-1562.
- ¹⁵ Frasci G, Comella P, Panza N, et al. Carboplatin-oral etoposide personalized dosing in elderly non-small cell lung cancer patients. Gruppo Oncologico Cooperativo Sud-Italia. *Eur J Cancer* 1998;34:1710-1714.
- ¹⁶ Danson S, Middleton MR, O'Byrne KJ, et al. Phase III trial of gemcitabine and carboplatin versus mitomycin, ifosfamide, and cisplatin or mitomycin, vinblastine, and cisplatin in patients with advanced non-small cell lung carcinoma. *Cancer* 2003;98:542-553.
- ¹⁷ Ohe Y, Ohashi Y, Kubota K, et al. Randomized phase III study of cisplatin plus irinotecan versus carboplatin plus paclitaxel, cisplatin plus gemcitabine, and cisplatin plus vinorelbine for advanced non-small-cell lung cancer: Four-Arm Cooperative Study in Japan. *Ann Oncol* 2007;18:317-323.
- ¹⁸ Scagliotti GV, Kortsik C, Dark GG, et al. Pemetrexed combined with oxaliplatin or carboplatin as first-line treatment in advanced non-small cell lung cancer: a multicenter, randomized, phase II trial. *Clin Cancer Res* 2005;11:690-696.
- ¹⁹ Cardenal F, Lopez-Cabrerizo MP, Anton A, et al. Randomized phase III study of gemcitabine-cisplatin versus etoposide-cisplatin in the treatment of locally advanced or metastatic non-small cell lung cancer. *J Clin Oncol* 1999;17:12-18.
- ²⁰ Scagliotti GV, Parikh P, von Pawel J, et al. Phase III study comparing cisplatin plus gemcitabine with cisplatin plus pemetrexed in chemotherapy-naïve patients with advanced-stage NSCLC. *J Clin Oncol* 2008;26:3543-3551.
- ²¹ Schiller JH, Harrington D, Belani CP, et al. Comparison of four chemotherapy regimens for advanced non-small cell lung cancer. *N Engl J Med* 2002;346:92-98.
- ²² Pujol JL, Breton JL, Gervais R, et al. Gemcitabine-docetaxel versus cisplatin-vinorelbine in advanced or metastatic non-small-cell lung cancer: a phase III study addressing the case for cisplatin. *Ann Oncol* 2005;16:602-610.
- ²³ Tan EH, Szczesna A, Krzakowski M, et al. Randomized study of vinorelbine--gemcitabine versus vinorelbine--carboplatin in patients with advanced non-small cell lung cancer. *Lung Cancer* 2005;49:233-240.
- ²⁴ Green M, Manikhas G, Orlov S, et al. Abraxane®, a novel Cremophor®-free, albumin-bound particle form of paclitaxel for the treatment of advanced non-small-cell lung cancer. *Ann Oncol* 2006;17:1263-1268.
- ²⁵ Rizvi N, Riely G, Azzoli C, et al. Phase I/II trial of weekly intravenous 130-nm albumin-bound paclitaxel as initial chemotherapy in patients with atage IV non-small-cell lung cancer. *J Clin Oncol* 2008;26:639-643.
- ²⁶ Socinski MA, Bondarenko I, Karaseva NA, et al. Weekly nab-paclitaxel in combination with carboplatin versus solvent-based paclitaxel plus carboplatin as first-line therapy in patients with advanced non-small cell lung cancer: final results of a phase III trial. *J Clin Oncol* 2012;30:2055-2062.
- ²⁷ Fossella FV, DeVore R, Kerr RN, et al. Randomized phase III trial of docetaxel versus vinorelbine or ifosfamide in patients with advanced non-small cell lung cancer previously treated with platinum-containing chemotherapy regimens. The TAX 320 Non-Small Cell Lung Cancer Study Group. *J Clin Oncol* 2000;18:2354-2362.
- ²⁸ Fidias PM, Dakhil SR, Lyss AP, et al. Phase III study of immediate compared with delayed docetaxel after front-line therapy with gemcitabine plus carboplatin in advanced non-small cell lung cancer. *J Clin Oncol* 2009;27:591-598.
- ²⁹ Zatloukal P, Kanitz E, Magyar P, et al. Gemcitabine in locally advanced and metastatic non-small cell lung cancer: the Central European phase II study. *Lung Cancer* 1998;22:243-250.
- ³⁰ Sederholm C, Hillerdal G, Lamberg K, et al. Phase III trial of gemcitabine plus carboplatin versus single agent gemcitabine in the treatment of locally advanced or metastatic non-small cell lung cancer: the Swedish Lung Cancer Study group. *J Clin Oncol* 2005;23:8380-8288.
- ³¹ Perol M, Chouaid C, Perol D, et al. Randomized, phase III study of gemcitabine or erlotinib maintenance therapy versus observation, with predefined second-line treatment, after cisplatin-gemcitabine induction chemotherapy in advanced non-small cell lung cancer. *J Clin Oncol* 2012;30:3516-3524.
- ³² Lilienbaum RC, Herndon JE, List MA, et al. Single-agent versus combination chemotherapy in advanced non-small cell lung cancer: the cancer and leukemia group B (study 9730). *J Clin Oncol* 2005;23:190-196.
- ³³ Ceresoli GL, Gregorc V, Cordio S, et al. Phase II study of weekly paclitaxel as second-line therapy in patients with advanced non-small cell lung cancer. *Lung Cancer* 2004;44:231-239.
- ³⁴ Yasuda K, Igishi T, Kawasaki Y, et al. Phase II study of weekly paclitaxel in patients with non-small cell lung cancer who have failed previous treatments. *Oncology* 2004;66:347-352.
- ³⁵ Hanna NH, Sheperd FA, Fossella FV, et al. Randomized phase III study of pemetrexed versus docetaxel in patients with non-small cell lung cancer previously treated with chemotherapy. *J Clin Oncol* 2004;22:1589-1597.
- ³⁶ Paz-Ares L, Luft A, Vicente D, et al. Pembrolizumab plus chemotherapy for squamous non-small-cell lung cancer. *N Engl J Med* 2018;379:2040-2051.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

**Table 1. Definitions for T, N, M**

T	Primary Tumor
TX	Primary tumor cannot be assessed, or tumor proven by the presence of malignant cells in sputum or bronchial washings but not visualized by imaging or bronchoscopy
T0	No evidence of primary tumor
Tis	Carcinoma <i>in situ</i> Squamous cell carcinoma <i>in situ</i> (SCIS) Adenocarcinoma <i>in situ</i> (AIS): adenocarcinoma with pure lepidic pattern, ≤3 cm in greatest dimension
T1	Tumor ≤3 cm in greatest dimension, surrounded by lung or visceral pleura, without bronchoscopic evidence of invasion more proximal than the lobar bronchus (i.e., not in the main bronchus)
T1mi	Minimally invasive adenocarcinoma: adenocarcinoma (≤3 cm in greatest dimension) with a predominantly lepidic pattern and ≤5 mm invasion in greatest dimension
T1a	Tumor ≤1 cm in greatest dimension. A superficial, spreading tumor of any size whose invasive component is limited to the bronchial wall and may extend proximal to the main bronchus also is classified as T1a, but these tumors are uncommon.
T1b	Tumor >1 cm but ≤2 cm in greatest dimension
T1c	Tumor >2 cm but ≤3 cm in greatest dimension
T2	Tumor >3 cm but ≤5 cm or having any of the following features: (1) Involves the main bronchus, regardless of distance to the carina, but without involvement of the carina; (2) Invades visceral pleura (PL1 or PL2); (3) Associated with atelectasis or obstructive pneumonitis that extends to the hilar region, involving part or all of the lung
T2a	Tumor >3 cm but ≤4 cm in greatest dimension
T2b	Tumor >4 cm but ≤5 cm in greatest dimension
T3	Tumor >5 cm but ≤7 cm in greatest dimension or directly invading any of the following: parietal pleura (PL3), chest wall (including superior sulcus tumors), phrenic nerve, parietal pericardium; or separate tumor nodule(s) in the same lobe as the primary
T4	Tumor >7 cm or tumor of any size invading one or more of the following: diaphragm, mediastinum, heart, great vessels, trachea, recurrent laryngeal nerve, esophagus, vertebral body, carina; separate tumor nodule(s) in an ipsilateral lobe different from that of the primary

Used with permission of the American College of Surgeons, Chicago, Illinois. The original source for this information is the AJCC Cancer Staging Manual, Eighth Edition (2017) published by Springer International Publishing.



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

Table 1. Definitions for T, N, M (continued)

N	Regional Lymph Nodes
NX	Regional lymph nodes cannot be assessed
N0	No regional lymph node metastasis
N1	Metastasis in ipsilateral peribronchial and/or ipsilateral hilar lymph nodes and intrapulmonary nodes, including involvement by direct extension
N2	Metastasis in ipsilateral mediastinal and/or subcarinal lymph node(s)
N3	Metastasis in contralateral mediastinal, contralateral hilar, ipsilateral or contralateral scalene, or supraclavicular lymph node(s)
M	Distant Metastasis
M0	No distant metastasis
M1	Distant metastasis
M1a	Separate tumor nodule(s) in a contralateral lobe; tumor with pleural or pericardial nodules or malignant pleural or pericardial effusion ^a
M1b	Single extrathoracic metastasis in a single organ (including involvement of a single nonregional node)
M1c	Multiple extrathoracic metastases in a single organ or in multiple organs

Table 2. AJCC Prognostic Groups

	T	N	M		T	N	M
Occult Carcinoma	TX	N0	M0	Stage IIIB	T1a	N3	M0
Stage 0	Tis	N0	M0		T1b	N3	M0
Stage IA1	T1mi	N0	M0		T1c	N3	M0
	T1a	N0	M0		T2a	N3	M0
Stage IA2	T1b	N0	M0		T2b	N3	M0
Stage IA3	T1c	N0	M0		T3	N2	M0
Stage IB	T2a	N0	M0	Stage IIIC	T4	N2	M0
Stage IIA	T2b	N0	M0		T3	N3	M0
Stage IIB	T1a	N1	M0		T4	N3	M0
	T1b	N1	M0	Stage IVA	Any T	Any N	M1a
	T1c	N1	M0		Any T	Any N	M1b
	T2a	N1	M0	Stage IVB	Any T	Any N	M1c
	T2b	N1	M0				
	T3	N0	M0				
Stage IIIA	T1a	N2	M0				
	T1b	N2	M0				
	T1c	N2	M0				
	T2a	N2	M0				
	T2b	N2	M0				
	T3	N1	M0				
	T4	N0	M0				
	T4	N1	M0				

^a Most pleural (pericardial) effusions with lung cancer are a result of the tumor. In a few patients, however, multiple microscopic examinations of pleural (pericardial) fluid are negative for tumor, and the fluid is nonbloody and not an exudate. If these elements and clinical judgment dictate that the effusion is not related to the tumor, the effusion should be excluded as a staging descriptor.

Used with permission of the American College of Surgeons, Chicago, Illinois. The original source for this information is the AJCC Cancer Staging Manual, Eighth Edition (2017) published by Springer International Publishing.



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

Table 3. Comparison of the Descriptors in the Eighth Edition of the TNM Classification of Lung Cancer Compared with the Seventh Edition**

Descriptor	7th Edition T/N/M	8th Edition T/N/M
T component		
0 cm (pure lepidic adenocarcinoma ≤3 cm in total size)	T1a if ≤2 cm; T1b if >2-3 cm	Tis (AIS)
≤0.5 cm invasive size (lepidic predominant adenocarcinoma ≤3 cm total size)	T1a if ≤2 cm; T1b if >2-3 cm	T1mi
≤1 cm	T1a	T1a
>1-2 cm	T1a	T1b
>2-3 cm	T1b	T1c
>3-4 cm	T2a	T2a
>4-5 cm	T2a	T2b
>5-7 cm	T2b	T3
>7 cm	T3	T4
Bronchus <2 cm from carina	T3	T2
Total atelectasis/pneumonitis	T3	T2
Invasion of diaphragm	T3	T4
Invasion of mediastinal pleura	T3	—
N component		
No assessment, no involvement, or involvement of regional lymph nodes	NX, N0, N1, N2, N3	No change
M component		
Metastasis within the thoracic cavity	M1a	M1a
Single extrathoracic metastasis	M1b	M1b
Multiple extrathoracic metastasis	M1b	M1c

*Rami-Porta R, Asamura H, Travis WD, Rusch VW. Lung cancer - major changes in the American Joint Committee on Cancer eighth edition cancer staging manual. CA Cancer J Clin 2017;67:138-155.

**The staging of tumor size in the AJCC Cancer Staging Manual, 7th Edition is based on the total tumor size (invasive and lepidic/noninvasive); whereas, in the AJCC Cancer Staging Manual, 8th Edition, staging is based on invasive size only for non-mucinous adenocarcinoma. However, in mucinous adenocarcinoma, the total tumor size is used.



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

ABBREVIATIONS

AIS	adenocarcinoma in situ	GGO	ground-glass opacity	PCR	polymerase chain reaction
AUC	area under the curve	GTV	gross tumor volume	PD-1	programmed cell death protein 1
BAC	bronchioloalveolar carcinoma	H&P	history and physical	PD-L1	programmed death ligand 1
BED	biologically effective dose	H1 blocker	Histamine-1 blocker	PFT	pulmonary function test
CBC	complete blood count	H2 blocker	Histamine-2 blocker	PORT	postoperative radiotherapy
CBCT	cone-beam CT	IFI	involved field radiation	PS	performance status
CHIP	clonal hematopoiesis of indeterminate potential	IGRT	image-guided radiation therapy	PTV	planning target volume
CLIA	Clinical Laboratory Improvement Amendments	IGTA	image-guided thermal ablation	ROSE	rapid on-site evaluation
CNS	central nervous system	IHC	immunohistochemistry	QUANTEC	Quantitative Analysis of Normal Tissue Effects in the Clinic
COPD	chronic obstructive pulmonary disease	IMPT	intensity-modulated proton therapy	SABR	stereotactic ablative radiotherapy
CPT	Current Procedural Terminology	IMRT	intensity-modulated radiation therapy	SBRT	stereotactic body radiation therapy
CRT	conformal radiation therapy	IO	immuno-oncology	SCLC	small cell lung cancer
ctDNA	circulating tumor DNA	IPF	idiopathic pulmonary fibrosis	SRS	stereotactic radiosurgery
CTV	clinical target volume	ITV	internal target volume	SRT	stereotactic radiation therapy
DLCO	diffusing capacity of the lung for carbon monoxide	LDCT	low-dose computed tomography	SUV	standardized uptake value
		LVEF	left ventricular ejection fraction	SVC	superior vena cava
EBUS	endobronchial ultrasound	MIA	minimally invasive adenocarcinoma	TBNA	transbronchial needle aspiration
ENI	elective nodal irradiation	MLD	mean lung dose	TKI	tyrosine kinase inhibitor
EUS	endoscopic ultrasound			TNM	tumor node metastasis
FDG	fluorodeoxyglucose	NGS	next-generation sequencing	TTNA	transthoracic needle aspiration
FEV1	forced expiratory volume in the first second	NOS	not otherwise specified	UIP	usual interstitial pneumonia
FFPE	formalin-fixed paraffin-embedded	NSCC	non-small cell carcinoma	VATS	video-assisted thoracic surgery
FISH	fluorescence in situ hybridization	NSCLC	non-small cell lung cancer	VMAT	volumetric modulated arc therapy
FNA	fine-needle aspiration	NUT	nuclear protein in testis	VUS	variant of uncertain significance
		OAR	organ(s) at risk	WBRT	whole brain radiation therapy



NCCN Categories of Evidence and Consensus	
Category 1	Based upon high-level evidence, there is uniform NCCN consensus that the intervention is appropriate.
Category 2A	Based upon lower-level evidence, there is uniform NCCN consensus that the intervention is appropriate.
Category 2B	Based upon lower-level evidence, there is NCCN consensus that the intervention is appropriate.
Category 3	Based upon any level of evidence, there is major NCCN disagreement that the intervention is appropriate.

All recommendations are category 2A unless otherwise indicated.

NCCN Categories of Preference	
Preferred intervention	Interventions that are based on superior efficacy, safety, and evidence; and, when appropriate, affordability.
Other recommended intervention	Other interventions that may be somewhat less efficacious, more toxic, or based on less mature data; or significantly less affordable for similar outcomes.
Useful in certain circumstances	Other interventions that may be used for selected patient populations (defined with recommendation).

All recommendations are considered appropriate.



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

Discussion

This discussion corresponds to the NCCN Guidelines for Non-Small Cell Lung Cancer. Last updated: April 13, 2023.

Table of Contents

Overview	MS-3	EGFR Mutations	MS-18
Risk Factors	MS-4	ERBB2 (HER2) Mutations	MS-20
Smoking Cessation	MS-4	KRAS Mutations	MS-21
Lung Cancer Screening	MS-5	MET Genomic Alterations	MS-21
Classification and Prognostic Factors	MS-6	NTRK1/2/3 Gene Fusions	MS-22
Guidelines Update Methodology	MS-6	RET Rearrangements	MS-22
Literature Search Criteria	MS-6	ROS1 Rearrangements	MS-23
Sensitive/Inclusive Language Usage	MS-6	Testing for Immune Biomarkers	MS-24
Diagnostic Evaluation	MS-7	PD-L1 Expression Levels	MS-24
Incidental Lung Nodules	MS-7	TMB	MS-24
Larger Tumors	MS-8	Treatment Approaches	MS-25
Pathologic Evaluation of Lung Cancer	MS-9	Surgery	MS-25
Adenocarcinoma	MS-10	Lymph Node Dissection	MS-26
Immunohistochemistry for Diagnosis of NSCLC	MS-11	Thoracoscopic Lobectomy	MS-26
Staging	MS-12	Stage IIIA N2 Disease	MS-27
Predictive and Prognostic Biomarkers	MS-12	Radiation Therapy	MS-28
Testing for Molecular Biomarkers	MS-15	General Principles	MS-28
ALK Gene Rearrangements	MS-16	Radiation Simulation, Planning, and Delivery	MS-28
BRAF V600E Mutations	MS-17	Target Volumes, Prescription Doses, and Normal Tissue Dose Constraints ...	MS-29
		General Treatment Information	MS-29
		Stereotactic Ablative Radiotherapy	MS-31
		Whole Brain RT and Stereotactic Radiosurgery	MS-33
		Combined Modality Therapy	MS-34



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

Surgery Followed by Adjuvant Therapy: Trial Data and NCCN Recommendations	34	Continuation of Targeted Therapy After Progression on Initial Therapy	MS-107
Preoperative Chemotherapy With or Without Immunotherapy Followed by Surgery	MS-38	Second-Line and Beyond (Subsequent) Systemic Therapy	MS-108
Chemoradiation: Trial Data and NCCN Recommendations	MS-39	Summary	MS-112
Chemotherapy: Trial Data and NCCN Recommendations	MS-42	References	MS-114
Targeted Therapies	MS-45		
Immune Checkpoint Inhibitors	MS-74		
Maintenance Therapy	MS-88		
Clinical Evaluation	MS-90		
Additional Pretreatment Evaluation	MS-91		
Initial Therapy	MS-92		
Stage I, Stage II, and Stage IIIA Disease	MS-93		
Multiple Lung Cancers	MS-95		
Stage IIIB and IIIC NSCLC	MS-96		
Limited Metastatic Disease	MS-96		
Neoadjuvant or Adjuvant Therapy	MS-97		
Chemotherapy, Chemoradiation, Immunotherapy, and Targeted Therapy	MS-97		
Radiation Therapy	MS-99		
Surveillance	MS-100		
Treatment of Recurrences and Distant Metastases	MS-101		
Clinical Trial Data	MS-105		
Number of Cycles of First-Line Systemic Therapy	MS-106		
Maintenance Therapy	MS-106		



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

Overview

Lung cancer is the leading cause of cancer death in the United States.^{1,2} In 2023, an estimated 238,340 new cases (117,550 in males and 120,790 in females) of lung and bronchial cancer will be diagnosed, and 127,070 deaths (67,160 in males and 59,910 in females) are estimated to occur because of the disease.¹ During the COVID pandemic, the diagnosis and treatment of lung cancer have been hampered; however, this has not been reflected in the 2023 estimates for incidence and mortality because of the typical delays in collecting and reporting the data.^{1,3} Only 22.9% of all patients with lung cancer are alive 5 years or more after diagnosis; this includes patients with non-small cell lung cancer (NSCLC) and those with small cell lung cancer (SCLC).⁴ From 2012 to 2018, the overall 5-year relative survival rate for NSCLC with adenocarcinoma histology was 29.8% in the United States.^{4,5} Since 2006 to 2007, the incidence of lung cancer decreased annually by 2.6% in men and 1.1% in women.¹ Common symptoms of lung cancer include cough, hemoptysis, dyspnea, weight loss, and chest pain; patients with symptoms are more likely to have chronic obstructive pulmonary disease (COPD).⁶

Much progress has been made recently for lung cancer, such as screening; minimally invasive techniques for diagnosis and treatment; advances in radiation therapy (RT), including stereotactic ablative radiotherapy (SABR); new targeted therapies; and new immunotherapies.⁷⁻¹² These new treatments are reflected in the improved survival rates for patients with NSCLC. From 2015 to 2016, 2-year relative survival for NSCLC was 42% compared with 34% from 2009 to 2010.¹³ From 1990 to 2020, the death rate from lung cancer dropped by 58% in men; from 2002 to 2020, the death rate dropped by 36% in women.¹ Patients with NSCLC who are eligible for targeted therapies or immunotherapies are now surviving longer; 5-year survival rates range from 15% to 50%, depending on the biomarker.^{12,14-30} Thus, death rates for lung cancer have been declining, although there are still more deaths from

lung cancer than from breast, prostate, colorectal, and brain cancers combined.¹

These NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) for NSCLC were first published in 1996.³¹ Subsequently, the NCCN Guidelines® have been updated at least once a year by the NCCN NSCLC Panel; there were 6 updates for the 2022 guidelines. The *Summary of the Guidelines Updates* describes the most recent revisions to the algorithms, which have been incorporated into this updated Discussion text (see the NCCN Guidelines for NSCLC and *Summary* in this Discussion). New references have been added, which support the new recommendations. For the 2023 update (Version 2), the NCCN NSCLC Panel added a recommendation for adjuvant pembrolizumab for eligible patients with completely resected early-stage NSCLC based on clinical trial data and FDA approval (see *Surgery Followed by Adjuvant Therapy: Trial Data and NCCN Recommendations* in this Discussion).³²

All the systemic therapy regimens have been categorized by preference—based on the biomedical literature and experience of the panel members—using the following categories: 1) preferred interventions; 2) other recommended interventions; and 3) interventions that are useful in certain circumstances (see the NCCN Guidelines for NSCLC). These preference categories emphasize the preferred regimens in clinical practice and do not replace the NCCN Categories of Evidence and Consensus, such as category 1 or category 2A. The NCCN Categories of Preference and NCCN Categories of Evidence and Consensus are two separate systems.

The NCCN Guidelines provide specific category designations for all treatment interventions in the guidelines, which are based on evidence from the biomedical literature and consensus among the panel members. Category 1 recommendations indicate uniform NCCN consensus (at least 85% of the NCCN Member Institutions on the panel) that the intervention



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

is appropriate based on high-level evidence, such as randomized phase 3 trials. Category 2A recommendations indicate uniform NCCN consensus that the intervention is appropriate based on lower-level evidence, such as phase 2 trials. It is important to note that all recommendations are category 2A in the NCCN Guidelines unless otherwise indicated. Category 2B recommendations indicate NCCN consensus (50%–<85% of the NCCN Member Institutions on the panel) that the intervention is appropriate based on lower-level evidence. Category 3 recommendations indicate major NCCN disagreement (at least 50% of the NCCN Member Institutions) that the intervention is appropriate based on any level of evidence. For a category 3 recommendation to remain in the guideline, at least 25% of the NCCN Member Institutions on the panel must vote that the intervention is appropriate. By definition, the NCCN Guidelines cannot incorporate all possible clinical variations and are not intended to replace good clinical judgment or individualization of treatments.

Risk Factors

The primary risk factor for lung cancer is smoking tobacco, which accounts for most lung cancer-related deaths.^{2,33-37} Approximately 350 people die every day from lung cancer in the United States; most of these deaths (81%) are caused by cigarette smoking.¹ Cigarette smoke contains many carcinogenic chemicals (eg, nitrosamines, benzo(a)pyrene diol epoxide).^{36,38} The risk for lung cancer increases with the number of packs of cigarettes smoked per day and with the number of years spent smoking (ie, pack-years of smoking history). Individuals who do not smoke also have an increased relative risk (RR = 1.24) of developing lung cancer if they are exposed to tobacco smoke from those who smoke, termed *second-hand smoke*; other studies have reported a modest risk (hazard ratio [HR], 1.05).^{34,38-41}

Other possible risk factors for lung cancer include disease history (eg, COPD), cancer history, family history of lung cancer, and exposure to

other carcinogens (see the NCCN Guidelines for Lung Cancer Screening, available at www.NCCN.org).^{42,43} The International Agency for Research on Cancer lists several agents known to cause lung cancer, including arsenic, asbestos, beryllium, cadmium, chromium, coal smoke, diesel fumes, nickel, silica, soot, and uranium.⁴⁴⁻⁴⁶ Asbestos is a known carcinogen that increases the risk for lung cancer in people exposed to airborne fibers, especially in individuals who smoke cigarettes. It is estimated that about 3% to 4% of lung cancers are caused by asbestos exposure.⁴⁷ Asbestos also causes pleural mesothelioma (see the NCCN Guidelines for Mesothelioma: Pleural, available at www.NCCN.org). Radon gas, a radioactive gas that is produced by the decay of radium 226, also seems to cause lung cancer.

It is not clear whether hormone replacement therapy (HRT) affects the risk for lung cancer in women. More than 20 studies have been published, but the results have been inconsistent. In a large randomized controlled study, no increase in the incidence of lung cancer was found among postmenopausal women treated with estrogen plus progestin HRT; however, the risk of death increased in those with NSCLC.⁴⁸ In women who received estrogen alone, the incidence or risk of death from lung cancer did not increase.⁴⁹

Smoking Cessation

Approximately 85% to 90% of cases of lung cancer are caused by cigarette smoking.³⁵ Of patients aged 20 to 49 years who were recently diagnosed with lung cancer, 81% of men and 72% of the women had smoked.³ Active smoking causes lung cancer; individuals who formerly smoked are at increased risk for lung cancer compared with those who have never smoked. There is a causal relationship between active smoking and lung cancer and also between other cancers (eg, bladder, cervical, colorectal, esophageal, gastric, kidney, laryngeal, oral cavity, ovarian cancer, pancreatic, pharyngeal) and other diseases and



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

conditions.³⁵ Smoking harms nearly every organ in the body; individuals who smoke have increased mortality compared with those who do not smoke.⁵⁰ Those who do not smoke, but live with someone who smokes, have an increased risk for lung cancer due to secondhand smoke.³⁹ Further complicating this problem, cigarettes also contain nicotine, which is a highly addictive substance.

Oncologists should encourage smoking cessation, especially in patients with cancer (see the NCCN Guidelines for Smoking Cessation, available at www.NCCN.org).⁵¹⁻⁵⁴ The 5 A's framework is a useful tool (that is, Ask, Advise, Assess, Assist, Arrange).⁵⁵ It is in the best interest of patients to quit smoking tobacco. Persistent smoking is associated with second primary cancers, treatment complications, and decreased survival.⁵⁶ Some surgeons will not operate on individuals who currently smoke, because active smoking may increase postoperative pulmonary complications.⁵⁷ However, active smoking should not be used to exclude patients with early-stage lung cancer from surgical treatment that will prolong survival. Programs using behavioral counseling combined with medications that promote smoking cessation (approved by the FDA) can be very useful.⁵⁸ The American Cancer Society (ACS) has resources on *How to Quit Using Tobacco*.

Agents that can be used to promote smoking cessation include nicotine replacement (eg, gum, inhaler, lozenge, nasal spray, patch), bupropion sustained release, and varenicline.^{59,60} A study suggests that cytisine is more efficacious than nicotine replacement therapy, although more side effects were reported with cytisine such as nausea, vomiting, and sleep disorders.⁶¹ Studies have shown that varenicline is better than bupropion or nicotine patch for smoking cessation.⁶²⁻⁶⁴ The effectiveness of varenicline for preventing relapse has not been clearly established.⁶⁵ The FDA has issued an alert for varenicline regarding neuropsychiatric symptoms. Varenicline has also been associated with visual disturbances,

movement disorders, unconsciousness, and cardiovascular disorders; therefore, it is banned in truck and bus drivers, pilots, and air traffic controllers.⁶⁶⁻⁶⁹ Other side effects of varenicline include nausea, abnormal dreams, insomnia, and headache.^{64,70,71} Bupropion may also be associated with similar serious neuropsychiatric symptoms. Nicotine replacement has fewer adverse effects than varenicline or bupropion.⁷² In spite of the potential adverse effects, it is probably more beneficial for motivated patients to use agents to promote smoking cessation.⁷²

Lung Cancer Screening

Lung cancer is the leading cause of cancer death worldwide in men, and late diagnosis is a major obstacle to improving lung cancer outcomes.^{2,73,74} The feasibility of lung cancer screening was assessed because localized cancer can be managed with curative intent and because the mortality rate in other solid tumors (eg, cervix, colon) seems to be decreased by screening, early detection, and prompt treatment.

The National Lung Screening Trial (NLST) (ACRIN Protocol A6654), a phase 3 randomized trial, assessed the risks and benefits of low-dose CT scans compared with chest radiographs for detecting lung cancer in more than 53,000 individuals with a long-term history of smoking cigarettes—including those who currently or previously smoked.⁷⁵ The NLST showed that screening individuals with high-risk factors using low-dose CT decreased the mortality rate from lung cancer by 20%.⁷⁶ Individuals who currently or previously smoked cigarettes were categorized as high risk for lung cancer if they had a 30 or more pack-year smoking history (individuals who previously smoked had quit smoking up to 15 years before enrollment), were 55 to 74 years of age, and had no evidence of lung cancer.^{75,77} NELSON, a phase 3 randomized trial, assessed low-dose CT screening in 15,789 individuals at high risk for lung cancer based on age and smoking history; 85% were men.⁷⁸ Individuals were aged 50 to 74 years and currently smoked or had quit smoking within the last 10 years.



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

At 10 years of follow-up, NELSON demonstrated a reduction in lung cancer mortality in men of 24% (cumulative rate ratio for death from lung cancer: 0.76; 95% CI, 0.61–0.94; $P = .01$).

The NCCN, ACS, U.S. Preventive Services Task Force (USPSTF), American College of Chest Physicians, European Society for Medical Oncology (ESMO), and other organizations recommend lung cancer screening using low-dose CT for select high-risk individuals who either currently or previously smoked cigarettes based on clinical trial data (see the NCCN Guidelines for Lung Cancer Screening, available at www.NCCN.org).⁷⁸⁻⁸³ Low-dose CT screening and follow-up are not a substitute for smoking cessation; patients should be offered smoking cessation counseling (see NCCN Guidelines for Smoking Cessation, available at www.NCCN.org).

Classification and Prognostic Factors

WHO divides lung cancer into two major classes based on its biology, therapy, and prognosis: NSCLC (discussed in these guidelines) and SCLC (see the NCCN Guidelines for Small Cell Lung Cancer, available at www.NCCN.org).⁸⁴⁻⁸⁷ NSCLC accounts for more than 80% of all lung cancer cases, and it includes two major types: 1) nonsquamous, including adenocarcinoma, large-cell carcinoma, and other subtypes; and 2) squamous cell (epidermoid) carcinoma.⁸⁸ Adenocarcinoma is the most common subtype of lung cancer seen in the United States and is also the most frequently occurring histology in individuals who have never smoked cigarettes. The lung adenocarcinoma classification was developed by an international panel and adopted by WHO (see the *Pathologic Evaluation of Lung Cancer* in this Discussion).^{84-87,89} All NSCLC should be classified according to subtype using the WHO Guidelines.^{84,85,87} The guidelines contain an extensive pathology section (see *Principles of Pathologic Review* in the NCCN Guidelines for NSCLC and *Pathologic Evaluation of Lung Cancer* in this Discussion). Certain prognostic factors are predictive

of survival in patients with NSCLC. Good prognostic factors include early-stage disease at diagnosis, good performance status (PS) (ECOG 0, 1), no significant weight loss (<5%), and female gender.⁹⁰

Guidelines Update Methodology

The complete details of the Development and Update of the NCCN Guidelines are available at www.NCCN.org.

Literature Search Criteria

Prior to the update of the NCCN Guidelines for Non-Small Cell Lung Cancer, an electronic search of the PubMed database was performed to obtain key literature in NSCLC published since the previous Guidelines update, using the search term: non-small cell lung cancer. The PubMed database was chosen because it is the most widely used resource for medical literature and indexes peer-reviewed biomedical literature. The search results were narrowed by selecting studies in humans published in English. Results were confined to the following article types: Clinical Trial, Phase 2; Clinical Trial, Phase 3; Clinical Trial, Phase 4; Guideline; Meta-Analysis; Randomized Controlled Trial; Systematic Reviews; and Validation Studies. Data from key PubMed articles as well as articles from additional sources deemed as relevant to these guidelines as discussed by the panel during the Guidelines update have been included in this version of the Discussion section. Recommendations for which high-level evidence is lacking are based on the panel's review of lower-level evidence and expert opinion.

Sensitive/Inclusive Language Usage

NCCN Guidelines strive to use language that advances the goals of equity, inclusion, and representation. NCCN Guidelines endeavor to use language that is person-first; not stigmatizing; anti-racist, anti-classist, anti-misogynist, anti-ageist, anti-ableist, and anti-fat-biased; and inclusive of individuals of all sexual orientations and gender identities. NCCN



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

Guidelines incorporate non-gendered language, instead focusing on organ-specific recommendations. This language is both more accurate and more inclusive and can help fully address the needs of individuals of all sexual orientations and gender identities. NCCN Guidelines will continue to use the terms *men*, *women*, *female*, and *male* when citing statistics, recommendations, or data from organizations or sources that do not use inclusive terms. Most studies do not report how sex and gender data are collected and use these terms interchangeably or inconsistently. If sources do not differentiate gender from sex assigned at birth or organs present, the information is presumed to predominantly represent cisgender individuals. NCCN encourages researchers to collect more specific data in future studies and organizations to use more inclusive and accurate language in their future analyses.

Diagnostic Evaluation

Incidental Lung Nodules

Lung cancer screening is recommended for early diagnosis in asymptomatic patients at high risk for cancer.^{75,78} Risk assessment is used to determine which individuals are at high risk for lung cancer and thus are candidates for screening with low-dose CT.⁹¹ Clinicians are referred to the NCCN Guidelines for Lung Cancer Screening for risk assessment criteria to determine which patients are eligible for screening and for how to evaluate and follow up on low-dose CT screening findings.⁹² The NCCN Guidelines for Lung Cancer Screening have been revised to harmonize with the LungRADs system developed by the American College of Radiology with the goal of decreasing the false-positive low-dose CT screening results reported in the NLST.⁹³

The diagnostic algorithm for incidental pulmonary nodules in the NCCN Guidelines for NSCLC incorporates information from the NCCN Guidelines for Lung Cancer Screening. The diagnostic algorithms for incidental solid and subsolid lung nodules, which are detected on chest CT, use cutoff

thresholds of 6 mm for a positive scan result based on the Fleischner criteria (see the NCCN Guidelines for NSCLC).⁹⁴⁻⁹⁸ Note that the Fleischner Society Guidelines do not specify whether a CT with contrast is necessary for follow-up or whether a low-dose CT is sufficient. Low-dose CT is a preferred recommendation in the NCCN Guidelines unless contrast enhancement is needed for better diagnostic resolution.

Solid and subsolid nodules are the two main types of pulmonary nodules that may be seen on chest CT scans. The Fleischner Society has recommendations for patients with solid and subsolid nodules.^{95,96} Subsolid nodules include: 1) nonsolid nodules also known as ground-glass opacities (GGOs) or ground-glass nodules (GGNs); and 2) part-solid nodules, which contain both ground-glass and solid components.^{96,99-101} Nonsolid nodules are mainly adenocarcinoma in situ (AIS) or minimally invasive adenocarcinoma (MIA), formerly known as bronchioloalveolar carcinoma (BAC) (see *Adenocarcinoma* in this Discussion); patients have 5-year disease-free survival of 100% if these nonsolid nodules are completely resected.^{89,96,99,100,102-104} Data suggest that many nonsolid nodules discovered incidentally on CT imaging will resolve and many of those that persist may not progress to clinically significant cancer.^{102,105,106} Solid and part-solid nodules are more likely to be invasive, faster-growing cancers, factors that are reflected in the increased suspicion and follow-up of these nodules (see the NCCN Guidelines for Lung Cancer Screening, available at www.NCCN.org).^{92,95,96}

All findings and factors for a patient need to be carefully evaluated in a multidisciplinary diagnostic team before establishing a diagnosis of lung cancer and before starting treatment. The NCCN Guidelines recommend biopsy or surgical excision for highly suspicious nodules seen on low-dose CT scans or further surveillance for nodules with a low suspicion of cancer, depending on the type of nodule and a multidisciplinary evaluation of other patient factors (see the NCCN Guidelines for Lung Cancer



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

Screening, available at www.NCCN.org). For patients having repeat scans, the most important radiologic factor is change or stability of a nodule when compared with a previous imaging study. False-positive results (eg, benign intrapulmonary lymph nodes, noncalcified granulomas) frequently occurred with low-dose CT when using the original cutoffs for nodule size deemed suspicious for malignancy from the NLST.⁷⁶ The revised cutoff values for suspicious nodules recommended by the American College of Radiology and incorporated into the LungRADs system have been reported to decrease the false-positive rate from low-dose CT.¹⁰⁷⁻¹⁰⁹

Larger Tumors

The NCCN Guidelines recommend that the diagnostic strategy should be individualized for each patient depending on the size and location of the tumor, the presence of mediastinal or distant disease, patient characteristics (eg, comorbidities), and local expertise. The diagnostic strategy needs to be decided in a multidisciplinary setting. Decisions about whether a biopsy (including what type of biopsy) or surgical excision is appropriate depend on several factors as outlined in the NSCLC algorithm, which were revised by the NCCN Panel for the 2023 update (Version 1) (see *Principles of Diagnostic Evaluation* in the NCCN Guidelines for NSCLC). For example, the NCCN Panel clarified that a preoperative biopsy may be useful for patients with early-stage NSCLC (ie, clinical stage IB or more) who may be candidates for systemic therapy before surgery (also known as neoadjuvant, induction, or preoperative therapy).¹¹⁰ A preoperative biopsy may be appropriate if an intraoperative diagnosis seems to be difficult or very risky (such as a small and central lesion, where it is difficult to do a wedge or intraoperative core needle biopsy). The preferred biopsy technique depends on the disease site and is described in the NSCLC algorithm. For example, radial endobronchial ultrasound (EBUS; also known as endosonography), navigational

bronchoscopy, or transthoracic needle aspiration (TTNA) are recommended for patients with suspected peripheral nodules.¹¹¹

PET/CT imaging is useful before selecting a biopsy site, because it is better to biopsy the site that will confer the highest stage. For patients with suspected nodal disease, pathologic mediastinal lymph node evaluation is recommended with either noninvasive or invasive staging methods, including endoscopic ultrasound–guided fine-needle aspiration (EUS-FNA), EBUS–guided transbronchial needle aspiration (EBUS-TBNA), navigational bronchoscopy, robotic bronchoscopy, or mediastinoscopy (see *Principles of Diagnostic Evaluation* in the NCCN Guidelines for NSCLC). Clinicians use both noninvasive and invasive methods when staging patients.¹¹² The panel decided that a preoperative bronchoscopy may also be preferred for tissue diagnosis and/or mediastinal staging (EBUS). EBUS provides access to nodal stations 2R/2L, 3P, 4R/4L, 7, 10R/10L, 11 to 13, and other hilar nodal stations. EUS provides access to nodal stations 5, 7, 8, and 9.

If pathology results from biopsy or surgical excision indicate a diagnosis of NSCLC, then further evaluation and staging need to be done so that the patient's health care team can determine the most appropriate and effective treatment plan (see *Pathologic Evaluation of Lung Cancer, Staging, and Clinical Evaluation* in this Discussion and the NCCN Guidelines for NSCLC). Diagnosis, staging, and planned resection (eg, lobectomy) are ideally one operative procedure for patients with early-stage disease (see the *Principles of Diagnostic Evaluation* in the NCCN Guidelines for NSCLC). A preoperative or intraoperative tissue diagnosis of lung cancer should be established before doing a lobectomy, bi-lobectomy, or pneumonectomy. If a preoperative or intraoperative tissue diagnosis appears risky or unreliable, multidisciplinary evaluation—that includes interventional radiology, thoracic surgery, and interventional pulmonology—is recommended to determine the safest and most efficient



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

approach, or to provide consensus that a biopsy is too risky or difficult and that anatomic resection can occur without tissue confirmation of lung cancer.

Pathologic Evaluation of Lung Cancer

Pathologic evaluation is done to determine whether patients have primary lung cancer or metastatic cancer, classify the histologic subtype of the lung cancer, determine the extent of invasion, establish whether the surgical margins contain cancer (ie, positive or negative margins), and conduct biomarker diagnostic studies to assess for certain somatic, disease-associated variants/mutations (eg, *EGFR* mutations) or immune biomarkers (eg, PD-L1) (see *Principles of Pathologic Review* in the NCCN Guidelines for NSCLC).¹¹³ Targeted therapy is potentially very effective in patients with NSCLC and specific driver mutations, such as *EGFR* mutations; therefore, tissue needs to be conserved for molecular testing (see *Principles of Molecular and Biomarker Analysis* in the NCCN Guidelines for NSCLC).^{10,114-123} All specimens should be assessed morphologically, including routine staining approaches such as hematoxylin and eosin (H&E) histology (or relevant stains for cytology specimens). Cytology may be sufficient to distinguish adenocarcinomas from squamous cell carcinomas.¹²⁴ Ideally, a diagnosis of NSCLC can be done using H&E findings, clinical findings, imaging studies, and the patient's history, which will conserve tissue for molecular analyses. If necessary, immunohistochemistry (IHC) should be used to distinguish adenocarcinoma, squamous cell carcinoma, metastatic malignancy, and primary pleural mesothelioma (particularly for pleural samplings) (see *Immunohistochemistry for Diagnosis of NSCLC* in this Discussion).¹²⁵ Typically, treatment is not recommended until the patient has been diagnosed with NSCLC.

Preoperative evaluations include examination of the following: bronchial brushings, bronchial washings, sputum, FNA biopsy, core needle biopsy,

endobronchial biopsy, and transbronchial biopsy.^{111,126} Minimally invasive techniques can be used to obtain specimens in patients with advanced unresectable NSCLC;^{127,128} however, diagnosis may be more difficult when using small biopsies and cytology.¹⁰³ When available, rapid on-site evaluation (ROSE) may be used to ensure that transbronchial needle aspirates or EBUS specimens are adequate for diagnosis and biomarker testing.¹²⁹⁻¹³³ The mediastinal lymph nodes are systematically sampled to determine the staging and therapeutic options. Other lung diseases also need to be ruled out, such as tuberculosis, sarcoidosis, and coccidioidomycosis.¹³⁴⁻¹³⁶ Lobectomy or pneumonectomy specimens are evaluated intraoperatively to determine the surgical resection margin status, diagnose incidental nodules discovered at the time of surgery, or evaluate the regional lymph nodes.

Postoperative evaluation provides the pathology characteristics necessary for the classification of tumor type, staging, and prognostic factors. The surgical pathology report should include the WHO histologic classification for carcinomas of the lung.^{85-87,137} The classification for lung adenocarcinoma was determined by an international panel and adopted by the WHO (see *Adenocarcinoma* in this Discussion).^{85-87,89} The classification recommends IHC and molecular studies (see *Principles of Pathologic Review* in the NCCN Guidelines for NSCLC).^{85,138} The use of general categories—such as non-small cell carcinoma (NSCC) or NSCC not otherwise specified (NOS)—should be minimized, because more effective treatment can be selected when the histology is known.

Major subtypes of NSCLC include adenocarcinoma, squamous cell carcinoma, adenosquamous carcinoma, large cell carcinoma, carcinoid tumor, and less common subtypes that are not discussed here (see *Principles of Pathologic Review* in the NCCN Guidelines for NSCLC). All NSCLC should be classified according to subtype using the WHO Guidelines.^{85,87} Ideally, the subtype should be specified. The general



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

terms NSCC or NSCC NOS should be used infrequently and only when a more specific diagnosis cannot be obtained by morphology and/or special staining. The purpose of the pathologic evaluation of NSCLC varies depending on whether the sample is 1) intended for initial diagnosis in a case of suspected NSCLC; 2) a definitive resection sample; or 3) obtained for molecular evaluation in the setting of an established NSCLC diagnosis. Further details are provided in the algorithm.

Adenocarcinomas include AIS, MIA, invasive adenocarcinomas, and invasive adenocarcinoma variants (see *Adenocarcinoma* in this Discussion and the NCCN Guidelines for NSCLC). Squamous cell carcinoma is a malignant epithelial tumor that 1) shows either keratinization and/or intercellular bridges; or 2) is an undifferentiated NSCC that demonstrates positivity for squamous cell carcinoma markers by IHC. Adenosquamous carcinomas are tumors with mixed adenocarcinoma and squamous cell carcinoma components; each component comprises at least 10% of the tumor. Molecular testing is recommended if any adenocarcinoma component is present in a biopsy specimen that is otherwise squamous. Large cell carcinomas are tumors lacking morphologic or IHC evidence of clear lineage, with negative or uninformative stains for squamous cell carcinoma and adenocarcinoma. The diagnosis of large cell carcinoma requires a thoroughly sampled resected tumor and cannot be made on non-resected or cytology specimens. Large cell carcinoma cannot be accurately identified using small samples because of challenges with complete assessment of the lesion.⁸⁹ Immunohistochemical stains that exclude adenocarcinoma (TTF1, Napsin A) and squamous cell carcinoma (p40, p63) need to be used before making a diagnosis of large cell carcinoma; mucin stain is also recommended to assess for occult glandular differentiation. Carcinoid tumors are treated using the neuroendocrine guidelines and not the NSCLC guidelines; however, they are part of the differential

diagnosis of pulmonary lesions (see the NCCN Guidelines for Neuroendocrine and Adrenal Tumors, available at www.NCCN.org). Care should be taken to distinguish typical carcinoids from atypical carcinoids by assessing for necrosis (see the NCCN Guidelines for Small Cell Lung Cancer, available at www.NCCN.org.)

The NCCN NSCLC Panel recommends molecular testing for eligible patients with metastatic NSCLC because FDA-approved agents for lung cancer are available for actionable biomarkers (see *Testing for Molecular Biomarkers* in this Discussion). Molecular testing is recommended for patients with metastatic adenocarcinoma, large cell carcinoma, and NSCLC NOS. Testing may be considered for patients with metastatic squamous cell carcinoma.^{139,140} The NCCN NSCLC Panel also recommends PD-L1 IHC testing (category 1) in all patients with metastatic NSCLC because FDA-approved immunotherapy agents are available for this immune biomarker (see *Testing for Immune Biomarkers* in this Discussion).¹⁴¹

Adenocarcinoma

Most lung carcinomas are adenocarcinomas. The categories for adenocarcinoma include: 1) AIS, which is a preinvasive, typically solitary lesion that is usually non-mucinous; 2) MIA, which is a solitary and discrete non-mucinous lesion with a maximum area of invasion no greater than 0.5 cm; and 3) invasive adenocarcinoma variants (see the NCCN Guidelines for NSCLC). Both AIS and MIA are associated with excellent survival if they are resected. The terms *AIS* and *MIA* should not be used for small samples because of challenges with complete assessment of the lesion.⁸⁹ The categories of BAC or mixed subtype adenocarcinoma are no longer used to classify adenocarcinoma.⁸⁹ The classification for lung adenocarcinoma was developed by an international panel and adopted by WHO.^{85-87,89} The lung classification recommends that use of general categories—NSCC and NSCC NOS—should be minimized, because more



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

effective treatment can be selected when the specific subtype is known (see *Principles of Pathologic Review* in the NCCN Guidelines for NSCLC).¹³⁸

Immunohistochemistry for Diagnosis of NSCLC

To diagnose NSCLC in small tissue samples, judicious use of IHC is strongly recommended to conserve tumor tissue for molecular studies, especially in patients with advanced disease (see *Principles of Pathologic Review* in the NCCN Guidelines for NSCLC).^{125,128,142} Note that the specific IHC analyses used to identify tumor type and lineage (eg, adenocarcinoma vs. squamous cell carcinoma) are distinct from the IHC analyses used to determine whether patients are candidates for anaplastic lymphoma kinase (*ALK*) inhibitor therapy or PD-L1 inhibitor therapy. If necessary, IHC should be used to distinguish adenocarcinoma, squamous cell carcinoma, metastatic malignancy, and primary pleural mesothelioma (particularly for pleural samplings).¹²⁵ IHC is useful for assessing poorly differentiated NSCLC in small biopsy and/or cytology specimens.^{89,143}

Adenocarcinomas are usually positive for thyroid transcription factor-1 (TTF-1), whereas squamous cell carcinomas are often negative for TTF-1 and positive for p40 (or alternatively p63).⁸⁹ Napsin A positivity occurs in more than 80% of lung adenocarcinomas and may also be useful in distinguishing adenocarcinoma from squamous cell carcinoma.^{144,145} Note that p63 can co-stain with TTF-1 or Napsin A in adenocarcinoma. In small biopsy specimens previously classified as NSCC NOS, a panel of TTF-1 (or alternatively Napsin A) and p40 (or alternatively p63) may be sufficient to refine the diagnosis to either adenocarcinoma or squamous cell carcinoma. Thus, two markers may be sufficient to distinguish adenocarcinomas from squamous cell carcinomas.^{89,143}

An appropriate panel of IHC stains should include those relevant for evaluation of metastatic carcinomas to the lung if the primary origin of the

carcinoma is uncertain. It is appropriate to first perform a limited panel of IHC to evaluate for NSCLC and, if negative, then proceed to additional IHC for evaluation of possible metastasis from a distant site. TTF-1 is very useful for distinguishing primary lung adenocarcinoma from metastatic adenocarcinoma, because most (70%–90%) non-mucinous primary adenocarcinomas are TTF-1 positive. TTF-1 is typically negative in squamous cell carcinoma.¹⁴³ However, TTF-1 is also positive in tumors such as thyroid cancer and rarely in a few other organ systems.¹⁴⁶ In addition, thyroglobulin and PAX8 are positive in tumors from patients with thyroid cancer, while they are negative in lung cancer. Immunomarkers that may be useful to assess for metastatic carcinoma to the lung include those for breast carcinoma (ER α , PR, GCDPF-15, mammaglobin, GATA-3), renal cell carcinoma (PAX8), papillary serous carcinoma (PAX8, PAX2, ER), and adenocarcinomas of the gastrointestinal tract (CDX2) or prostate gland (NKX3.1). All typical and atypical carcinoid tumors are positive for chromogranin and synaptophysin.

Although the cytologic diagnosis of NSCLC is generally reliable, it is more difficult to diagnose SCLC (see the NCCN Guidelines for Small Cell Lung Cancer, available at www.NCCN.org).^{111,143,147} Many patients with SCLC have characteristic CT and clinical findings (eg, massive lymphadenopathy, mediastinal invasion). Most SCLCs are immunoreactive for TTF-1; they are typically negative for CK34 β E12 and p63.^{148,149} Many SCLCs also stain positively for markers of neuroendocrine differentiation, including insulinoma-associated protein 1 (INSM1), CD56/NCAM, chromogranin, and synaptophysin. IHC should be used to confirm neuroendocrine differentiation only when appropriate morphologic features—speckled chromatin pattern, nuclear molding, and peripheral palisading—are present. CD56/NCAM, INSM1, chromogranin, and synaptophysin are used to identify neuroendocrine tumors if morphologic suspicion of neuroendocrine differentiation exists.¹⁵⁰ One



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

positive marker is sufficient if the staining is not ambiguous in more than 10% of the tumor cells (TCs).

Malignant pleural mesothelioma is a rare disease (see the NCCN Guidelines for Mesothelioma: Pleural, available at www.NCCN.org).¹⁵¹⁻¹⁵³

The NCCN NSCLC Panel feels that malignant mesothelioma and lung adenocarcinoma can be distinguished using clinical impression, imaging, and a limited panel of immunomarkers (if needed) to preserve tissue for molecular testing. Commonly used immunostains sensitive and specific for adenocarcinoma include pCEA, Claudin-4, TTF-1, and Napsin A (negative in mesothelioma). Other potentially useful markers include B72.3, Ber-EP4, MOC31, and CD15; however, these markers generally do not have the sensitivity and specificity of the commonly used markers. Immunostains sensitive and specific for pleural mesothelioma include WT-1, calretinin, cytokeratin 5/6, and D2-40 (podoplanin antibody) (negative in adenocarcinoma).¹⁵¹⁻¹⁵⁴ Broad epithelial markers such as keratin(s), as well as other lineage-specific markers, should be used when the differential diagnosis includes non-pulmonary and non-mesothelial lesions. Other markers can be useful in the differential diagnosis between mesothelioma and metastatic carcinoma to the lung (see *Principles of Pathologic Review* in the NCCN Guidelines for NSCLC).

Staging

The AJCC Cancer Staging Manual (8th edition) is effective for all cancer cases recorded on or after January 1, 2018.^{155,156} The lung cancer staging system was revised by the International Association for the Study of Lung Cancer (IASLC)¹⁵⁷⁻¹⁵⁹ and was adopted by the AJCC.^{155,156,160,161} The definitions for TNM and the stage grouping for the eighth edition are summarized in Tables 1 and 2 of the staging tables (see *Staging* in the NCCN Guidelines for NSCLC). The descriptors of the TNM classification scheme are summarized in Table 3 of the staging tables.¹⁶² Early-stage disease is stages I and II with negative nodes (N0), whereas locally

advanced disease is stages II and III with positive nodes (N+);¹⁶³ advanced or metastatic disease is stage IV. Pathologic staging uses both clinical staging information (which is noninvasive and includes medical history and physical [H&P] examination, and imaging) and other invasive staging procedures (eg, thoracotomy, examination of lymph nodes using mediastinoscopy).¹⁶⁴

From 2012 to 2018, the overall 5-year relative survival rate for adenocarcinoma was 29.6% in the United States; the 5-year survival rate for squamous cell carcinoma was 23%.⁴ The corresponding 5-year relative survival rates for adenocarcinoma were 70.1% for localized, 44.7% for regional, 9.6% for distant, and 20% for unstaged.⁴ Five-year survival after lobectomy for pathologic stage I NSCLC ranges from 45% to 65%, depending on whether the patient has stage 1A or 1B disease and on the location of the tumor.¹⁶⁵ Another study in patients with stage I disease (n = 19,702) found that 82% had surgical resection and their 5-year overall survival was 54%; for untreated stage I NSCLC, 5-year overall survival was only 6%.¹⁶⁶ Of patients with stage I disease who refused surgery (although it was recommended), 78% died of lung cancer within 5 years.

Predictive and Prognostic Biomarkers

Several biomarkers have emerged as predictive and prognostic markers for NSCLC. A *predictive* biomarker is indicative of therapeutic efficacy, because there is an interaction between the biomarker and therapy on patient outcome. A *prognostic* biomarker is indicative of patient survival independent of the treatment received, because the biomarker is an indicator of the innate tumor behavior (see *KRAS Mutations* in this Discussion). The NSCLC Panel recommends testing for certain molecular and immune biomarkers in all appropriate patients with NSCLC to assess whether patients are eligible for targeted therapies or immunotherapies based on data showing improvement in overall survival for patients receiving targeted therapies or immunotherapies compared with traditional



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

chemotherapy regimens.²¹⁻²⁸ Biomarker testing is recommended in eligible patients with stage IV disease, including M1a, M1b, and M1c. Testing for certain biomarkers is now also recommended in eligible patients with resectable early-stage NSCLC (see *Combined Modality Therapy* in this Discussion).

Predictive molecular biomarkers include *ALK* rearrangements, *BRAF* p.V600E point mutations, *EGFR* mutations, v-erb-b2 avian erythroblastic leukemia viral oncogene homolog 2 (*ERBB2*) (also known as human epidermal growth factor receptor 2 [*HER2*]) mutations, Kirsten Rat Sarcoma virus (*KRAS*) mutations, mesenchymal-epithelial transition factor exon 14 (*MET*ex14) skipping mutations, neurotrophic tyrosine receptor kinase 1/2/3 (*NTRK1/2/3*) gene fusions, rearranged during transfection (*RET*) rearrangements, and *ROS* proto-oncogene 1 (*ROS1*) gene rearrangements; PD-L1 expression is the predictive immune biomarker (see *Principles of Molecular and Biomarker Analysis* in the NCCN Guidelines for NSCLC). For the 2023 update (Version 1), the NCCN NSCLC Panel added testing recommendations for *ERBB2* (*HER2*) mutations. Emerging predictive molecular biomarkers include high-level *MET* amplifications (see *Emerging Biomarkers to Identify Novel Therapies for Patients with Metastatic NSCLC* in the NCCN Guidelines for NSCLC). Targeted agents are available for patients with NSCLC who have high-level *MET* amplifications.¹⁶⁷⁻¹⁷² However, there are less data to support using these agents and they may not be FDA approved for NSCLC; therefore, they are referred to as emerging biomarkers. In 2020, the NCCN Panel deleted tumor mutational burden (TMB) as an emerging immune biomarker based on clinical trial data and other issues (see *TMB* in this Discussion).^{173,174}

The NCCN NSCLC Panel recommends molecular testing based on clinical trial data, but strongly advises broader molecular profiling, to identify these and other rare driver mutations for which targeted therapies may be

available to ensure that patients receive the most appropriate treatment; patients may be eligible for clinical trials for some of these targeted agents.¹⁷⁵ Several online resources are available that describe NSCLC driver events, such as *My Cancer Genome*. Resources are available to assess whether the *HER2* mutations are oncogenic or likely to be oncogenic (see oncoKB.org).¹⁷⁶

The presence of *EGFR* exon 19 deletions or *EGFR* exon 21 L858R mutations is predictive of treatment benefit from *EGFR* tyrosine kinase inhibitor (*EGFR* TKI) therapy, such as osimertinib (see *EGFR Mutations* in this Discussion).^{177,178} Previously, these mutations were referred to as *sensitizing EGFR* mutations; however, the specific mutations are now described. The presence of *EGFR* exon 19 deletions or *EGFR* exon 21 L858R mutations does not appear to be prognostic of survival for patients with NSCLC, independent of therapy.¹⁷⁹ Molecular testing is also recommended in eligible patients with metastatic NSCLC for less common *EGFR* mutations—such as *EGFR* S768I, L861Q, and G719X alterations—based on data showing the efficacy of certain *EGFR* TKIs (see *Oral TKIs that Inhibit EGFR S768I, L861Q, and G719X Alterations* in this Discussion). The panel also recommends testing for *EGFR* exon 20 insertion mutations in eligible patients with metastatic NSCLC based on data showing the efficacy of novel agents as subsequent therapy options (see *Agents that Inhibit EGFR Exon 20 Insertion Mutations* in this Discussion).^{180,181} All of these *EGFR* mutations can be assessed in the same assay, if the assay has been appropriately validated (see *Principles of Molecular and Biomarker Analysis* in the NCCN Guidelines for NSCLC). Because targeted polymerase chain reaction (PCR)-based methods for detecting *EGFR* mutations may under-detect *EGFR* exon 20 insertions, NGS-based strategies are preferred. The phrase *subsequent* therapy is used in these NCCN Guidelines instead of *second-line* or *beyond* therapy, because the line of therapy may vary depending on previous treatment with targeted agents.



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

ALK rearrangements predict for benefit from targeted therapy such as alectinib, brigatinib, or lorlatinib (see *ALK Gene Rearrangements* in this Discussion). Testing for *ALK* rearrangements and *EGFR* mutations is recommended (category 1 for both) for patients with metastatic nonsquamous NSCLC or NSCLC NOS so that patients with these driver mutations can receive effective treatment with targeted agents (see *Targeted Therapies* in this Discussion and the NCCN Guidelines for NSCLC).^{175,182-185} Testing for the other actionable mutations—including *BRAF* p.V600E, *ERBB2* (*HER2*) mutations, *KRAS*, *MET*ex14 skipping, *NTRK1/2/3*, *RET*, and *ROS1*—is also recommended for nonsquamous NSCLC or NSCLC NOS because effective targeted agents are available.

Patients with metastatic NSCLC squamous cell carcinoma can also have actionable biomarkers, such as *EGFR* mutations, although at a lower incidence than those with metastatic NSCLC adenocarcinoma.^{139,140,186,187} Molecular testing for actionable alterations can be considered in patients with metastatic squamous cell carcinoma based on the effectiveness of targeted therapies.^{186,187} The NCCN Panel recommends that molecular testing be considered in all patients with metastatic NSCLC squamous cell carcinoma and not just those with certain characteristics, such as those who have never smoked cigarettes, with small biopsy specimens, and with mixed histology.

For patients with metastatic nonsquamous NSCLC, the NCCN NSCLC Panel currently recommends that a minimum of the following biomarkers should be assessed, including *ALK* rearrangements, *BRAF* mutations, *EGFR* mutations, *ERBB2* (*HER2*) mutations, *KRAS* mutations, *MET*ex14 skipping mutations, *NTRK1/2/3* fusions, *RET* rearrangements, *ROS1* rearrangements, and PD-L1 expression levels; molecular testing can be considered in those with metastatic squamous cell carcinoma. This list of recommended biomarkers has been revised as new oncogenic driver mutations were identified and new agents were approved. The NCCN

NSCLC Panel also recommends molecular testing in eligible patients with metastatic NSCLC for *EGFR* exon 20 insertion mutations and for less common *EGFR* mutations, such as *EGFR* S768I, L861Q, and G719X. Patients with metastatic NSCLC may have other somatic genomic alterations for which targeted therapies may be available even if they are not FDA approved for NSCLC, such as high-level *MET* amplifications; these are referred to as emerging biomarkers (see *Emerging Biomarkers to Identify Novel Therapies for Patients with Metastatic NSCLC* in the NCCN Guidelines for NSCLC).¹⁶⁷⁻¹⁷² In 2020, the NCCN Panel deleted TMB as an emerging immune biomarker based on clinical trial data and other issues (see *TMB* in this Discussion).¹⁷³ The NCCN Guidelines for NSCLC provide recommendations for individual biomarkers that should be tested and recommend testing techniques but do not endorse any specific commercially available biomarker assays or commercial laboratories.¹⁸⁸ Biomarker testing should be done at properly accredited laboratories (minimum of Clinical Laboratory Improvement Amendments [CLIA] accreditation) (see *Principles of Molecular and Biomarker Analysis* in the NCCN Guidelines for NSCLC).

ALK, *BRAF* p.V600E, *EGFR*, *KRAS*, *MET*ex14 skipping mutations, *RET* rearrangements, and *ROS1* rearrangements do not usually overlap; thus, testing for *KRAS* mutations may identify patients who will not benefit from further molecular testing (also known as tiered testing approaches).^{187,189-193} The *KRAS* oncogene is a prognostic biomarker. The presence of *KRAS* mutations is prognostic of poor survival for patients with NSCLC when compared to the absence of *KRAS* mutations, independent of therapy (see *KRAS Mutations* in this Discussion).¹⁹⁴

Information about biomarker testing and plasma circulating tumor DNA (ctDNA) testing (so-called “liquid biopsy”) for actionable mutations is included in the algorithm (see *Principles of Molecular and Biomarker Analysis* in the NCCN Guidelines for NSCLC). Briefly, the panel feels that



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

plasma ctDNA testing should not be used to diagnose NSCLC; tissue should be used to diagnose NSCLC. Standards and guidelines for plasma ctDNA testing for somatic variants/mutations have not been published, there is up to a 30% false-negative rate, and variants can be detected that are not related to the tumor (eg, clonal hematopoiesis of indeterminate potential [CHIP]).^{195,196} For example, an *IDH1* mutation identified by plasma ctDNA testing is likely unrelated to NSCLC, given exceptionally low incidence, and is more likely to represent CHIP. Rare examples of CHIP with *KRAS* mutations have been described, suggesting caution in the interpretation of ctDNA findings.¹⁹⁷ In addition, CHIP can be identified following prior chemotherapy or radiotherapy, further confounding interpretation of variants such as in *TP53*.¹⁹⁸ Given the previous caveats, careful consideration is required to determine whether ctDNA findings reflect a true oncogenic driver or an unrelated finding. For the 2023 update (Version 1), the NCCN Panel added a caveat that many, but not all, ctDNA tests use next-generation sequencing (NGS)-based technology.¹⁹⁹

However, plasma ctDNA testing can be used in specific circumstances if 1) the patient is not medically fit for invasive tissue sampling; or 2) there is insufficient tissue for molecular analysis and follow-up tissue-based analysis will be done if an oncogenic driver is not identified.^{200,201} Data suggest that plasma ctDNA testing is a useful minimally invasive test that can be used to identify *ALK*, *BRAF*, *EGFR*, *HER2*, *MET* exon 14 skipping, *RET*, *ROS1*, and other oncogenic biomarkers that would not otherwise be identified in patients with metastatic NSCLC.^{199,202-204} Molecular testing of plasma ctDNA should be done using clinically validated tests.¹⁹⁹

Testing for Molecular Biomarkers

Molecular testing is used to test for oncogenic genomic driver events for which targeted therapies are available; these somatic genomic alterations (also known as molecular biomarkers) include gene mutations and

fusions.²⁰⁵ Testing for certain biomarkers is also recommended for eligible patients with resectable early-stage and locally advanced NSCLC (see *Surgery Followed by Adjuvant Therapy: Trial Data and NCCN Recommendations* in this Discussion). The panel defines broad molecular profiling for NSCLC as molecular testing that identifies all of the classic actionable biomarkers described in the algorithm [eg, *ALK*, *BRAF*, *EGFR*, *ERBB2* (*HER2*), *KRAS*, *MET*ex14 skipping, *NTRK1/2/3*, *RET*, *ROS1*]—using either a single assay or a combination of a limited number of assays—and optimally also identifies the emerging biomarkers (eg, high-level *MET* amplifications) (see *Summary of the Guidelines Updates* and *Principles of Molecular and Biomarker Analysis* in the NCCN Guidelines for NSCLC). Tiered *KRAS* testing approaches, based on the low prevalence of co-occurring biomarkers, are acceptable (see *KRAS Mutations* in this Discussion).^{175,206} Broad genomic profiling may be used to assess for mechanisms of resistance in patients who have had disease progression on targeted therapy. In addition, broad molecular profiling may be used to distinguish separate primary lung cancers from intrapulmonary metastases (see *Multiple Lung Cancers* in this Discussion). Broad genomic profiling may also help determine eligibility for certain molecularly driven clinical trials.

The various testing methods that may be used to assess for the different biomarkers are described in the algorithm (see *Principles of Molecular and Biomarker Analysis* in the NCCN Guidelines for NSCLC). Broad molecular profiling systems may be used to simultaneously test for multiple biomarkers. NGS (also known as massively parallel sequencing) is a type of broad molecular profiling system that can detect panels of mutations and gene fusions if the NGS platforms have been designed and validated to detect these somatic genomic alterations.²⁰⁷⁻²¹⁵ It is important to recognize that NGS requires quality control as much as any other diagnostic technique; because it is design dependent, the panel of genes and abnormalities detected with NGS will vary depending on the design of



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

the NGS platform. For example, some NGS platforms can detect both mutations and gene fusions, as well as copy number variation, but they are not uniformly present in all NGS assays being conducted either commercially or in institutional laboratories.

Other mutation screening assays are available for detecting multiple biomarkers simultaneously, which can detect more than 50 point mutations; NGS platforms can detect even more biomarkers. However, multiplex PCR systems do not typically detect gene fusions. *ROS1* and *ALK* gene rearrangements can be detected using fluorescence in situ hybridization (FISH), NGS, and other methods (see *Principles of Molecular and Biomarker Analysis* in the NCCN Guidelines for NSCLC).

To minimize tissue use and potential wastage, the NCCN NSCLC Panel recommends that broad molecular profiling be done as part of biomarker testing using a validated test(s) that assesses a minimum of the following potential genetic variants: *ALK* rearrangements, *BRAF* mutations, *EGFR* mutations, *ERBB2* (*HER2*) mutations, *KRAS* mutations, *MET*ex14 skipping mutations, *NTRK1/2/3* gene fusions, *RET* rearrangements, and *ROS1* rearrangements. Both FDA and laboratory-developed test platforms are available that evaluate these and other analytes. Broad molecular profiling is also recommended to identify emerging biomarkers for which effective therapy may be available, such as high-level *MET* amplifications. Although clinicopathologic features—such as smoking status, ethnicity, and histology—are associated with specific somatic, disease-associated variants/mutations (eg, *EGFR* mutations), these features should not be used to select patients for testing. The NCCN Guidelines for NSCLC provide recommendations for individual biomarkers that should be tested and recommend testing techniques, but do not endorse any specific commercially available biomarker assays or commercial laboratories.

Several systems are available to classify the pathogenicity of variants. One classification system uses 1) variants with strong clinical significance

(Tier I); 2) variants with potential clinical significance (Tier II); 3) variants of unknown clinical significance (Tier III); and 4) variants that are benign or likely benign (Tier IV).¹¹⁴ Another classification system uses pathogenic, likely pathogenic, variants of uncertain significance (VUS), likely not pathogenic (likely benign), and not pathogenic (benign); this schema is most commonly applied to germline alterations, with some adoption in somatic testing interpretation.^{216,217} Laboratories that adopt either approach (or others) typically do not report alterations that are classified as not pathogenic/Tier IV. Certain molecular testing methods—such as NGS or Sanger—can identify VUS alterations, while targeted assays generally do not detect them. The NCCN Guidelines note that any variant that is classified as VUS should not be used to select targeted therapy even if the VUS occurs in a gene in which other variants are clinically actionable (see *Principles of Molecular and Biomarker Analysis* in the NCCN Guidelines for NSCLC).

***ALK* Gene Rearrangements**

About 5% of patients with NSCLC have *ALK* gene rearrangements.¹²³ Patients with *ALK* rearrangements are resistant to *EGFR* TKIs but have similar clinical characteristics to those with *EGFR* mutations, such as adenocarcinoma histology and either a light or never smoking history.²¹⁸ The NCCN NSCLC Panel recommends testing for *ALK* rearrangements in patients with metastatic nonsquamous NSCLC based on data showing the efficacy of alectinib, brigatinib, ceritinib, crizotinib, or lorlatinib for *ALK* rearrangements and on FDA approvals.²¹⁹⁻²²³ If patients appear to have squamous cell NSCLC, then *ALK* testing can be considered because *ALK* rearrangements also occur in squamous cell NSCLC, although at a lower rate than nonsquamous NSCLC.^{139,140} For the 2023 update (Version 1), the NCCN Panel now recommends testing for *ALK* rearrangements, in addition to *EGFR* mutations, in eligible patients with resectable early-stage NSCLC (stages IB–IIIA, stage IIIB [only T3,N2]) to assess whether adjuvant therapy with atezolizumab or pembrolizumab is an



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

option. PD-L1 inhibitors are less beneficial in patients with some oncogenic drivers (ie, *EGFR* exon 19 deletions, *EGFR* exon 21 L858R mutations, or *ALK* rearrangements).¹⁸⁶

The different testing methods for *ALK* rearrangements are described in the algorithm (see *Principles of Molecular and Biomarker Analysis* in the NCCN Guidelines for NSCLC). A molecular diagnostic FISH test has been approved by the FDA for detecting *ALK* rearrangements. Rapid prescreening with IHC to assess for *ALK* rearrangements can be done.^{185,193,224-231} An IHC assay for *ALK* rearrangements has also been approved by the FDA. NGS can also be used to assess whether *ALK* rearrangements are present, if the platform has been appropriately designed and validated to detect *ALK* rearrangements.²³²⁻²³⁴ If an actionable oncogenic genetic variant occurs in a patient, usually only one variant is present.^{187,189-193,235,236} Therefore, tiered approaches may identify patients who will not benefit from further molecular testing (see *KRAS Mutations* in this Discussion).

The NCCN Panel has preference stratified the first-line therapy options for patients with *ALK* rearrangement–positive metastatic NSCLC. Alectinib, brigatinib, or lorlatinib are recommended as preferred first-line monotherapy options for patients with *ALK* rearrangement–positive metastatic NSCLC (see *Oral TKIs that Inhibit ALK Rearrangements* in this Discussion). Ceritinib is an “other recommended” option, whereas crizotinib is “useful in certain circumstances.” Data suggest that patients with *ALK* rearrangement–positive metastatic NSCLC do not respond to single-agent immune checkpoint inhibitors (ICIs).¹⁸⁶

Patients typically have disease progression after first-line therapy with alectinib, brigatinib, ceritinib, crizotinib, or lorlatinib; subsequent therapy recommendations are described in the algorithm and often include continuing the first-line targeted therapies, depending on the type of progression [see *Second-Line and Beyond (Subsequent) Systemic*

Therapy in this Discussion and the NCCN Guidelines for NSCLC]. Patients with *ALK* rearrangements often have brain metastases after progression on the initial targeted therapies. Treatment of limited brain lesions—clinical trials have included up to 3–5 progressing sites—in patients with NSCLC differs from that recommended in the NCCN Guidelines for Central Nervous System Cancers, because patients with NSCLC and brain lesions often have long-term survival; therefore, the potential neurocognitive issues that may occur with whole brain RT are a concern.²³⁷ Clinicians are using whole brain RT less often in patients with NSCLC and limited brain lesions.²³⁸ For multiple lesions, whole brain RT is recommended; stereotactic radiosurgery (SRS) may be preferred for patients who have good PS and low systemic tumor burden (see the NCCN Guidelines for Central Nervous System Cancers, available at www.NCCN.org).²³⁹⁻²⁴²

BRAF V600E Mutations

BRAF (v-Raf murine sarcoma viral oncogene homolog B) is a serine/threonine kinase that is part of the MAP/ERK signaling pathway. The *BRAF* p.V600E mutation occurs in 1% to 2% of patients with lung adenocarcinoma; it is the most common of the *BRAF* point mutations when considered across all tumor types.^{190,243} Rare *BRAF* mutations include p.V600K, p.V600D, and other mutations. Patients with *BRAF* p.V600E mutations typically either currently or previously smoked cigarettes, whereas those with *EGFR* mutations or *ALK* rearrangements typically have never smoked.²⁴⁴ Mutations in *BRAF* typically do not overlap with *EGFR* mutations, *MET*ex14 skipping mutations, *RET* rearrangements, *ALK* rearrangements, or *ROS1* rearrangements.^{190,191} Testing for *BRAF* mutations is recommended in patients with metastatic nonsquamous NSCLC. Testing may be considered in patients with metastatic NSCLC squamous cell carcinoma because *BRAF* mutations also occur in squamous cell NSCLC, although at a lower rate than nonsquamous NSCLC.^{139,140,190,191} Real-time PCR, Sanger sequencing, and NGS are the



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

most commonly used methods to assess for *BRAF* mutations (see *Principles of Molecular and Biomarker Analysis* in the NCCN Guidelines for NSCLC).

The NCCN NSCLC Panel recommends testing for *BRAF* mutations in patients with metastatic nonsquamous NSCLC based on data showing the efficacy of dabrafenib plus trametinib for patients with *BRAF* p.V600E mutations and on FDA approval (see *Oral Agents that Inhibit BRAF Mutations* in this Discussion).¹⁹⁰ The NCCN Panel has preference stratified the first-line therapy options for patients with *BRAF* p.V600E mutation–positive metastatic NSCLC. Dabrafenib plus trametinib is a preferred treatment option for patients with *BRAF* p.V600E mutations. If combination therapy with dabrafenib plus trametinib is not tolerated, single-agent therapy with dabrafenib or vemurafenib are treatment options; therefore, these agents are categorized as “useful in certain circumstances.”^{190,191,245} Chemotherapy regimens used for initial systemic therapy (eg, carboplatin plus paclitaxel) are also “useful in certain circumstances.” In patients with *BRAF* p.V600E mutation–positive metastatic NSCLC, the response rate is about 24% to single-agent ICIs.¹⁸⁶

EGFR Mutations

The NCCN NSCLC Panel recommends testing for *EGFR* mutations, including common and uncommon mutations, in eligible patients with metastatic NSCLC based on clinical trial data as described in the following sections. Molecular testing for *EGFR* mutations is also recommended for eligible patients with resectable stage IB to IIIA and stage IIIB (only T3,N2) NSCLC to determine whether adjuvant therapy with osimertinib is an option (see *Surgery Followed by Adjuvant Therapy: Trial Data and NCCN Recommendations* in this Discussion).

EGFR Exon 19 Deletions and EGFR Exon 21 L858R Mutations

In patients with NSCLC, the two most commonly found *EGFR* gene mutations are deletions in exon 19 (with conserved deletion of the LREA

sequence) in 45% of patients with *EGFR* mutations and a point mutation in exon 21 (L858R in 40%). Both mutations result in activation of the tyrosine kinase domain, and both are associated with sensitivity to the small-molecule *EGFR* TKIs, such as afatinib, dacomitinib, erlotinib, gefitinib, and osimertinib (see *Targeted Therapies* in this Discussion).²⁴⁶ Because these *EGFR* mutations are sensitive to the *EGFR* TKIs, they were previously referred to as sensitizing *EGFR* mutations; however, the specific mutations are now described. These common *EGFR* mutations are found in approximately 10% of white patients with NSCLC and up to 50% of Asian patients.²⁴⁷ Other less common mutations (approximately 10%), which are also sensitive to *EGFR* TKIs, include exon 20 p.S768I, exon 21 p.L861Q, and/or exon 18 p.G719X (see *EGFR S768I, L861Q, and G719X Mutations* in this section and *Principles of Molecular and Biomarker Analysis* in the NCCN Guidelines for NSCLC).^{248,249} Data suggest that patients harboring tumors without these specific *EGFR* mutations should not be treated with *EGFR* TKIs in any line of therapy, although there are exceptions.

Most patients with the common *EGFR* mutations have adenocarcinoma histology and either have never smoked cigarettes or previously lightly smoked. Data suggest that *EGFR* mutations can occur in patients with adenosquamous carcinoma, which is harder to discriminate from squamous cell carcinoma in small specimens.²⁵⁰ Patients with pure squamous cell carcinoma are less likely to have the common *EGFR* mutations; those with adenosquamous carcinoma may have mutations.^{139,140,250} However, smoking status, ethnicity, and histology should not be used in selecting patients for testing. The NCCN Panel recommends that molecular testing be considered in all patients with metastatic NSCLC squamous cell carcinoma because these patients may also have actionable biomarkers, such as *EGFR* mutations, although at a lower incidence than those with metastatic NSCLC adenocarcinoma.^{139,140,186,187}



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

The predictive effects of the *EGFR* exon 19 deletions and *EGFR* exon 21 L858R mutations are well-defined. Patients with these common *EGFR* mutations have a significantly better response to afatinib, dacomitinib, erlotinib, gefitinib, or osimertinib.²⁴⁶ Data show that *EGFR* TKI therapy is as effective as first-line monotherapy in patients with advanced NSCLC and common *EGFR* mutations (see *Targeted Therapies* in this Discussion).²⁵¹⁻²⁵⁶ Progression-free survival (PFS) is longer with use of *EGFR* TKI monotherapy in patients with the common *EGFR* mutations when compared with cytotoxic systemic therapy, although overall survival is not statistically different for afatinib, erlotinib, or gefitinib.^{251,252,257,258} Patients with *EGFR* exon 20 insertion mutations are usually resistant to afatinib, dacomitinib, erlotinib, or gefitinib, although there are rare exceptions (eg, p.A763_Y764insFQEA) (see *Principles of Molecular and Biomarker Analysis* in the NCCN Guidelines for NSCLC).²⁵⁹⁻²⁶⁴ Patients typically have disease progression after first-line *EGFR* TKI monotherapy; subsequent therapy recommendations are described in the algorithm [see *Second-Line and Beyond (Subsequent) Systemic Therapy* in this Discussion and the NCCN Guidelines for NSCLC]. The phrase *subsequent therapy* is used in these NCCN Guidelines instead of *second-line or beyond therapy*, because the line of therapy may vary depending on previous treatment with targeted agents.

Most patients with the common *EGFR* mutations become resistant to afatinib, erlotinib, or gefitinib; PFS is about 9.7 to 13 months.^{252,258,265-267} *EGFR* p.Thr790Met (T790M) is an *EGFR* exon 20 mutation that is associated with acquired resistance to *EGFR* TKI therapy and has been reported in about 60% of patients with disease progression after initial response to afatinib, erlotinib, or gefitinib.^{212,267-273} Studies suggest T790M may rarely occur in patients who have not previously received afatinib, erlotinib, or gefitinib.²⁷⁴ Germline T790M confers a high risk for lung cancer regardless of smoking status.²⁷⁵⁻²⁷⁷ Therefore, genetic counseling is recommended for patients if p.T790M is identified before treatment.

Acquired resistance to *EGFR* TKIs may also be associated with histologic transformation from NSCLC to SCLC and with epithelial to mesenchymal transition.²⁷⁸⁻²⁸² The NCCN NSCLC Panel recommends that a tissue biopsy should be considered at progression to rule out SCLC transformation (approximately 6%) and to evaluate mechanisms of resistance.²⁷⁹ Acquired resistance can also be mediated by other molecular events, such as acquisition of *ALK* rearrangement, *MET* or *ERBB2* amplification, and other biomarkers.²⁸³

The NCCN NSCLC Panel recommends testing for *EGFR* mutations (category 1) and other biomarkers in patients with metastatic nonsquamous NSCLC or NSCLC NOS based on data showing the efficacy of afatinib, dacomitinib, erlotinib, gefitinib, or osimertinib and on FDA approvals (see *Agents that Inhibit EGFR Mutations* in this Discussion).^{21,251-256} Molecular testing can be considered for *EGFR* mutations and other biomarkers in patients with squamous cell carcinoma as previously described.

DNA mutational analysis is used to assess for *EGFR* status; IHC is not recommended for detecting *EGFR* mutations.²⁸⁴⁻²⁸⁷ Real-time PCR, Sanger sequencing (paired with tumor enrichment), and NGS are the most commonly used methods to assess *EGFR* mutation status (see *Principles of Molecular and Biomarker Analysis* in the NCCN Guidelines for NSCLC).^{185,284} Direct sequencing of DNA corresponding to exons 18 to 21 (or just testing for exons 19 and 21) is a reasonable approach; however, more sensitive methods are available.^{247,286,288-290} Mutation screening assays using multiplex PCR can simultaneously detect more than 50 point mutations.²⁹¹ NGS is a preferred method for detecting *EGFR* variants, because targeted PCR approaches may miss some *EGFR* exon 20 insertion mutations.²¹⁴

The NCCN Panel has preference stratified the first-line therapy options for patients with *EGFR* mutation-positive (exon 19 deletion, exon 21



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

L858R) metastatic NSCLC. Osimertinib is a preferred first-line EGFR TKI option for patients with *EGFR*-positive metastatic NSCLC (see *Agents that Inhibit EGFR Mutations* in this Discussion). Erlotinib (± bevacizumab or ramucirumab), afatinib, dacomitinib, or gefitinib are “other recommended” EGFR TKI options for first-line therapy. Osimertinib is recommended (category 1) as second-line and beyond (subsequent) therapy for patients with *EGFR* T790M–positive metastatic NSCLC and disease progression on erlotinib (± bevacizumab or ramucirumab), afatinib, dacomitinib, or gefitinib.^{266,292}

EGFR S768I, L861Q, and G719X Alterations

Less common *EGFR* mutations (approximately 10%) that are also sensitive to first-, second-, and third-generation EGFR TKIs (eg, afatinib, erlotinib, gefitinib, osimertinib; classic EGFR TKIs) include exon 20 p.S768I, exon 21 p.L861Q, and exon 18 p.G719X (see *Principles of Molecular and Biomarker Analysis* in the NCCN Guidelines for NSCLC).^{248,249,257,293} The NCCN NSCLC Panel recommends testing for *EGFR* S768I, L861Q, and G719X mutations in eligible patients with metastatic NSCLC based on data showing the efficacy of afatinib or osimertinib as preferred first-line therapy options for patients with *EGFR* S768I, L861Q, and G719X mutation-positive metastatic NSCLC (see *Oral TKIs that Inhibit EGFR S768I, L861Q, and G719X Alterations* in this Discussion).^{257,293} Other recommended TKI options in this setting include dacomitinib, erlotinib, or gefitinib.^{294,295}

EGFR Exon 20 Insertion Mutations

Exon 20 insertions are the third most common *EGFR* mutations; they occur in approximately 2% of patients with NSCLC and 4% to 12% of patients with *EGFR* mutations.^{181,261,296,297} Although there are many different *EGFR* exon 20 insertion mutations, three are more common (insASV, insSVD, and insNPH).¹⁸¹ Most patients with *EGFR* exon 20 insertion mutations have low response rates (≤9%) to afatinib, erlotinib,

or gefitinib.^{180,181} An exception is the p.A763_Y764insFQEA mutation; afatinib, erlotinib, or gefitinib are effective for patients with this *EGFR* exon 20 insertion.²⁵⁹ When used at high doses (160 mg/day), osimertinib is associated with response rates of about 25% in patients with *EGFR* exon 20 insertion mutations, which is much lower than with *EGFR* exon 19 deletions or *EGFR* exon 21 L858R mutations.²⁹⁸ First-line platinum-based chemotherapy (± immunotherapy) is a recommended option for patients with *EGFR* exon 20 mutations (eg, carboplatin plus paclitaxel).²⁹⁹⁻³⁰¹ Patients with *EGFR* exon 20 insertion mutations who receive first-line platinum-based chemotherapy have shorter median overall survival (about 16 months) compared with patients with *EGFR* exon 19 deletions or *EGFR* exon 21 L858R mutations who receive targeted therapy with afatinib, erlotinib, or gefitinib (about 39 months).^{180,302,303} The response rates (0%–25%) to immunotherapy regimens vary, depending on the specific *EGFR* exon 20 insertion mutation.^{181,304,305}

The NCCN NSCLC Panel recommends testing for *EGFR* exon 20 insertion mutations in eligible patients with metastatic NSCLC based on data showing the efficacy of amivantamab-vmjw or mobocertinib as subsequent therapy options for patients with *EGFR* exon 20 insertion mutation-positive metastatic NSCLC and on FDA approvals (see *Agents that Inhibit EGFR Exon 20 Insertion Mutations* in this Discussion).^{180,181} NGS is preferred for detecting *EGFR* exon 20 variants because PCR-based strategies may miss some variants (see *Principles of Molecular and Biomarker Analysis* in the NCCN Guidelines for NSCLC).

ERBB2 (HER2) Mutations

For the 2023 update (Version 1), the NCCN Panel added content about *ERBB2 (HER2)* mutations (see *Principles of Molecular and Biomarker Analysis* in the NCCN Guidelines for NSCLC). *ERBB2* encodes for HER2, which is a receptor tyrosine kinase that is found on the surface of normal



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

epithelial cells that is often overexpressed or mutated in various human malignancies, such as NSCLC. *ERBB2* (*HER2*) alterations are commonly insertion or duplication events in exon 20 but other activating mutations are also observed. The panel added a caveat that while some *ERBB2* (*HER2*) mutations are known to be activating, not all single- or double-nucleotide changes are activating. *ERBB2* (*HER2*) exon 20 mutations occur in approximately 3% of patients (median age, 62 years) with advanced nonsquamous NSCLC.³⁰⁶ Patients tend to be women who have never smoked cigarettes; they have a higher incidence of brain metastases than those with other actionable mutations.^{306,307} Although clinicopathologic features—such as smoking status and histology—are associated with *ERBB2* (*HER2*) activating mutations, these features should not be used to select patients for testing. For the 2023 update (Version 1), the NCCN Panel added testing recommendations for *ERBB2* (*HER2*) mutations. NGS-based approaches are best able to survey the broad spectrum of genomic *ERBB2* (*HER2*) alterations that may occur, although Sanger sequencing and targeted PCR approaches may also be used. The NCCN NSCLC Panel recommends testing for *ERBB2* (*HER2*) mutations in all patients with metastatic nonsquamous NSCLC or NSCLC NOS based on clinical trial data and FDA approval of fam-trastuzumab deruxtecan-nxki [see *Agents that Inhibit ERBB2 (HER2) Mutations* in this Discussion]. Testing for *ERBB2* (*HER2*) mutations can be considered in patients with metastatic squamous cell carcinoma. Resources are available to assess whether the *ERBB2* (*HER2*) mutations are oncogenic or likely to be oncogenic (see oncoKB.org). Data suggest that patients with *ERBB2* mutations respond to first-line immunotherapy regimens.³⁰⁸

KRAS Mutations

KRAS is a G-protein with GTPase activity that is part of the MAP/ERK pathway; point mutations in KRAS most commonly occur at codon 12. Approximately 25% of patients with adenocarcinomas in a North American population have *KRAS* mutations; *KRAS* is the most common mutation in

this population.^{121,177,213,309,310} *KRAS* mutation prevalence is associated with cigarette smoking, unlike many of the other actionable mutations (eg, *EGFR* mutations, *ALK* rearrangements).³¹¹ Patients with *KRAS* mutations appear to have a shorter survival than patients with wild-type *KRAS*; therefore, *KRAS* mutations are prognostic biomarkers.^{194,310,312} *KRAS* mutations do not generally overlap with *EGFR*, *ROS1*, *BRAF*, and *ALK* genetic variants.^{187,190-193,313} Therefore, a tiered approach using *KRAS* testing may identify patients who may not benefit from further molecular biomarker testing.^{175,206} *KRAS* mutations may infrequently overlap with *EGFR* mutations or *RET* rearrangements.^{314,315} In patients with *KRAS* mutation-positive metastatic NSCLC, data suggest the response rate is about 26% for single-agent ICIs.^{186,316} First-line platinum-based chemotherapy (± immunotherapy) is a recommended option for patients with *KRAS* mutations (eg, carboplatin plus paclitaxel).

The NCCN NSCLC Panel recommends testing for *KRAS* mutations in eligible patients with metastatic NSCLC based on data showing the efficacy of adagrasib or sotorasib as subsequent therapy options for patients with *KRAS* p.G12C mutations and on FDA approvals (see *Oral Agents that Inhibit KRAS Mutations* in this Discussion).³¹⁷⁻³¹⁹ Responsiveness to adagrasib or sotorasib has not been assessed for mutations other than *KRAS* G12C. NGS, real-time PCR, and Sanger sequencing (ideally with tumor enrichment) are the most commonly used methods to assess for *KRAS* mutations (see *Principles of Molecular and Biomarker Analysis* in the NCCN Guidelines for NSCLC).

MET Genomic Alterations

C-MET, the hepatocyte growth factor (HGF) receptor, is a tyrosine kinase receptor that is involved in cell survival and proliferation; oncogenic driver genomic alterations in MET include *MET*ex14 skipping mutations, MET gene copy number (GCN) gain or amplification, and MET protein overexpression.¹⁸⁹ *MET* genomic alterations do not typically overlap with



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

EGFR, *ROS1*, *BRAF*, and *ALK* genetic variants.³²⁰ However, *MET*ex14 skipping mutations and *MET* amplification may occur together. *MET*ex14 skipping mutations occur in 3% to 4% of patients with adenocarcinoma NSCLC and 1% to 2% of patients with other NSCLC histologies.^{321,322} *MET*ex14 skipping mutations are more frequent in older women who have never smoked cigarettes.³²³

Several different types of *MET*ex14 skipping mutations may occur, such as mutations, base substitutions, and deletions, which makes it difficult to test for all of the mutations. NGS is the primary method of detecting *MET*ex14 skipping mutations; RNA-based NGS may have improved detection. IHC should not be used to detect *MET*ex14 skipping mutations. In patients with *MET*ex14 skipping mutation–positive metastatic NSCLC, data suggest the response rate is about 16% for single-agent ICIs, even with high PD-L1 levels.^{186,324} Data suggest that patients with *MET* amplification respond to immunotherapy.³²⁵

The NCCN NSCLC Panel recommends testing for *MET*ex14 skipping mutations in eligible patients with metastatic NSCLC based on data showing the efficacy of several agents for patients with *MET*ex14 skipping mutations and on FDA approvals for capmatinib and tepotinib (see *Oral TKIs that Inhibit METex14 Skipping Mutations* in this Discussion).^{326,327} The NCCN Panel has preference stratified the first-line therapy options for patients with *MET*ex14 skipping mutation–positive metastatic NSCLC. The NCCN Panel voted that capmatinib or tepotinib are preferred first-line monotherapy options for patients with *MET*ex14 skipping mutation–positive metastatic NSCLC. The panel also voted that crizotinib or systemic therapy options (such as carboplatin plus paclitaxel) are useful in certain circumstances.

NTRK1/2/3 Gene Fusions

NTRK gene fusions encode tropomyosin receptor kinase (TRK) fusion proteins (eg, TRKA, TRKB, TRKC) that act as oncogenic drivers for solid

tumors including lung, salivary gland, thyroid, and sarcoma.³²⁸⁻³³⁰ A diverse range of solid tumors in children and adults may be caused by *NTRK* gene fusions (eg, *NTRK1*, *NTRK2*, *NTRK3*). It is estimated that *NTRK1/2/3* fusions occur in 0.2% of patients with NSCLC and do not typically overlap with other oncogenic drivers such as *EGFR*, *ALK*, or *ROS1*.³²⁹ Various methods can be used to detect *NTRK1/2/3* gene fusions, including NGS, FISH, IHC, and PCR assays (see *Principles of Molecular and Biomarker Analysis* in the NCCN Guidelines for NSCLC). NGS testing can detect a broad range of *NTRK* gene fusions; however, RNA-based NGS may improve detection. DNA-based NGS may not detect some *NTRK1* and *NTRK3* fusions; RNA-based NGS may be considered to assess for fusions.³³¹ In a clinical trial, *NTRK* gene fusions were detected with NGS (50 patients) and FISH (5 patients).³³⁰ Larotrectinib and entrectinib are oral TKIs that inhibit TRK across a diverse range of solid tumors in patients with *NTRK* gene–fusion positive disease, regardless of age.^{330,332}

The NCCN NSCLC Panel recommends *NTRK1/2/3* gene fusion testing in patients with metastatic NSCLC based on clinical trial data showing the efficacy of larotrectinib or entrectinib for patients with *NTRK* gene fusion–positive disease and on FDA approvals; however, clinical data are limited in NSCLC to support this recommendation (see *Oral TKIs that Inhibit NTRK1/2/3 Gene Fusions* in this Discussion).^{330,333}

RET Rearrangements

RET is a tyrosine kinase receptor that affects cell proliferation and differentiation. Rearrangements may occur in NSCLC between the *RET* gene and other domains, especially kinesin family 5B (*KIF5B*) and coiled coil domain containing-6 (*CCDC6*), which lead to overexpression of the RET protein.^{334,335} *RET* rearrangements occur in about 1% to 2% of patients with NSCLC and are more frequent in patients with adenocarcinoma histology.³³⁴⁻³³⁸ In European patients, *RET*



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

rearrangements occur in individuals who currently smoke cigarettes and those who have never nonsmoked.³³⁶ *RET* rearrangements do not typically overlap with *EGFR*, *ROS1*, *BRAF*, *MET*ex14 skipping, and *ALK* genetic variants.³³⁵ However, a few studies suggest that *RET* rearrangements may infrequently overlap with *EGFR* or *KRAS* mutations.^{314,315} NGS, FISH, and RT-PCR can be used to detect *RET* rearrangements (see *Principles of Molecular and Biomarker Analysis* in the NCCN Guidelines for NSCLC).³³⁵ NGS has high specificity; however, RNA-based NGS is preferable to DNA-based NGS for fusion detection. In patients with *RET*-positive metastatic NSCLC, data suggest the response rate is about 6% to single-agent ICIs.¹⁸⁶

The NCCN NSCLC Panel recommends testing for *RET* rearrangements in eligible patients with metastatic NSCLC based on data showing the efficacy of several agents for patients with *RET* rearrangements and on FDA approvals for selpercatinib and pralsetinib.³³⁹⁻³⁴² Previously, the panel deleted vandetanib for patients with *RET* rearrangements because there are better therapy options.^{343,344} The NCCN Panel has preference stratified the therapy options for patients with *RET* rearrangement-positive metastatic NSCLC. The NCCN Panel voted that selpercatinib or pralsetinib are preferred monotherapy options for patients with *RET* rearrangement-positive metastatic NSCLC; cabozantinib is useful in certain circumstances.

***ROS1* Rearrangements**

Although *ROS1* is a distinct receptor tyrosine kinase, it is very similar to *ALK* and members of the insulin receptor family.^{345,346} It is estimated that *ROS1* gene rearrangements occur in about 1% to 2% of patients with NSCLC.³⁴⁶⁻³⁴⁹ The NCCN NSCLC Panel recommends *ROS1* testing in patients with metastatic nonsquamous NSCLC or NSCLC NOS based on data showing the efficacy of crizotinib, ceritinib, and entrectinib for patients with *ROS1* rearrangements.^{192,346,350,351} *ROS1* testing can be

considered in patients with metastatic squamous cell NSCLC because *ROS1* rearrangements also occur in metastatic squamous cell NSCLC, although at a lower rate than nonsquamous NSCLC.^{139,140} Various methods can be used to detect *ROS1* rearrangements, including NGS, FISH, IHC, and PCR assays, although some methods are more effective (see *Principles of Molecular and Biomarker Analysis* in the NCCN Guidelines for NSCLC).^{224,346,348,351-355} False-negative results may occur with FISH, IHC, PCR and DNA-based NGS.³⁵⁶ RNA-based NGS may be considered to assess for fusions.

The NCCN NSCLC Panel recommends crizotinib, entrectinib, or ceritinib as first-line monotherapy options for patients with *ROS1*-positive metastatic NSCLC based on clinical trial data (see *Oral TKIs that Inhibit ROS1 Rearrangements* in this Discussion). The NCCN Panel has preference stratified the first-line therapy options for patients with *ROS1*-positive metastatic NSCLC. The NCCN NSCLC Panel voted that crizotinib and entrectinib are preferred first-line therapy options for patients with *ROS1*-positive metastatic NSCLC because they are better tolerated, have been assessed in more patients, and are approved by the FDA.^{332,350,351,357} Although entrectinib has better central nervous system (CNS) penetration than crizotinib, it is more toxic. The NCCN NSCLC Panel voted that ceritinib is an “other recommended” first-line therapy option for patients with *ROS1*-positive metastatic NSCLC. If *ROS1* rearrangements are discovered during first-line systemic therapy (eg, carboplatin plus paclitaxel), then the planned therapy may be either completed or interrupted followed by crizotinib (preferred), entrectinib (preferred), or ceritinib.

The NCCN NSCLC Panel recommends lorlatinib as a subsequent therapy option for select patients with *ROS1*-positive metastatic NSCLC and disease progression after treatment with crizotinib, entrectinib, or ceritinib (see *Oral TKIs that Inhibit ROS1 Rearrangements* in this Discussion).³⁵⁸



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

However, the panel clarified that entrectinib is recommended as a subsequent therapy option for patients with symptomatic brain lesions after progression on crizotinib or ceritinib.³⁵⁹ Initial systemic therapy options that are used for adenocarcinoma or squamous cell carcinoma are also an option in this setting (eg, carboplatin plus paclitaxel). In patients with *ROS1*-positive metastatic NSCLC, data suggest the response rate is about 17% for single-agent ICIs.¹⁸⁶ Local therapy may be considered for limited progression.

Testing for Immune Biomarkers

PD-L1 Expression Levels

Human ICI antibodies inhibit the PD-1 receptor or PD-L1, which improves antitumor immunity; PD-1 receptors are expressed on activated cytotoxic T cells.³⁶⁰⁻³⁶² Cemiplimab-rwlc, nivolumab, and pembrolizumab inhibit PD-1 receptors.^{141,363,364} Atezolizumab and durvalumab inhibit PD-L1.^{365,366} The NCCN NSCLC Panel recommends IHC testing for PD-L1 expression (category 1) ideally before first-line treatment (if clinically feasible) in all patients with metastatic NSCLC to assess whether the ICI regimens are an option based on clinical data showing the efficacy of these regimens (see *Immune Checkpoint Inhibitors* in this Discussion).^{141,367}

Although it is not an optimal biomarker, PD-L1 expression is currently the best available biomarker to assess whether patients are candidates for PD-1 or PD-L1 inhibitors (also known as ICIs, immuno-oncology [IO] agents, immunotherapy).^{368,369} PD-L1 expression is continuously variable and dynamic; thus, a cutoff value for a positive result is artificial. Patients with PD-L1 expression levels just below and just above 50% will probably have similar responses.³⁶⁸ Unique anti-PD-L1 IHC assays have been developed for each one of the different ICIs.^{368,370-372} The definition of a positive or negative PD-L1 test result depends on the individual antibody, clone, and platform, which may be unique to each ICI.³⁷² Extensive effort has been undertaken to examine the cross-comparability of different

clones with regard to each other to facilitate adoption of testing. While some clones for PD-L1 are FDA-approved for specific indications, using multiple IHC tests is not necessary if the individual IHC test has been validated against the FDA-approved clone. Testing for PD-L1 is not required for prescribing first-line therapy with certain ICI regimens—such as cemiplimab-rwlc monotherapy or atezolizumab with or without chemotherapy—or for subsequent therapy with single-agent nivolumab or atezolizumab.

The NCCN NSCLC Panel emphasizes that clinicians should obtain molecular testing results for actionable biomarkers before administering first-line ICI therapy, if clinically feasible, including *ALK*, *BRAF*, *EGFR*, *ERBB2 (HER2)*, *KRAS*, *MET*ex14 skipping, *NTRK1/2/3*, *RET*, and *ROS1* variants. If it is not feasible to do molecular testing, then patients are treated as though they do not have driver oncogenes. Patients with metastatic NSCLC and PD-L1 expression levels of 1% or more—but who also have a targetable driver oncogene molecular variant—should receive first-line targeted therapy for that oncogene and not first-line ICIs, because targeted therapies yield higher response rates (eg, osimertinib, 80%) than ICIs (lower response rates) in the first-line setting, targeted therapy is better tolerated, and these patients are less likely to respond to single-agent ICIs.^{186,373-376}

TMB

TMB is an approximate measure of the total number of somatic mutations.³⁷⁷ Theoretically, high TMB levels will correlate with high neoantigen levels that will activate an antitumor immune response.³⁷⁸ TMB levels are typically high in patients with NSCLC who currently or previously smoked cigarettes. Low TMB is more commonly detected in individuals who have never smoked cigarettes.^{331,379} Preliminary data for PFS from CHECKMATE 227, a phase 3 randomized trial with a complex design, suggested that TMB might be a useful immune biomarker for deciding



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

whether to use immunotherapy in patients with metastatic NSCLC.³⁸⁰ However, updated data from CHECKMATE 227 showed that overall survival was improved with nivolumab plus ipilimumab regardless of TMB or PD-L1 expression levels.³⁸¹ In addition, combining TMB with PD-L1 expression level also did not correlate with overall survival. Several trials have shown that high TMB levels do not correlate with PD-L1 expression levels in patients with NSCLC.³⁸⁰⁻³⁸³ KEYNOTE 158, a phase 2 trial, assessed TMB levels in patients with solid tumors who received pembrolizumab as second-line therapy; however, none of the patients had NSCLC.³⁸⁴ TMB does not identify patients who will respond to chemotherapy; therefore, TMB has limited value for assessing combination immunotherapy plus chemotherapy regimens.³⁷⁸ TMB is also not an ideal immune biomarker because some patients with low TMB levels respond to immunotherapy and others with high levels do not respond to immunotherapy.³⁷⁸

In addition to the lack of clinical data to support use of TMB as an immune biomarker, there are technical problems with measuring TMB.³⁷⁷ These problems include: 1) lack of agreement on the definition of a cut off for designating high TMB levels; and 2) lack of standardization of TMB measurements across laboratories.³⁷⁷ PD-L1 expression level is a more useful immune biomarker than TMB for deciding how to use immunotherapy, because test results are obtained more quickly, less tissue is needed for testing, and data demonstrate relative reproducibility across platforms and individuals. In 2020, the NCCN Panel removed TMB as an emerging immune biomarker for patients with metastatic NSCLC based on clinical trial data, concerns about variable TMB measurements, and other issues as described here.^{173,377,381} The NCCN Guidelines do not recommend measurement of TMB levels before deciding whether to use nivolumab plus ipilimumab regimens or to use other ICIs, such as pembrolizumab.¹⁷³

Treatment Approaches

Surgery, RT, and systemic therapy are the three modalities most commonly used to treat patients with NSCLC. They can be used either alone or in combination depending on the disease status. In the following sections, the clinical trials are described that have led to the recommended treatments. For tools to aid optimal assessment and care of older adults, see the NCCN Guidelines for Older Adult Oncology (available at www.NCCN.org). Older adults may be at risk for treatment-related adverse events.³⁸⁵

Surgery

In general, surgery provides the best chance for cure in patients with stage I or II disease.³⁸⁶ Thoracic surgical oncology consultation should be part of the evaluation of any patient being considered for curative local therapy. The overall plan of treatment and the necessary imaging studies should be determined before any nonemergency treatment is initiated. It is essential to determine whether patients can tolerate surgery or whether they are medically inoperable; some patients deemed inoperable may be able to tolerate minimally invasive surgery and/or sublobar resection.³⁸⁶⁻³⁹² Although frailty is an increasingly recognized predictor of surgical and other treatment morbidity, a preferred frailty assessment system has not been established.³⁹³⁻³⁹⁵

The *Principles of Surgical Therapy* are described in the NSCLC algorithm and are summarized here. Determination of resectability, surgical staging, and pulmonary resection should be performed by thoracic surgeons who should participate in multidisciplinary clinics and/or tumor boards for patients with lung cancer. Surgery may be appropriate for select patients with uncommon types of lung cancer (eg, superior sulcus, chest wall involvement) (see the NCCN Guidelines for NSCLC).³⁹⁶ Patients with clinical stage IB or greater disease, or high-risk factors, should be referred to a medical oncologist for evaluation. For patients with stage IIIA NSCLC



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

that is deemed resectable, consider referral to a radiation oncologist. Treatment delays, because of poor coordination among specialists, should be avoided.

The surgical procedure used depends on the extent of disease and on the cardiopulmonary reserve of the patient. A preoperative or intraoperative tissue diagnosis of lung cancer should be established before doing a lobectomy, bilobectomy, or pneumonectomy. If a preoperative or intraoperative tissue diagnosis appears risky or unreliable, multidisciplinary evaluation is recommended to determine the safest and most efficient approach, or to provide consensus that a biopsy is too risky or difficult and that anatomic resection can occur without prior tissue confirmation of lung cancer.

Lung-sparing anatomic resection (sleeve lobectomy [also known as sleeve resection]) is preferred over pneumonectomy, if anatomically appropriate and if margin-negative resection can be achieved; lobectomy or pneumonectomy should be done if physiologically feasible.^{386,397,398}

Sublobar resection, either segmentectomy (preferred) or wedge resection, is appropriate in select patients: 1) those who are not eligible for lobectomy; and 2) those with a peripheral nodule 2 cm or less with very low-risk features (see *Principles of Surgical Therapy* in the NCCN Guidelines for NSCLC).^{387,399-404} Segmentectomy (preferred) or wedge resection should achieve parenchymal resection margins that are: 1) 2 cm or more; or 2) the size of the nodule or larger. Resection (including wedge resection) is preferred over ablation.^{386,398} Wide wedge resection may improve outcomes.⁴⁰⁵

Patients with medically inoperable early-stage NSCLC may be candidates for definitive RT, preferably SABR, also known as stereotactic body RT (SBRT).^{406,407} If SABR is considered for patients at high risk for surgical morbidity, a multidisciplinary evaluation is recommended (see *Stereotactic Ablative Radiotherapy* in this Discussion).^{406,408-410}

Lymph Node Dissection

ACOSOG Z0030, a randomized phase 3 trial, compared systematic mediastinal lymph node sampling versus complete lymphadenectomy during pulmonary resection in patients with NSCLC who had either N0 (no demonstrable metastasis to regional lymph nodes) or N1 (metastasis to lymph nodes in the ipsilateral peribronchial and/or hilar region, including direct extension) disease. In patients with early-stage NSCLC who had negative nodes by systematic lymph node dissection, complete mediastinal lymph node dissection did not improve survival.^{411,412} Thus, systematic lymph node sampling is appropriate during pulmonary resection; one or more nodes should be sampled from all mediastinal stations. For right-sided cancers, an adequate mediastinal lymphadenectomy should include stations 2R, 4R, 7, 8, and 9. For left-sided cancers, stations 4L, 5, 6, 7, 8, and 9 should be sampled.⁴¹¹ Patients should have N1 and N2 node resection and mapping (American Thoracic Society map) with a minimum of three N2 stations sampled or a complete lymph node dissection.¹⁵⁵ The lymph node map from the IASLC may be useful.⁴¹³ Formal ipsilateral mediastinal lymph node dissection is indicated for patients undergoing resection for stage IIIA (N2) disease. For patients undergoing sublobar resection, the appropriate N1 and N2 lymph node stations should be sampled unless not technically feasible because sampling would substantially increase the surgical risk.

Thoracoscopic Lobectomy

Video-assisted thoracic surgery (VATS), which is also known as thoracoscopic lobectomy, is a minimally invasive surgical treatment (see *Principles of Surgical Therapy* in the NCCN Guidelines for NSCLC).^{414,415} Published studies suggest that thoracoscopic lobectomy has several advantages over lobectomy by thoracotomy.⁴¹⁶⁻⁴²⁰ Acute and chronic pain associated with thoracoscopic lobectomy is minimal; thus, this procedure requires a shorter length of hospitalization.^{421,422} Thoracoscopic lobectomy is also associated with low postoperative morbidity and mortality, minimal



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

risk of intraoperative bleeding, or minimal locoregional recurrence.⁴²³⁻⁴²⁷ Thoracoscopic lobectomy is associated with less morbidity, fewer complications, and more rapid return to function than lobectomy by thoracotomy.⁴²⁸⁻⁴³¹

In patients with stage I NSCLC who had thoracoscopic lobectomy with lymph node dissection, the 5-year survival rate, long-term survival, and local recurrence rate were comparable to those achieved by routine open lung resection.⁴³²⁻⁴³⁶ Thoracoscopic lobectomy has also been shown to improve discharge independence in older populations and patients at high risk.^{437,438} Data show that thoracoscopic lobectomy improves the ability of patients to complete postoperative chemotherapy regimens.^{439,440} Based on its favorable effects on postoperative recovery and morbidity, thoracoscopic lobectomy (including robotic-assisted approaches) is recommended in the NSCLC algorithm as an acceptable approach for patients who are surgically resectable (and have no anatomic or surgical contraindications) as long as principles of thoracic surgery are not compromised (see *Principles of Surgical Therapy* in the NCCN Guidelines for NSCLC).⁴⁴¹⁻⁴⁴⁴ Robotic VATS seems to be more expensive with longer operating times than conventional VATS.^{445,446}

Stage IIIA N2 Disease

The role of surgery in patients with pathologically documented stage IIIA (N2) disease is described in the NSCLC algorithm (see *Principles of Surgical Therapy* in the NCCN Guidelines for NSCLC) and summarized here. Before treatment, it is essential to carefully evaluate for N2 disease using radiologic and invasive staging (ie, EBUS-guided procedures, mediastinoscopy, thoracoscopic procedures) and to discuss whether surgery is appropriate in a multidisciplinary team, which should include a thoracic surgeon.^{447,448} Randomized controlled trials suggest that surgery does not increase survival in these patients.^{449,450} However, one of these trials (EORTC) only enrolled patients with unresectable disease.⁴⁵⁰ Most

clinicians agree that resection is appropriate for patients with negative preoperative mediastinal nodes and with a single positive node (<3 cm) found at thoracotomy.⁴⁵¹ Neoadjuvant (also known as preoperative or induction) systemic therapy is recommended for select patients with resectable NSCLC (see *Combined Modality Therapy* in this Discussion).

The optimal timing of RT in trimodality therapy (preoperative with chemotherapy or postoperative) is not established and is controversial.⁴⁵²⁻⁴⁵⁴ Adding RT to induction regimens for patients with stage IIIA (N2) disease has been associated with higher pathological response but similar overall survival when compared with using preoperative chemotherapy (generally followed by postoperative RT).⁴⁵⁴ For patients with completely resected stage IIIA (N2) NSCLC who received neoadjuvant or adjuvant chemotherapy, data from the LungART and PORT-C trials show that postoperative RT (also known as PORT) did not improve survival compared with no postoperative RT, although locoregional control was significantly improved.^{452,455} Postoperative RT may be considered for select patients with negative margins and high-risk N2 disease (see *Chemoradiation: Trial Data and NCCN Recommendations* in this Discussion and *Principles of Radiation Therapy* in the algorithm). It is controversial whether pneumonectomy after preoperative chemoradiotherapy is appropriate.^{449,456-462} Clinicians also agree that resection is not appropriate for patients with multiple pathologically proven malignant lymph nodes greater than 3 cm; definitive chemoradiotherapy is recommended for these patients. Patients with resectable stage IIIA (N2) disease should not be excluded from surgery, because some of them may have long-term survival or may be cured.^{457,463}

The NCCN NSCLC Panel believes that surgery may be appropriate for select patients with N2 disease.^{447,457,464} The NCCN Member Institutions were surveyed in 2021 regarding their approach to patients with N2



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

disease (see *Principles of Surgical Therapy* in the NCCN Guidelines for NSCLC). For example, more institutions now use neoadjuvant chemotherapy compared with neoadjuvant chemoradiation in patients with N2 disease. Many (66%) of the NCCN Member Institutions use induction chemotherapy, whereas 33% use induction chemoradiation.^{465,466} Before surgery, most institutions require at least stable disease after induction therapy but do not require radiologic or pathologic response. All NCCN institutions consider surgery for single-station non-bulky N2 disease. However, 50% consider surgery for single-station bulky disease, 39% for multi-station non-bulky disease, and 21% for multi-station bulky disease.

Radiation Therapy

The *Principles of Radiation Therapy* in the NSCLC algorithm include the following: 1) general principles for early-stage, locally advanced, and advanced/metastatic NSCLC; 2) target volumes, prescription doses, and normal tissue dose constraints for early-stage, locally advanced, and advanced/metastatic NSCLC; and 3) RT simulation, planning, and delivery.⁴⁶⁷⁻⁴⁷⁷ These RT principles are summarized in this section. Whole brain RT and SRS for brain metastases are also discussed in this section. The RT abbreviations are defined in the NSCLC algorithm (see Table 1 in *Principles of Radiation Therapy* in the NCCN Guidelines for NSCLC). For the 2023 update (Version 1), the NCCN NSCLC Panel revised some of the RT recommendations in the algorithm (see *Summary of the Guidelines Updates* and *Principles of Radiation Therapy* in the NCCN Guidelines for NSCLC). For example, recommendations about postoperative RT have been revised based on clinical trial data.^{452,455,473-477}

General Principles

Treatment recommendations should be made by a multidisciplinary team. Because RT has a potential role in all stages of NSCLC, as either definitive or palliative therapy, input from radiation oncologists who perform lung cancer RT as a prominent part of their practice should be

part of the multidisciplinary evaluation for all patients with NSCLC. RT recommendations for NSCLC include: 1) definitive therapy for locally advanced NSCLC, generally combined with chemotherapy; 2) definitive therapy for early-stage NSCLC in patients with contraindications for surgery; 3) neoadjuvant or adjuvant therapy (also known as preoperative or postoperative therapy) for selected patients treated with surgery; 4) therapy for limited progression; and/or 5) palliative therapy for patients with incurable NSCLC.^{410,452,455,478-485} The goals of RT are to maximize tumor control and to minimize treatment toxicity. Advanced technologies such as 4D-CT simulation, intensity-modulated RT/volumetric modulated arc therapy (IMRT/VMAT), image-guided RT (IGRT), motion management strategies, and proton therapy have been shown to reduce toxicity and increase survival in nonrandomized trials.^{474,486-492} A secondary analysis of the RTOG 0617 randomized trial reported that 2-year overall survival, PFS, local failure, and distant metastasis-free survival were not significantly different for IMRT when compared with older 3D-conformal RT (3D-CRT) technique despite higher tumor burden in patients treated with IMRT. However, IMRT yielded lower rates of severe pneumonitis compared with 3D-CRT (3.5% vs. 7.9%; $P = .039$).⁴⁹³ CT-planned 3D-CRT is now considered to be the minimum technological standard, with IMRT preferred.

Radiation Simulation, Planning, and Delivery

Simulation should be performed using CT scans obtained in the RT treatment position. Intravenous contrast CT scans, with or without oral contrast, are recommended for better target delineation whenever possible, especially in patients with central tumors or nodal involvement. FDG PET/CT can significantly improve target delineation accuracy, especially when there is atelectasis or contraindications to intravenous CT contrast.^{494,495} Ideally, target delineation should be based on PET/CT that is obtained no more than 4 weeks before treatment because of the potential for rapid progression of NSCLC.^{496,497} In the NSCLC algorithm,



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

recommendations are provided for patients receiving chemoradiation (including those with compromised lung or cardiac function), photon beams, or IMRT (see the *Principles of Radiation Therapy* in the NCCN Guidelines for NSCLC).^{491,498-501} Respiratory motion should be managed. The report from the AAPM Task Group 76 is a useful reference for implementing a broad range of motion management strategies as described in the NSCLC algorithm.⁵⁰²

Target Volumes, Prescription Doses, and Normal Tissue Dose Constraints

Commonly used prescription RT (or SABR) doses and normal tissue dose constraints are summarized in the *Principles of Radiation Therapy* in the NSCLC algorithm (see Tables 2–5 in the NCCN Guidelines for NSCLC).^{468,470,482,503-508} Reports 50, 62, and 83 from the International Commission on Radiation Units and Measurements provide a formalism for defining RT target volumes based on grossly visible disease, potential microscopic extension, and margins for target motion and daily positioning uncertainty.^{509,510} the ACR Practice Parameters and Technical Standards are also a helpful reference.^{488,511,512} It is essential to evaluate the dose-volume histogram (DVH) of critical structures and to limit the doses to the organs at risk (such as spinal cord, lungs, heart, esophagus, and brachial plexus) to minimize normal tissue toxicity (see Table 5 in *Principles of Radiation Therapy*).⁵¹³ For patients receiving postoperative RT (also known as PORT), stricter DVH parameters should be considered for the lungs. The QUANTEC review provides the most comprehensive estimates from clinical data of dose-response relationships for normal tissue complications.⁵¹⁴⁻⁵¹⁸

The normal tissue dose constraints for conventionally fractionated RT are based on a survey of radiation oncologists at NCCN Member Institutions (see Table 5 in *Principles of Radiation Therapy* in the NCCN Guidelines for NSCLC).^{475,519-524} These constraints are mainly empirical and have not, for the most part, been validated rigorously.^{503,523,525-530} Therefore, the

doses and constraints provided in the tables are not specific prescriptive recommendations; they are useful reference doses that have been commonly used or are from previous clinical trials. A caveat was also added that these constraints represent doses that generally should not be exceeded. Because the risk of toxicity increases progressively with dose to normal tissues, a key principle of radiation treatment planning is to keep normal tissue doses "as low as reasonably achievable" while adequately covering the target. The doses to any given organ at risk should typically be lower than these constraints, approaching them only when there is close proximity to the target volume. After surgery, lung tolerance to RT is much less than for patients with intact lungs; therefore, more conservative constraints should be used for postoperative RT.

For definitive RT, the commonly prescribed dose is 60 to 70 Gy in 2 Gy fractions over 6 to 7 weeks (see *Principles of Radiation Therapy* in the NCCN Guidelines for NSCLC).^{531,532} RTOG 0617, a phase 3 randomized trial, suggests that high-dose radiation using 74 Gy with concurrent chemotherapy does not improve survival, and might be harmful, when compared with a dose of 60 Gy.^{520,533-537} In this trial and subsequent retrospective analyses, higher dose to the heart and specific cardiac substructures, such as the coronary arteries, was associated with increased mortality.⁵³⁷⁻⁵⁴³ Although optimal RT dose intensification remains a valid question, the NCCN Panel does not currently recommend a high dose of 74 Gy for routine use.^{534,536,537,539-547}

General Treatment Information

The RT recommendations for patients with stages I to IV are described in the NSCLC algorithm (see *Principles of Radiation Therapy* in the NCCN Guidelines for NSCLC). Definitive RT, preferably SABR, is recommended for patients with early-stage NSCLC (ie, stage I–II, N0) who are medically inoperable or those who refuse surgery (see *Stereotactic Ablative Radiotherapy* in this Discussion).^{406,407,410,485,548,549} SABR is also an option



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

for patients at high surgical risk who cannot tolerate a lobectomy because of a major medical comorbidity and/or severely limited lung function. After definitive RT (preferably SABR), adjuvant chemotherapy may be considered in patients with high-risk factors for recurrence (eg, large tumor size), similar to the postoperative setting.^{408,550-552} Image-guided thermal ablation (eg, cryotherapy, microwave, radiofrequency ablation [RFA]) is an option for selected patients who are medically inoperable and will not receive SABR or definitive RT (see *Principles of Image-Guided Thermal Ablation Therapy* in the NCCN Guidelines for NSCLC).^{386,553-559} Resection is recommended for patients with early-stage NSCLC who are medically fit (see *Principles of Surgical Therapy* in the NCCN Guidelines for NSCLC).⁵⁶⁰ The indications for using preoperative or postoperative chemoradiation, systemic therapy, or RT alone are described in the NSCLC algorithm (see *Principles of Radiation Therapy* in the NCCN Guidelines for NSCLC). In patients with clinical stage I or II NSCLC who are upstaged to N2+ after surgery, postoperative chemotherapy can be administered, with concurrent postoperative RT for patients with positive margins, or followed by postoperative RT in selected patients with high-risk features.

Definitive chemoradiation is recommended for patients with stage II to III disease who are not appropriate surgical candidates.⁵⁶¹ For patients with locally advanced NSCLC (stage III), the most commonly prescribed conventionally fractionated doses for definitive RT are 60 to 70 Gy in 2 Gy fractions. Doses of at least 60 Gy should be given.⁵⁶² Involved-field RT (also known as involved-field irradiation or IFI) is recommended for treating nodal disease in patients with locally advanced NSCLC (over the older standard of elective nodal irradiation, or ENI).⁵⁶³⁻⁵⁷⁰

The optimal care of patients with potentially operable stage IIIA (N2) NSCLC is controversial and is discussed in detail in the algorithm (see *Principles of Surgical Therapy* in the NCCN Guidelines for

NSCLC).^{447,449,454,461,571} Before surgical resection of stage IIIA NSCLC, induction systemic therapy or induction chemoradiotherapy is recommended to potentially shrink the tumor(s).^{454,465} Preoperative systemic therapy and postoperative RT is an option for patients with resectable stage IIIA NSCLC with minimal N2 disease who can be treated with lobectomy.^{453,466} Preoperative concurrent chemoradiation is recommended for resectable superior sulcus tumors and is an option for other resectable stage IIA to IIIA NSCLC.⁵⁰⁵ In a 2021 survey of the NCCN Member Institutions, we found that 66% use induction chemotherapy, whereas 33% use induction chemoradiation before surgery for patients with stage IIIA N2 disease.^{465,466} The NCCN NSCLC Panel recommends a preoperative RT dose of 45 to 54 Gy in 1.8 to 2 Gy fractions over 5 weeks.^{453,572} Definitive RT doses delivered as preoperative chemo/RT can safely be administered and achieve promising nodal clearance and survival rates;^{506-508,573} the risk of surgical complications after high-dose RT can be minimized with expert thoracic surgical techniques. About 50% of NCCN Member Institutions would consider pneumonectomy after preoperative chemotherapy, whereas only 25% do not consider pneumonectomy absolutely contraindicated after preoperative induction chemoradiation.

Surgery is associated with potentially greater risk of complications, particularly stump breakdown and bronchopleural fistula, in a field that has had high-dose RT (eg, 60 Gy). Thus, surgeons are often wary of resection in areas that have previously received RT doses of more than 45 to 50 Gy, especially in patients who have received definitive doses of preoperative concurrent chemoradiation (ie, ≥60 Gy). Soft tissue flap coverage and reduced intraoperative fluid administration and ventilator pressures can reduce the risk of these complications.⁵⁰⁶⁻⁵⁰⁸ When giving preoperative RT to less than definitive doses (eg, 45 Gy), one should be prepared up front to continue to a full definitive dose of RT without interruption if the patient does not proceed to surgery for some reason.



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

For these reasons, when considering trimodality therapy, the treatment plan—including assessment for resectability and the type of resection—should be decided before initiation of any therapy. In postoperative RT, the clinical target volume (CTV) includes the bronchial stump and high-risk draining lymph node stations.⁵⁷⁴ Standard RT doses after complete resection are 50 to 54 Gy in 1.8 to 2 Gy fractions over 5 to 6 weeks, but a boost may be considered to high-risk regions including areas of nodal extracapsular extension or microscopic positive margins.^{469,575,576} Lung dose constraints should be more conservative, because tolerance appears to be reduced after surgery. The LungART and PORT-C trials provide useful guidelines for postoperative RT technique.^{452,455,577} Highly conformal techniques (such as IMRT or proton therapy) to minimize lung and heart doses are preferred.

For patients with advanced lung cancer (ie, stage IV) with extensive metastases, systemic therapy is recommended; palliative RT can be used for symptom relief and potentially for prophylaxis at primary or distant sites (such as pain, bleeding, or obstruction) (see *Therapy for Recurrence and Metastases* in the algorithm).^{485,578-580} Shorter courses of palliative RT are preferred for patients with symptomatic chest disease who have poor PS and/or shorter life expectancy (eg, 17 Gy in 8.5 Gy fractions in 1–2 weeks), because they provide similar pain relief as longer courses, although there is a higher potential need for retreatment (see Table 4 in the *Principles of Radiation Therapy* in the NCCN Guidelines for NSCLC).⁵⁸¹⁻⁵⁸⁴ Higher dose and longer course thoracic RT (eg, ≥30 Gy in 10 fractions) are associated with modestly improved survival and symptoms, especially in patients with good PS.^{578,585} When higher doses (>30 Gy) are warranted, technologies to reduce normal tissue irradiation may be used (at least 3D-CRT and including IMRT or proton therapy as appropriate).

Oligometastatic disease is heterogenous and refers to limited metastatic sites or disease burden; management is evolving. Definitive local therapy to oligometastases (including brain and lung) achieves prolonged survival in a small proportion of well-selected patients with PS 0 to 2 who have also received radical therapy to the intrathoracic disease.⁵⁸⁶ Definitive RT to oligometastases, particularly SABR, is an appropriate option in such cases if it can be delivered safely to the involved sites.^{587,588} In two randomized phase II trials, significantly longer PFS was found for local consolidative therapy (RT or surgery) to primary and oligometastatic lesions versus maintenance systemic therapy or observation for patients not progressing on systemic therapy.^{589,590} Updated data from one of the trials also show that median overall survival was longer for patients with oligometastatic NSCLC who received local consolidative therapy (median, 41.2 months; 95% CI, 18.9 months–not reached) compared with those receiving maintenance therapy or observation (median, 17.0 months; 95% CI, 10.1–39.8 months; $P = .017$).⁵⁹¹ A phase 2 trial of consolidative RT for oligometastatic NSCLC ($n = 29$) reported median overall survival of 28.4 months (95% CI, 14.5–45.8 months).⁵⁹² The NCCN Guidelines recommend that local therapy (RT, SABR, or surgery) to primary and oligometastatic lesions should be considered for patients without progression on systemic therapy.⁵⁸⁹⁻⁵⁹¹

Stereotactic Ablative Radiotherapy

SABR (also known as SBRT) uses short courses of highly conformal and dose-intensive (ablative) RT precisely delivered to limited-size targets.^{406,593-596} SABR has achieved good primary tumor control rates and overall survival that is higher than conventionally fractionated RT.⁴⁰⁶ Studies, including prospective multi-institutional trials, have demonstrated the efficacy of SABR for patients with inoperable stage I NSCLC or for those who refuse surgery.^{410,597-601} With conventionally fractionated RT, 3-year survival is only about 20% to 35% in these patients, with local failure rates of about 40% to 60%.⁴⁰⁷ In prospective



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

clinical trials, local control and overall survival appear to be considerably increased with SABR, generally more than 85%, and about 60% at 3 years (median survival, 4 years), respectively, in patients who are medically inoperable.^{386,407,501,559,560,600,602-607} A 7-year follow-up of 65 patients with medically inoperable stage I NSCLC reported that overall survival rates were 55.7% at 5 years and 47.5% at 7 years.⁵⁴⁸ In 12 patients (18.5%), a second primary lung carcinoma developed after SABR at a median of 35 months (range, 5–67 months); 27% (18/65) had disease recurrence a median of 14.5 months (range, 4.3–71.5 months) after SABR.

Although SABR is not proven equivalent to lobectomy for patients with operable early-stage NSCLC, some prospective series have shown similar overall survival and cancer-specific survival.^{473,597,599,608} A combined analysis of two randomized trials (that individually did not complete accrual) compared SABR to lobectomy, with 3-year overall survival of 95% and 79%, respectively.⁶⁰⁸ A single-arm SABR expansion cohort of one of the trials with longer follow-up compared non-randomly to a propensity-matched contemporary cohort treated with VATS lobectomy with mediastinal lymph node dissection and found 5-year overall survival of 87% and 84%, respectively.⁴⁷³ These analyses do not provide sufficient data to change the standard of care for good surgical candidates but help to confirm the indication for SABR in patients with relative contraindications to surgery or those who refuse surgery. SABR can also be used for patients with limited lung metastases or limited metastases to other body sites.^{594,601,609-615}

SABR is recommended in the NSCLC algorithm for patients with stage I and II (T1–3,N0,M0) NSCLC who are medically inoperable; SABR is a reasonable alternative to surgery for patients with potentially operable disease who are high risk, older, or refuse surgery after appropriate consultation (see the NCCN Guidelines for NSCLC).^{386,601,603,608,616} If

possible, biopsy should confirm NSCLC before use of SABR.^{406,617,618} Decisions about whether to recommend SABR should be based on multidisciplinary discussion. If therapy is contemplated based on a clinical diagnosis without tissue confirmation of NSCLC, then multidisciplinary evaluation is required to provide consensus on the risk-benefit ratio of biopsy and/or the clinical diagnosis.^{619,620} By analogy with surgical treatment, adjuvant chemotherapy may be considered after definitive RT, preferably SABR, in patients with high-risk factors for recurrence (eg, large tumor size).^{408,550-552}

Locoregional recurrences can occur after SABR.^{560,599,621-626} Late recurrences have been reported more than 5 years after SABR, highlighting the need for careful surveillance.^{627,628} After SABR, assessment of recurrences by imaging can be challenging because of benign inflammatory/fibrotic changes that can remain FDG-PET avid for 2 or more years after treatment, emphasizing the importance of follow-up by a team with experience interpreting such post-treatment effects.^{629,630} This careful follow-up is particularly relevant, because selected patients with recurrences after SABR may benefit from surgery or re-treatment with SABR.^{627,631-635}

SABR fractionation regimens and a limited subset of historically used maximum dose constraints are provided in the NSCLC algorithm; 1 to 5 fractions are generally used (see Tables 2 and 3 in the *Principles of Radiation Therapy* in the NCCN Guidelines for NSCLC).^{406,598,600,607,636-646}

In the United States, only regimens of 5 fractions or less meet the arbitrary billing code definition for SBRT; however, slightly more protracted ablative regimens are also appropriate.^{646,647} Prescription doses do not completely describe the actual delivered doses.^{648,649} These maximum dose constraints are for reference and are not intended to be prescriptive; they are used commonly or have been used in clinical trials. Although none of these dose constraints has been validated as a maximally tolerated dose,



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

outcomes of clinical trials to date suggest that they are safe constraints. The bronchial tree, esophagus, and brachial plexus are critical structures for SABR. For centrally located tumors—defined variably as within 2 cm of the proximal bronchial tree and/or abutting mediastinal pleura—regimens of 54 to 60 Gy in 3 fractions are not safe and should be avoided; 4- to 10-fraction SABR regimens appear to be effective and safe (see *Principles of Radiation Therapy* in the NCCN Guidelines for NSCLC).^{408,639,650-652} The RTOG 0813 trial evaluated 5-fraction regimens and found no high-grade toxicities at 50 Gy in 5 fractions.^{653,654}

SRS or SABR for limited oligometastases to the brain or other body sites, respectively, is recommended for patients with good PS if their thoracic disease can be treated with definitive therapy (see *Stage IVA, M1b* in the NCCN Guidelines for NSCLC).^{396,587,588,601,655-658} SRS or SABR can be considered for select patients with stage M1c disease who have limited metastases that are amenable to treatment with definitive local therapy; clinical trials have included up to 3 to 5 metastatic sites.^{655,656} For patients with disease progression on targeted therapy for *ALK* rearrangements, *ROS1* rearrangements, or the common *EGFR* mutations (ie, exon 19 deletion or exon 21 L858R), consideration of local therapy (eg, surgery or SABR [or SRS]) is recommended for limited progression.⁶⁵⁹⁻⁶⁶² Hypofractionated or dose-intensified conventional 3D-CRT is a less preferred option if an established SABR program is not available.⁶⁶³⁻⁶⁶⁵ Nonrandomized clinical data indicate that local tumor control with SABR is higher than with image-guided thermal ablation techniques. Image-guided thermal ablation (ie, cryotherapy, microwave, radiofrequency) may be an option for selected patients who will not be receiving SABR or definitive RT.^{386,410,559}

Whole Brain RT and Stereotactic Radiosurgery

Many patients with NSCLC have brain metastases (30%–50%), which substantially affect their quality of life.^{6,666} Control of brain metastases

preserves neurocognitive function.^{667,668} However, whole brain RT is itself associated with measurable declines in neurocognitive function in clinical trials, particularly with increasing dose and advanced age of the patient.⁶⁶⁹⁻⁶⁷¹ For limited brain metastases, randomized trials have found that the addition of whole brain RT to SRS decreases intracranial recurrence but does not improve survival and may increase the risk of cognitive decline.^{668,672} Thus, SRS alone is recommended for patients with limited brain metastases.²³⁹ A randomized trial assessed cognitive function in 213 patients with 1 to 3 brain metastases who received SRS alone versus SRS with whole brain RT; most patients had lung cancer.²³⁸ At 3 months after SRS alone, patients had less cognitive deterioration (40/63 patients [63.5%]) than those receiving SRS plus whole brain RT (44/48 patients [91.7%]; difference, -28.2%; 90% CI, -41.9% to -14.4%; $P < .001$). Resection followed by SRS to the cavity (instead of resection followed by whole brain RT) may decrease the risk of neurocognitive impairment.^{673,674} A randomized trial demonstrated that using IMRT to avoid the hippocampus decreases memory impairment after whole brain RT.⁶⁷⁵ A phase 3 randomized trial assessed optimal supportive care (including dexamethasone) with whole brain RT versus optimal supportive care alone in patients with NSCLC and brain metastases who were not eligible for brain surgery or SRS.⁶⁷⁶ Overall survival was similar between the groups (HR, 1.06; 95% CI, 0.90–1.26). Overall quality of life, use of dexamethasone, and reported adverse events were also similar between the arms. Two retrospective analyses have reported increased survival in patients with brain metastases who received SRS and concurrent ICI therapy.^{677,678}

Treatment options for limited brain metastases in patients with NSCLC include: 1) SRS alone; and 2) surgical resection for selected patients followed by SRS or whole brain RT (see the NCCN Guidelines for Central Nervous System Cancers, available at www.NCCN.org). Selected patients include those with symptomatic lesions or whose tumor tissue is



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

needed for diagnosis.^{238,612,666,679-685} Decisions about whether to recommend SRS alone or brain surgery followed by whole brain RT or SRS for limited brain metastases should be based on multidisciplinary discussion, weighing the potential benefit over the risk for each individual patient.^{679,686-688} Treatment should be individualized for patients with recurrent or progressive brain lesions.⁶⁸⁹ For extensive brain metastases, whole brain RT (with hippocampal avoidance if eligible) and memantine are recommended (see the NCCN Guidelines for Central Nervous System Cancers, available at www.NCCN.org).²³⁹⁻²⁴²

Combined Modality Therapy

As previously mentioned, surgery provides the best chance for cure for patients with stage I or II disease who are medically fit and can tolerate surgery. SABR can be considered for patients with medically inoperable or high-risk stage I or II (T1–3,N0) disease or those who refuse surgery if their disease is node negative (see *Stereotactic Ablative Radiotherapy* in this Discussion and see the NCCN Guidelines for NSCLC). In patients with completely resected NSCLC, adjuvant (postoperative) chemotherapy has been shown to improve survival in patients with early-stage disease.⁶⁹⁰⁻⁶⁹³ Some studies suggest that preoperative chemotherapy (also referred to as neoadjuvant chemotherapy or induction chemotherapy) is as effective as and better tolerated than postoperative chemotherapy (see *Preoperative Chemotherapy With or Without Immunotherapy Followed by Surgery* in this Discussion).^{447,694-700} A randomized trial found no difference in survival with preoperative versus postoperative chemotherapy.⁷⁰¹ The NCCN Guidelines state that patients with stage II or IIIA (T3,N1) disease may be treated with induction systemic therapy before surgery if they would have been candidates for adjuvant chemotherapy after surgery.^{386,702} Concurrent chemoradiation is more efficacious than sequential chemoradiation for patients with unresectable stage III disease.⁷⁰³⁻⁷⁰⁶ Cytotoxic chemotherapeutic agents can cause hair loss, which is distressing for patients. Hair loss varies depending on the systemic

regimen and other factors. Data in women with non-metastatic breast cancer suggest that a scalp cooling device may help reduce hair loss in patients receiving cytotoxic chemotherapy regimens.⁷⁰⁷⁻⁷¹¹

For patients with stage IV disease who have a good PS, platinum-based chemotherapy with or without immunotherapy is beneficial.⁷¹²⁻⁷¹⁹ Data show that early palliative care combined with systemic therapy improved quality of life, mood, and survival in patients with metastatic NSCLC, even if these patients had less aggressive end-of-life care, when compared with those receiving systemic therapy alone.^{720,721} Patients should receive treatment for debilitating symptoms.^{6,722,723} A study also suggests that social support, such as being married, is as effective as systemic therapy.⁷²⁴ Data suggest that systematic symptom monitoring during outpatient chemotherapy treatment increases overall survival when compared with usual care.⁷²⁵⁻⁷²⁷ Surgery is rarely recommended for patients with stage IV disease. However, surgical resection of limited brain progression may improve survival in selected patients with stage IV disease and is recommended for selected patients in the NCCN Guidelines (see the NCCN Guidelines for NSCLC).⁷²⁸ Definitive local therapy with surgical resection or RT is recommended for limited single-organ progression located in sites other than the brain if definitive thoracic therapy is feasible (see *Stage IVA, M1b* in the NCCN Guidelines for NSCLC).^{396,586,589,591,601,655,656} The trials supporting the recommendations for combined modality therapy are discussed in the following sections.

Surgery Followed by Adjuvant Therapy: Trial Data and NCCN Recommendations

Chemotherapy

The International Adjuvant Lung Cancer Trial (IALT) assessed cisplatin-based postoperative therapy in patients with completely resected stage I, II, or III NSCLC.⁶⁹¹ The study included 1867 patients with



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

surgically resected lung cancer who were randomly assigned either to cisplatin-based postoperative chemotherapy or to observation, with a median follow-up duration of 56 months. The survival rate at 5 years was 45% for cisplatin-based therapy versus 40% for observation (HR for death, 0.86; 95% CI, 0.76–0.98; $P < .03$); the disease-free survival rate was 39% versus 34% at 5 years (HR, 0.83; 95% CI, 0.74–0.94; $P < .003$). However, after 7.5 years of follow-up, there were more deaths in the chemotherapy group and the benefit of chemotherapy decreased over time.⁷²⁹ Data show that postoperative chemotherapy prevents recurrences.

The NCIC CTG JBR.10 trial and the ANITA trial compared the effectiveness of postoperative vinorelbine plus cisplatin versus observation in early-stage NSCLC. In the JBR.10 trial, 482 patients (ECOG PS of 0–1) with completely resected stage IB (T2a,N0) or stage II (T1,N1, or T2,N1) NSCLC were randomly assigned either to vinorelbine plus cisplatin or to observation.⁶⁹² Postoperative chemotherapy significantly prolonged overall survival compared with observation alone (94 vs. 73 months; HR for death, 0.69; $P = .04$) and relapse-free survival (not reached vs. 47 months, HR for recurrence, 0.60; $P < .001$). The 5-year survival rates were 69% and 54%, respectively ($P = .03$). When compared with observation alone, postoperative chemotherapy was beneficial for patients with stage II disease but not for stage IB disease as shown by updated data from JBR.10 after 9 years of follow-up.⁷³⁰ In patients with stage II disease receiving postoperative chemotherapy, median survival was 6.8 versus 3.6 years in those who were only observed. Of note, patients receiving chemotherapy did not have an increased death rate.

In the ANITA trial, 840 patients with stage IB (T2a,N0), II, or IIIA NSCLC were randomly assigned either to postoperative vinorelbine plus cisplatin or to observation.⁶⁹³ Grade 3 and 4 toxicities were manageable in the chemotherapy group; seven toxic deaths were reported. After a median follow-up of 76 months, median survival was 66 months in the

chemotherapy group and 44 months in the observation group.⁶⁹³ Postoperative chemotherapy significantly improved (8.6%) the 5-year overall survival in patients with completely resected stage II and IIIA disease, although no benefit was observed in stage I. Some clinicians consider vinorelbine plus cisplatin to be the preferred regimen for completely resected early-stage NSCLC based on the number of trials and the amount of use;⁷³¹ however, most clinicians in the United States prefer to use regimens with less toxicity.^{732,733}

A meta-analysis of 4584 patients (LACE) found that postoperative cisplatin-based chemotherapy increased survival over 5 years (absolute benefit of 5.4%); there was no difference among the chemotherapy regimens (vinorelbine, etoposide, and others).⁷³⁴ A subgroup analysis found that cisplatin plus vinorelbine also increased survival.⁷³¹ The benefit was greater in patients with stage II and III disease and with good PS. Postoperative chemotherapy benefited older patients up to 80 years of age.^{391,735}

The CALGB 9633 trial assessed paclitaxel plus carboplatin in patients with stage IB (T2a,N0,M0) lung cancer.⁷³⁶⁻⁷³⁸ In this trial, 344 patients were randomly assigned either to paclitaxel plus carboplatin or to observation (within 4–8 weeks of resection) with a median follow-up duration of 74 months. Postoperative chemotherapy was well tolerated with no chemotherapy-related toxic deaths. Overall survival at 6 years was not significantly different (however, a subset analysis showed a benefit for tumors ≥ 4 cm), although 3-year survival was significant (80% vs. 73%, $P = .02$).^{737,738} Thus, the carboplatin and paclitaxel regimen is only recommended for early-stage disease if patients cannot tolerate cisplatin (see *Perioperative Systemic Therapy* in the NCCN Guidelines for NSCLC).⁷³⁹ It is important to note that the CALGB trial was underpowered for patients with stage 1B disease.⁷⁴⁰



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

TREAT, a phase 2 randomized study, assessed cisplatin plus pemetrexed versus cisplatin plus vinorelbine as postoperative therapy for patients with completely resected stages IB to III NSCLC.⁷³² Data showed that cisplatin plus pemetrexed was an effective, less toxic regimen compared with cisplatin plus vinorelbine; in addition, patients were able to receive more cycles of cisplatin plus pemetrexed compared with cisplatin plus vinorelbine.⁷³² Overall survival at 3 years was similar between the arms (75% vs. 77%; $P = .858$).⁷⁴¹

The NCCN NSCLC Panel does not recommend postoperative chemotherapy for resected stage IA disease based on the trials described in the previous paragraphs.⁷⁴² Postoperative chemotherapy is recommended for high-risk, margin-negative, stage IB disease. Recommended chemotherapy regimens for preoperative and postoperative (perioperative) chemotherapy for patients with completely resected stages IB to III NSCLC are provided in the NCCN Guidelines; the regimens also include specific dosing (see *Perioperative Systemic Therapy* in the NCCN Guidelines for NSCLC).^{690,742}

The NCCN NSCLC Panel has preference stratified all the systemic therapy regimens and decided that cisplatin plus pemetrexed is the preferred neoadjuvant or adjuvant option for nonsquamous NSCLC if patients are not candidates for ICIs (see *Perioperative Systemic Therapy* in the NCCN Guidelines for NSCLC).^{732,741} Cisplatin plus gemcitabine or cisplatin plus docetaxel are the preferred perioperative chemotherapy options for patients with squamous cell NSCLC.^{743,744} Other recommended options include cisplatin plus vinorelbine or cisplatin plus etoposide.⁶⁹¹⁻⁶⁹³ Perioperative therapy options for patients who are not candidates for cisplatin-based therapy are preference stratified as useful in certain circumstances and include: 1) carboplatin plus paclitaxel; 2) carboplatin plus gemcitabine; and 3) carboplatin plus pemetrexed (but only for nonsquamous NSCLC).⁷⁴⁵⁻⁷⁴⁸ Neoadjuvant therapy is also known as

preoperative (or induction) therapy; adjuvant therapy is also known as postoperative therapy.

Targeted Therapy

ADAURA, a phase 3 randomized trial, assessed adjuvant therapy with osimertinib versus placebo in 682 patients with resected stage IB to IIIA *EGFR* mutation-positive NSCLC.⁷⁴⁹ At 24 months, 89% (95% CI, 85%–92%) of the osimertinib group and 52% (95% CI, 46%–58%) of the placebo group were alive and disease-free (overall HR for disease recurrence or death, 0.20; 95% CI, 0.14–0.30; $P < .001$). At 24 months, 98% (95% CI, 95%–99%) of the osimertinib group and 85% (95% CI, 80%–89%) of the placebo group were alive and did not have CNS disease (overall HR for disease recurrence or death, 0.18; 95% CI, 0.10–0.33). For patients with stage IB NSCLC, 88% (95% CI, 78%–94%) of the osimertinib group and 71% (95% CI, 60%–80%) of the placebo group were alive and disease-free at 24 months (overall HR for disease recurrence or death, 0.39; 95% CI, 0.18–0.76). For patients with stage II NSCLC, 91% (95% CI, 82%–95%) of the osimertinib group and 56% (95% CI, 45%–65%) of the placebo group were alive and disease-free at 24 months (overall HR, 0.17; 95% CI, 0.08–0.31). For those with stage IIIA NSCLC, 88% (95% CI, 79%–94%) of the osimertinib group and 32% (95% CI, 23%–41%) of the placebo group were alive and disease-free at 24 months (overall HR, 0.12; 95% CI, 0.07–0.20). Data were also reported for those who received adjuvant chemotherapy and those who did not; 60% (410/682) of patients received adjuvant chemotherapy.⁷⁵⁰ Overall survival data are not yet available. Serious adverse events were reported in 16% (54/339) of patients receiving osimertinib versus 12% (42/343) receiving placebo. Ten patients (3%) receiving osimertinib had interstitial lung disease versus no patients in the placebo group. There were no deaths in the osimertinib group versus one in the placebo group.



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

The NCCN Panel recommends osimertinib as an adjuvant therapy option for eligible patients with completely resected (R0) stage IB to IIIA, and stage IIIB (only T3,N2) *EGFR* mutation-positive NSCLC who have previously received adjuvant chemotherapy or are ineligible to receive platinum-based chemotherapy based on clinical trial data and FDA approval.⁷⁴⁹ Osimertinib is recommended in this setting for *EGFR* exon 19 deletions or *EGFR* exon 21 L858R mutations, which are the most common *EGFR* mutations. Although it is technically easier to use a resected specimen to do molecular testing, it is also acceptable to use initial diagnostic biopsy specimens. For the 2023 update (Version 1), the NCCN Panel expanded the criteria to include patients with stage IIIB (only T3,N2) NSCLC based on differences between the AJCC staging systems for the 7th and 8th editions (see Table 3 in the algorithm).¹⁶² Although the 8th edition of the AJCC staging manual is currently being used, the clinical trial used the 7th edition.

Immunotherapy

Trial Data

IMpower010, a phase 3 randomized trial, assessed adjuvant therapy with atezolizumab versus best supportive care in 1005 patients with resected early-stage NSCLC and various PD-L1 levels who had received adjuvant chemotherapy.⁷⁵¹ In patients with resected stage II to IIIA NSCLC and PD-L1 of 1% or more, disease-free survival was improved in those receiving adjuvant atezolizumab compared with best supportive care (HR, 0.66; 95% CI, 0.5–0.88; *P* = .0039). Treatment-related grade 3 and 4 adverse events were reported in 11% (53/495) of patients; 4 deaths occurred (1%, 4/495). Atezolizumab is an ICI that inhibits PD-L1 receptors.

PEARLS/KEYNOTE 091, a phase 3 randomized trial, assessed adjuvant therapy with pembrolizumab versus placebo in 1177 patients with completely resected stage IB to IIIA NSCLC (based on AJCC 7th edition).³² Most patients (86%) received adjuvant platinum-based chemotherapy after

resection but had not received neoadjuvant chemotherapy or radiotherapy. In patients who received adjuvant chemotherapy after resection, the median disease-free survival was 53.6 months (95% CI, 39.2–not reached) in those receiving adjuvant pembrolizumab compared with 42.0 months (95% CI, 31.3–not reached) in those receiving placebo (HR, 0.76; 95% CI, 0.63–0.91). Grade 3 or worse events included hypertension (6%) and pneumonia (2%). In patients receiving pembrolizumab, there were four fatal (1%) treatment-related adverse reactions, including myocarditis and cardiogenic shock, septic shock and myocarditis, pneumonia, and sudden death. Pembrolizumab is an ICI that inhibits PD-1 receptors.

NCCN Recommendations

The NCCN Panel recommends atezolizumab as an adjuvant therapy option for eligible patients with completely resected (R0) stage IIB to IIIA, stage IIIB (only T3,N2), or high-risk stage IIA NSCLC and with PD-L1 of 1% or more who are negative for certain biomarkers—*EGFR* exon 19 deletions, *EGFR* exon 21 L858R mutations, and *ALK* rearrangements—and who have previously received adjuvant chemotherapy based on clinical trial data and FDA approval.⁷⁵¹ For the 2023 update (Version 2), the NCCN Panel added a recommendation for adjuvant pembrolizumab for eligible patients with completely resected stage IIB to IIIA, stage IIIB (only T3,N2), or high-risk stage IIA NSCLC who are negative for *EGFR* exon 19 deletions, *EGFR* exon 21 L858R mutations, or *ALK* fusions and who have previously received adjuvant chemotherapy.³² The panel added a caveat that the benefit of adjuvant pembrolizumab is unclear for patients with PD-L1 levels less than 1%.

For the 2023 update (Version 1), the NCCN Panel expanded the molecular testing criteria to include *ALK* rearrangements because patients with *ALK* rearrangements, *EGFR* exon 19 deletions, or *EGFR* exon 21 L858R mutations have less benefit from ICIs.¹⁸⁶ Clinical trials are assessing whether *ALK* inhibitors are effective in this setting.⁷⁵²⁻⁷⁵⁸ The



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

NSCLC Panel also clarified that single-agent therapy with osimertinib is recommended if patients have both *EGFR* mutations and PD-L1 of 1% or more; the panel does not recommend using single-agent therapy with atezolizumab or combination therapy with osimertinib plus atezolizumab in this setting. The NCCN Panel also added a recommendation that patients should be evaluated for perioperative (neoadjuvant or adjuvant) therapy before surgery. For the 2023 update (Version 1), the NCCN Panel expanded the criteria to include patients with stage IIIB (only T3,N2) disease as previously described.

Preoperative Chemotherapy With or Without Immunotherapy Followed by Surgery

Trial Data

Data from clinical trials in patients with resected NSCLCs indicate that delivery of chemotherapy is an important problem. In the postoperative setting, significant comorbidities and incomplete recovery after surgery often make it difficult for patients to tolerate systemic therapy. This problem was demonstrated in NATCH, a phase 3 randomized trial—which compared surgery alone to preoperative or postoperative chemotherapy with paclitaxel/carboplatin—because 90% of the preoperative cohort completed 3 cycles of chemotherapy but only 61% of the postoperative cohort completed chemotherapy; however, survival was equivalent among all three arms.⁶⁹⁹ IFCT 0002, a randomized trial, found no difference in 3-year overall survival (67.4% vs. 67.7%) with preoperative versus postoperative chemotherapy in patients with early-stage NSCLC; response rate and quality of life were similar in both arms.⁷⁰¹

Several trials suggest that preoperative therapy is beneficial in patients with N2 disease.^{447,454,698} Other trials suggest that preoperative therapy is beneficial in patients with earlier stage disease.^{695,696,700} A randomized intergroup trial (SWOG 9900) evaluated preoperative paclitaxel plus carboplatin in 354 patients with stage IB to IIIA (but not N2) disease

versus surgery alone. The trial closed prematurely because of practice changes and was therefore not appropriately powered. This SWOG trial did show a trend toward improved PFS (33 vs. 20 months) and overall survival (62 vs. 41 months) with preoperative chemotherapy, and no difference in resection rates between the two arms.⁷⁰⁰

Scagliotti et al published a phase 3 trial of preoperative cisplatin plus gemcitabine versus surgery alone in 270 patients with stage IB to IIIA disease. Although the trial closed early, a significant survival benefit was seen in patients with stages IIB and IIIA disease who received preoperative chemotherapy (HR, 0.63).⁶⁹⁵ Song et al published a meta-analysis of 13 randomized clinical trials evaluating preoperative chemotherapy followed by surgery versus surgery alone in resectable NSCLCs (overall survival: HR, 0.84; 95% CI, 0.77–0.92; $P = .0001$).⁶⁹⁴ These results are similar to those reported in another meta-analysis (HR, 0.89; 95% CI, 0.81–0.98; $P = .02$).⁶⁹⁵ The benefit from preoperative chemotherapy is similar to that attained with postoperative chemotherapy.^{695,699,701,734}

CheckMate 816, a phase 3 randomized trial, assessed neoadjuvant therapy with nivolumab plus platinum-doublet chemotherapy versus chemotherapy alone in 358 patients with resectable (tumors ≥ 4 cm or node positive) NSCLC (stage IB–IIIA resectable NSCLC using AJCC staging, 7th edition).¹¹⁰ Chemotherapy regimens included 1) cisplatin plus either pemetrexed (nonsquamous only) or gemcitabine (squamous only); or 2) carboplatin plus paclitaxel (any histology). If their patients could not tolerate cisplatin, clinicians could substitute carboplatin-based regimens. Patients had good PS (0–1) and did not have *EGFR* or *ALK* alterations. The median event-free survival was 31.6 months (95% CI, 30.2–not reached) for nivolumab plus chemotherapy versus 20.8 months (95% CI, 14.0–26.7) for chemotherapy alone (HR, 0.63; 97% CI, 0.43–0.91; $P = .005$). The pathologic complete response was 24% (95% CI, 18%–



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

31%) with nivolumab plus chemotherapy versus 2.2% (95% CI, 0.6%–5.6%) with chemotherapy alone (odds ratio 13.9; 99% CI, 3.49–55.7; $P<.001$). The major pathologic response was 36.9% with nivolumab plus chemotherapy versus 8.9% with chemotherapy alone (major pathologic response, $\leq 10\%$ viable tumor in lung and lymph nodes). The overall response rate was 53.6% for nivolumab plus chemotherapy versus 37.4% with chemotherapy alone. Surgery was done for 83% of patients receiving nivolumab plus chemotherapy versus 75% of those receiving chemotherapy alone. Grade 3 to 4 treatment-related adverse events occurred in 33.5% of patients receiving nivolumab plus chemotherapy versus 36.9% of those receiving chemotherapy alone.

NADIM, a single-arm phase 2 study, assessed neoadjuvant therapy with nivolumab plus paclitaxel plus carboplatin in 46 patients with surgically resectable stage IIIA NSCLC; 41 patients had resection.^{759,760} At 36 months, updated results show that overall survival was 81.9% (95% CI, 66.8%–90.6%).⁷⁵⁹ TMB and PD-L1 levels were not predictive of survival. Grade 3 or higher adverse events were reported in 30% (14/46) of patients including increased lipase (3/46, 7%) and febrile neutropenia (3/46, 7%). None of the adverse events were associated with death or delays in surgery.

NCCN Recommendations

Based on clinical trial data, the NCCN Panel recommends that certain patients who are likely to receive adjuvant chemotherapy may instead be treated with preoperative (also known as induction or neoadjuvant) systemic therapy after surgical evaluation.^{695,699,701,734} The NCCN Panel recommends nivolumab plus platinum-doublet chemotherapy as a neoadjuvant systemic therapy option for eligible patients with resectable (tumors ≥ 4 cm or node positive) NSCLC and no contraindications to treatment with PD-1/PD-L1 inhibitors based on clinical trial data and FDA approval.^{110,760} For the 2023 update (Version 1), the panel added a

recommendation that all patients should be evaluated for neoadjuvant therapy with strong consideration for those with node-positive disease or tumors 4 cm or more and no contraindications for ICIs. However, neoadjuvant therapy should not be used to attempt to induce resectability in patients who do not already meet criteria for resectability on initial evaluation. Some oncogenic drivers—such as *EGFR* exon 19 deletions, *EGFR* exon 21 L858R mutations, or *ALK* rearrangements—have been shown to be associated with less benefit from PD-1/PD-L1 inhibitors.¹⁸⁶ Testing for PD-L1 status, *EGFR* mutations, and *ALK* rearrangements is recommended before administering neoadjuvant nivolumab plus platinum-doublet chemotherapy in eligible patients with stage IB (only tumors = 4 cm) to IIIA, or stage IIIB (only T3, N2) NSCLC. For the 2023 update (Version 1), the NCCN Panel expanded the criteria to include patients with stage IIIB (only T3,N2) NSCLC as previously described. The chemotherapy regimens that may be used with neoadjuvant nivolumab include: 1) cisplatin plus either pemetrexed (nonsquamous only), gemcitabine (squamous only), or paclitaxel (any histology); or 2) carboplatin plus either pemetrexed (nonsquamous only), gemcitabine (squamous only), or paclitaxel (any histology).¹¹⁰

Chemoradiation: Trial Data and NCCN Recommendations

The major controversies in NSCLC relate to the care of patients with stage IIIA disease (see the *Role of Surgery in Patients with Stage IIIA (N2) NSCLC* in *Principles of Surgical Therapy* in the NCCN Guidelines for NSCLC). All three treatment modalities—surgical resection, chemotherapy, and radiation—may be used when treating stage III disease. The ongoing debate centers on which modalities to use and in what sequence.⁷⁶¹⁻⁷⁶⁶

Two randomized trials assessed postoperative RT in patients with clinical stage I or II disease that was upstaged surgically to N2.^{452,455} LungART, a phase 3 randomized trial, assessed postoperative RT versus no



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

postoperative RT in 501 patients with completely resected stage IIIA (N2) NSCLC who had received either neoadjuvant (14%) or adjuvant chemotherapy (75%).⁴⁵² The median disease-free survival was 30.5 months (95% CI, 24–49) in the postoperative RT group versus 22.8 months (95% CI, 17–37) in the control group (HR, 0.86; 95% CI, 0.68–1.08; $P = .18$). Common grade 3 to 4 adverse events included pneumonitis (postoperative RT: 5% [13/241]; control: <1% [1/246]); lymphopenia (postoperative RT: 4% [9/241]; control: 0); and fatigue (postoperative RT: 3% [6/241]; control: <1%). Late-grade 3 to 4 cardiopulmonary toxicity occurred in 11% (26/241) of patients in the postoperative RT group versus 5% (12/246) in the control group. Three patients died in the postoperative RT group (2, pneumonitis; 1, sepsis).

PORT-C, a phase 3 randomized trial, assessed postoperative RT versus no postoperative RT in 394 patients with completely resected stage IIIA (N2) NSCLC who had received adjuvant chemotherapy.⁴⁵⁵ The 3-year disease-free survival rates were 40.5% with postoperative RT versus 32.7% with no RT (median, 22.1 vs. 18.6 months) (HR, 0.84; 95% CI, 0.65–1.09; $P = .20$). The 3-year overall survival rates were 78.3% in patients receiving postoperative RT versus 82.8% with no RT (HR, 1.02; $P = .93$). The 3-year local recurrence only rates were 9.5% and 18.3%, respectively (Fine-Gray HR, 0.55; Gray test $P = .04$). No radiotherapy-related grade 4 or 5 adverse events were reported.

Data from the LungART and PORT-C trials show that postoperative RT (also known as PORT) did not improve survival compared with no postoperative RT, although locoregional control was significantly improved.^{452,455} For the 2023 update (Version 1), the NCCN Panel clarified that postoperative RT may be considered for select patients with high-risk N2 disease such as extracapsular extension, multi-station involvement, inadequate lymph node dissection or sampling, and/or refusal of or intolerance to adjuvant systemic therapy.^{452,455,761,767} To minimize

potential lung and heart toxicities, highly conformal RT techniques such as IMRT or proton therapy are preferred.^{468,532,768,769}

For patients with unresectable stage IIIA or stage IIIB disease, combined modality therapy (chemoradiation) is more efficacious than RT alone.^{762,763,765,766,770} Concurrent chemoradiation is more efficacious than sequential chemoradiation.^{703-706,771} However, concurrent chemoradiation has a higher rate of grade 3 or 4 esophagitis than sequential chemoradiation. Selection of patients should be based not only on the anticipated response to therapy but also on how well the patient is anticipated to tolerate therapy. Accelerated RT regimens may be useful if concurrent chemoradiation would not be tolerated.^{772,773} Sequential chemoradiation or RT alone is recommended for frail patients who cannot tolerate concurrent chemoradiation.^{389,774}

JCOG0301, a phase 3 randomized trial, assessed chemoradiation using low-dose carboplatin versus RT alone in patients (>70 years) with unresectable NSCLC.⁷⁷⁵ Median overall survival was 22.4 months (95% CI, 16.5–33.6) for chemoradiotherapy with carboplatin and 16.9 months (95% CI, 13.4–20.3) for RT alone (HR, 0.68; 95.4% CI, 0.47–0.98; $P = .0179$). In the chemoradiation group, 3% (3/100) of patients died, whereas 4% (4/100) of patients died in the RT group. Grade 3 to 4 hematologic effects occurred at a greater rate in the chemoradiation arm than in the RT alone arm, including leucopenia (61 [63.5%] vs. none), neutropenia (55 [57.3%] vs. none), and thrombocytopenia (28 [29.2%] vs. 2 [2.0%]). In older patients receiving chemoradiation versus RT alone, long-term follow-up data for overall survival continue to support using chemoradiation (HR, 0.74; 95% CI, 0.55–0.99; $P = .024$).⁷⁷⁶ Although the optimal sequence is not established, postoperative RT is generally administered after adjuvant chemotherapy or concurrently for positive resection margins.^{468,470,471,768}



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

Concurrent chemoradiation regimens recommended for all histologies for initial treatment include 1) cisplatin plus etoposide; and 2) carboplatin plus paclitaxel (see *Concurrent Chemoradiation Regimens* in the NCCN Guidelines for NSCLC).^{534,703,705,777-782} For nonsquamous NSCLC, additional recommended concurrent chemoradiation regimens include pemetrexed plus either carboplatin or cisplatin.⁷⁸³⁻⁷⁸⁵ A weekly paclitaxel plus carboplatin regimen is another chemoradiation option.⁵³⁴ The different options for preoperative, definitive, and postoperative chemotherapy/RT are described in detail in the algorithm. The NCCN NSCLC Panel has preference stratified all the systemic therapy regimens and decided that the following concurrent chemoradiation options are preferred for patients with NSCLC: 1) pemetrexed plus either carboplatin or cisplatin for nonsquamous NSCLC only; 2) carboplatin plus paclitaxel for all histologies; and 3) cisplatin plus etoposide for all histologies. The cisplatin plus vinblastine concurrent regimen is rarely used in the United States; therefore, it is not recommended in the NCCN Guidelines.

The recommended sequential chemoradiation options include cisplatin combined with pemetrexed (nonsquamous only), docetaxel, etoposide, gemcitabine, or vinorelbine. Carboplatin regimens are recommended for patients with comorbidities or those not able to tolerate cisplatin, including 1) carboplatin plus gemcitabine; 2) carboplatin plus paclitaxel; and 3) carboplatin plus pemetrexed (nonsquamous only). The NCCN NSCLC Panel has preference stratified all the systemic therapy regimens and decided that the following sequential chemoradiation options are preferred for patients with NSCLC: 1) pemetrexed plus cisplatin for nonsquamous NSCLC only; 2) cisplatin plus gemcitabine for squamous histology; and 3) cisplatin plus docetaxel for squamous histology.

Durvalumab

Durvalumab is a human ICI antibody that inhibits PD-L1 (see *PD-L1 Expression Levels* and *Immune Checkpoint Inhibitors* in this

Discussion).^{360-362,365} PACIFIC, a phase 3 randomized trial, compared adjuvant treatment with durvalumab (also known as consolidation immunotherapy in this setting) versus placebo in eligible patients with unresectable stage III NSCLC (PS 0–1) but without disease progression after treatment with 2 or more cycles of definitive concurrent platinum-based chemoradiation.^{365,786} Eligible patients received adjuvant durvalumab after treatment with concurrent chemoradiation (1–42 days). Most patients either currently or previously smoked cigarettes and did not have *EGFR* mutations; their PD-L1 status was typically less than 25% or unknown. Grade 3 or 4 adverse events occurred at a similar rate in both groups of patients (durvalumab, 30.5% vs. placebo, 26.1%). Pneumonia was the most common grade 3 or 4 adverse event (durvalumab, 4.4% vs. placebo, 3.8%). Durvalumab did not compromise patient-reported outcomes.⁷⁸⁷ An updated analysis of this trial reported that overall survival was increased after durvalumab consolidation (47.5 months; 95% CI, 38.4–52.6) compared with placebo (29.1 months; 95% CI, 22.1–35.1) (stratified HR for death, 0.72; 95% CI, 0.45–0.68).^{19,786} After 5 years, 42.9% of patients who received durvalumab were alive versus 33.4% of those who received placebo.¹⁶ In addition, 42.9% were alive after 5 years if they had received durvalumab and 33% were alive and free of disease progression compared with 19% of those who received placebo. The subgroup analyses were limited by small sample size and were not powered to assess efficacy. Patients receiving durvalumab received less subsequent immunotherapy (11.6%) compared with those who received placebo (28.3%).

The NCCN NSCLC Panel recommends durvalumab (category 1) as a consolidation immunotherapy option (regardless of PD-L1 status) for eligible patients (PS 0–1) with unresectable stage III NSCLC and without disease progression after treatment with definitive concurrent platinum-based chemoradiation based on this trial and FDA approval.^{365,786} It is important to note that durvalumab is used as an adjuvant treatment



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

option in this setting; it is not used as second-line therapy. Durvalumab may be used as a consolidation immunotherapy option after treatment with any of the concurrent chemoradiation regimens described in the algorithm (eg, cisplatin plus etoposide, carboplatin plus paclitaxel) (see *Concurrent Chemoradiation Regimens* in the NCCN Guidelines for NSCLC). Ten patients with stage II NSCLC were included in the PACIFIC trial, which used the older AJCC staging criteria (7th edition). Therefore, the NCCN NSCLC Panel also recommends durvalumab as a consolidation immunotherapy option (regardless of PD-L1 status) for eligible patients (PS 0–1) with unresectable stage II NSCLC and without disease progression after treatment with definitive concurrent platinum-based chemoradiation.

A few of the recommended concurrent chemoradiation regimens have an option for additional cycles (2 or 4) of chemotherapy after radiation, which is termed consolidation chemotherapy (see *Concurrent Chemoradiation Regimens* in the NCCN Guidelines for NSCLC).^{505,534,778,783} If patients will be receiving durvalumab, the NCCN NSCLC Panel does not recommend consolidation chemotherapy based on concerns that adding consolidation chemotherapy will increase the risk of pneumonitis if patients are also receiving durvalumab. Durvalumab should be discontinued for patients with severe or life-threatening pneumonitis and should be withheld or discontinued for other severe or life-threatening immune-mediated adverse events when indicated (see prescribing information). If patients will not be receiving durvalumab because of medical contraindications or other reasons, consolidation chemotherapy is an option after concurrent chemoradiation, depending on the initial regimen.

Chemotherapy: Trial Data and NCCN Recommendations

Patients with metastatic (stage IV) NSCLC (who have a good PS) benefit from chemotherapy, usually with a platinum-based regimen, which was the only treatment option for many years before the advent of targeted therapy

and immunotherapy regimens.⁷¹⁶⁻⁷¹⁸ If patients are not eligible for the targeted therapy or immunotherapy regimens, then chemotherapy regimens are recommended. Combination chemotherapy regimens produce 1-year survival rates of 30% to 40% and are more efficacious than single agents.^{739,744,788-790} However, survival rates are higher for patients with stage IV NSCLC who are eligible for either the newer targeted therapy or immunotherapy regimens.^{12,17-30} Phase 3 randomized trials have shown that many of the platinum-doublet combinations yield similar objective response rates and survival.^{791,792} The platinum-doublet regimens differ slightly for toxicity, convenience, and cost; thus, clinicians can individualize therapy for their patients.^{744,793-799} Non-platinum-based regimens are also options, such as 1) gemcitabine plus vinorelbine, or 2) gemcitabine plus docetaxel.⁸⁰⁰⁻⁸⁰³ The prognosis for stage IV inoperable lung cancer remains poor if patients are not candidates for targeted therapy or immunotherapy regimens.

In the United States, frequently used initial cytotoxic regimens for stage IV nonsquamous NSCLC include: 1) cisplatin (or carboplatin) plus pemetrexed; or 2) carboplatin plus paclitaxel with (or without) bevacizumab.^{777,804,805} Gemcitabine plus cisplatin (or carboplatin) is often used for patients with stage IV squamous cell NSCLC.^{790,795,804,805} These chemotherapy regimens are recommended based on phase 3 randomized trials (see *Systemic Therapy for Advanced or Metastatic Disease* in the NCCN Guidelines for NSCLC).^{790,806} A phase 3 randomized trial suggests that conventional cytotoxic agents should not be continued beyond 4 to 6 cycles of therapy; however, many patients assigned to a longer duration of therapy did not receive the planned number of cycles (see *Maintenance Therapy* in this Discussion).^{807,808}

A phase 3 randomized trial assessed cisplatin plus pemetrexed versus cisplatin plus gemcitabine as first-line therapy in patients with stage IIIB or IV NSCLC.⁷⁹⁰ For patients with adenocarcinoma who received cisplatin



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

plus pemetrexed, median overall survival was 12.6 months compared with 10.9 months for those receiving cisplatin plus gemcitabine (HR, 0.84; 95% CI, 0.71–0.99; $P = .03$). In contrast, for patients with squamous cell NSCLC who received cisplatin plus pemetrexed, overall survival was 9.4 versus 10.8 months for those receiving cisplatin plus gemcitabine (HR, 1.23; 95% CI, 1.00–1.51; $P = .05$). Patients with nonsquamous NSCLC receiving cisplatin plus pemetrexed had less toxicity when compared with those receiving cisplatin plus gemcitabine.⁸⁰⁹ Median overall survival was similar for both regimens when histologies were combined (8.6 vs. 9.2 months, respectively; HR, 1.08; 95% CI, 0.81–1.45; $P = .586$).

TAX 326, a phase 3 randomized trial, assessed docetaxel plus cisplatin (or carboplatin) versus vinorelbine plus cisplatin as first-line therapy for patients with stage IIIB or IV NSCLC.⁷⁴⁴ Docetaxel plus cisplatin was associated with similar overall survival (11.3 vs. 10.1 months; $P = .044$; HR, 1.183; 97.2% CI, 0.989–1.416) and better response rate (31.6%) when compared with cisplatin plus vinorelbine (24.5%; $P = .029$); docetaxel plus cisplatin was associated with better quality of life and was better tolerated.

Many oncologists use pemetrexed-based regimens for stage IV adenocarcinomas (if patients are not candidates for targeted therapy or PD-1/PD-L1 inhibitors), because taxane-based regimens are associated with more toxicity (eg, neurotoxicity).^{790,810} There are no agents for the prevention of peripheral neuropathy, and few agents are useful for treatment.⁸¹¹ The POINTBREAK trial showed that carboplatin plus pemetrexed plus bevacizumab is a reasonable option for patients with metastatic NSCLC and confirmed that taxane-based regimens are more toxic than pemetrexed-based regimens.⁸¹² The POINTBREAK trial also showed that both regimens are similar in regard to overall survival rates; therefore, oncologists may return to using taxane-based regimens, which are well established. A retrospective cohort study suggests that the

addition of bevacizumab to carboplatin plus paclitaxel does not increase survival in patients ≥ 65 years with advanced nonsquamous NSCLC.⁸¹³ However, another retrospective cohort study reported increased survival in older patients.⁸¹⁴ A combined analysis of the ECOG 4599 and POINTBREAK trials found a survival benefit with the addition of bevacizumab to carboplatin plus paclitaxel in patients < 75 years but no benefit in those > 75 years.⁸¹⁵

Albumin-bound paclitaxel (also known as nab-paclitaxel) can be substituted for paclitaxel or docetaxel for patients: 1) who have experienced hypersensitivity reactions after receiving paclitaxel or docetaxel despite premedication; or 2) in whom premedications (ie, dexamethasone, H2 blockers, H1 blockers) to prevent hypersensitivity are contraindicated.^{816,817} A phase 3 randomized trial in patients with advanced NSCLC reported that an albumin-bound paclitaxel plus carboplatin regimen is associated with less neurotoxicity and improved response rate, when compared with the control arm of paclitaxel plus carboplatin.⁸¹⁸ Based on the trial and FDA approval, the NCCN NSCLC Panel recommends an albumin-bound paclitaxel plus carboplatin regimen as initial cytotoxic therapy for patients with advanced NSCLC and good PS.

Chemotherapy is recommended for patients with stage IV NSCLC and negative test results for actionable driver mutations (ie, *ALK*, *BRAF* p.V600E, *ERBB2*, *EGFR*, *KRAS*, *MET*ex14 skipping, *NTRK1/2/3*, *RET*, and *ROS1*), PD-L1 expression less than 1%, and contraindications to PD-1 or PD-L1 inhibitors. Chemotherapy regimens may be used as first-line or subsequent therapy options in eligible patients with actionable driver mutations as shown in the algorithm, although they are generally not preferred options. Targeted therapy is a preferred first-line therapy option for patients with actionable driver mutations, except for *EGFR* exon 20 mutations or *KRAS* mutations. Recommended



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

chemotherapy regimens are based on PS and include platinum agents (eg, cisplatin, carboplatin), taxanes (eg, paclitaxel, albumin-bound paclitaxel, docetaxel), vinorelbine, etoposide, pemetrexed, and gemcitabine. To clarify use of systemic therapy, the NCCN Guidelines list all the combination systemic therapy regimens and single agents that are recommended for patients with metastatic NSCLC based on histology and PS (see *Systemic Therapy for Advanced or Metastatic Disease* in the NCCN Guidelines for NSCLC).

For patients with advanced NSCLC who have a PS of 2, platinum-based combinations and a few single-agent chemotherapy agents are recommended in the NCCN Guidelines; cisplatin-based regimens are not recommended in this setting.⁸¹⁹ For nonsquamous NSCLC or NSCLC NOS, single-agent chemotherapy includes gemcitabine, pemetrexed, or taxanes; combination chemotherapy regimens include carboplatin plus paclitaxel or carboplatin plus pemetrexed.⁸²⁰⁻⁸²² Patients with a PS of 2 are often just treated with single-agent chemotherapy because of concerns about toxicity.⁸²³ Treatment with carboplatin plus pemetrexed increased median overall survival when compared with pemetrexed alone (9.3 vs. 5.3 months, $P = .001$) in patients with a PS of 2; however, 4 treatment-related deaths occurred in the carboplatin plus pemetrexed arm.^{820,824}

The NCCN NSCLC Panel has preference stratified all the systemic therapy regimens for patients with NSCLC. The newer chemotherapy plus pembrolizumab regimens are preferred for eligible patients with metastatic NSCLC who do not have contraindications to immunotherapy and are not candidates for targeted therapy (see *Systemic Therapy for Advanced or Metastatic Disease* in the NCCN Guidelines for NSCLC and *Immune Checkpoint Inhibitors* in this Discussion). For patients with metastatic nonsquamous NSCLC and PS 0 to 1 who have contraindications to immunotherapy, the panel decided that the following

chemotherapy regimens are “useful in certain circumstances,” including 1) carboplatin with paclitaxel (or albumin-bound paclitaxel), docetaxel, etoposide, gemcitabine, or pemetrexed; all are category 1; 2) cisplatin with paclitaxel (or albumin-bound paclitaxel), docetaxel, etoposide, gemcitabine, or pemetrexed; all are category 1; 3) bevacizumab with carboplatin and either paclitaxel or pemetrexed; and 4) gemcitabine with either docetaxel or vinorelbine. The panel also preference stratified the regimens for patients with metastatic nonsquamous NSCLC and PS 2; carboplatin plus pemetrexed is preferred for patients with adenocarcinoma. The regimens for patients with metastatic squamous cell NSCLC have also been preference stratified.

The initial cytotoxic systemic therapy regimens do not include options that are less effective, more toxic, and/or infrequently used in the United States based on each panel member’s experience and data generated by surveying the NCCN NSCLC Panel (see the NCCN Guidelines with Evidence Blocks™ for NSCLC, available at www.NCCN.org). For patients with metastatic nonsquamous NSCLC and NSCLC NOS, the following regimens are not recommended in the NCCN Guidelines: carboplatin plus vinorelbine, cisplatin plus vinorelbine, etoposide, irinotecan, and vinorelbine. For patients with metastatic squamous cell NSCLC, the following regimens are not recommended in the NCCN Guidelines: carboplatin plus vinorelbine, cisplatin plus gemcitabine plus necitumumab, cisplatin plus vinorelbine, etoposide, irinotecan, and vinorelbine.

The NCCN NSCLC Panel voted unanimously to not recommend the necitumumab plus cisplatin plus gemcitabine regimen for patients with metastatic squamous cell NSCLC. The NCCN NSCLC Panel feels the addition of necitumumab to the regimen is not beneficial based on toxicity, cost, and limited improvement in efficacy when compared with cisplatin plus gemcitabine. A phase 3 randomized trial only showed a slight improvement in overall survival (11.5 months; 95% CI, 10.4–12.6; vs. 9.9



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

months; 95% CI, 8.9–11.1).⁸²⁵ The stratified HR was 0.84 (95% CI, 0.74–0.96; $P = .01$). In addition, there were more grade 3 or higher adverse events in patients receiving the necitumumab regimen (388 [72%] of 538 patients) than in patients receiving only gemcitabine plus cisplatin (333 [62%] of 541). Although it has been suggested that adding necitumumab to cisplatin plus gemcitabine adds value and is cost-effective, the NCCN NSCLC Panel does not agree.⁸²⁶

Targeted Therapies

Specific targeted therapies are available for the treatment of eligible patients with metastatic NSCLC.^{184,827,828} Afatinib, adagrasib, alectinib, brigatinib, cabozantinib, capmatinib, ceritinib, crizotinib, dabrafenib, dacomitinib, entrectinib, erlotinib, gefitinib, larotrectinib, lorlatinib, osimertinib, pralsetinib, selpercatinib, sotorasib, tepotinib, and trametinib are oral TKIs. Alectinib inhibits *ALK* and *RET* rearrangements.⁸²⁹ Brigatinib inhibits various *ALK* rearrangements and other targets.⁸³⁰ Ceritinib inhibits *ALK* and *ROS1* rearrangements. Crizotinib inhibits *ALK* rearrangements, *ROS1* rearrangements, and *MET* tyrosine kinases (ie, high-level *MET* amplification, *MET*ex14 skipping mutation). Erlotinib, gefitinib, afatinib, and dacomitinib inhibit *EGFR* mutations (eg, exon 19 deletions, exon 21 L858R, exon 18 G719X, exon 21 L861Q, exon 20 S768I); osimertinib inhibits these *EGFR* mutations and T790M.²⁵² Lorlatinib inhibits *ALK* and *ROS1* rearrangements.^{357,358,831-833} Dabrafenib inhibits *BRAF* p.V600E mutations; trametinib inhibits MEK; both agents inhibit different kinases in the RAS/RAF/MEK/ERK pathway.^{190,191} Entrectinib and larotrectinib inhibit *TRK* fusion proteins.^{330,333,834} Capmatinib and tepotinib inhibit several *MET* tyrosine kinases including *MET*ex14 skipping mutations and high-level *MET* amplification.^{169,172,326,835} Selpercatinib, pralsetinib, and cabozantinib inhibit *RET* rearrangements.³³⁹⁻³⁴² Adagrasib and sotorasib inhibit *KRAS* G12C mutations.^{317,319} Bevacizumab and ramucirumab are recombinant monoclonal antibodies that target the vascular endothelial growth factor (VEGF) or VEGF receptor, respectively. Cetuximab is a monoclonal

antibody that targets *EGFR*. Other targeted therapies are being developed (see *Emerging Biomarkers to Identify Novel Therapies for Patients with Metastatic NSCLC* in the NCCN Guidelines for NSCLC). Flare phenomenon may occur in some patients who discontinue targeted therapies for driver mutations such as *ALK*, *EGFR*, *MET*ex14 skipping, *RET*, or *ROS1*. If disease flare occurs, then the targeted therapies should be restarted.⁸³⁶⁻⁸³⁹

Targeted therapies are recommended for patients with metastatic NSCLC and specific oncogenic drivers, independent of PD-L1 levels. Patients with metastatic NSCLC and PD-L1 expression levels of 1% or more—but who also have a targetable driver oncogene molecular variant (ie, *EGFR* exon 19 deletions, *EGFR* exon 21 L858R mutations, *ALK* rearrangements)—should receive first-line targeted therapy for that oncogene and not first-line ICIs, because targeted therapies yield higher response rates (eg, osimertinib, 80%) than ICIs (poor response rates) in the first-line setting, targeted therapy is better tolerated, and these patients are unlikely to respond to ICIs.^{186,374,840} Molecular testing results for the actionable oncogenic mutations should be known before starting systemic therapy with ICI regimens in eligible patients with advanced NSCLC, if clinically feasible. If it is not feasible to do molecular testing, then patients are treated as though they do not have driver oncogenes.^{186,373-375,840} If molecular testing results are pending and patients require an urgent start to therapy, clinicians should consider holding immunotherapy for one cycle (ie, just use platinum-based chemotherapy regimens). Monitoring is recommended during initial therapy with response assessment with CT, with or without contrast, of known or high-risk sites of disease after 2 cycles and then every 2 to 4 cycles. Likewise, monitoring is also recommended during maintenance or subsequent therapy with CT, with or without contrast, every 6 to 12 weeks.



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

Oral TKIs that Inhibit ALK Rearrangements

About 5% of patients with NSCLC have *ALK* gene rearrangements. The NCCN NSCLC Panel recommends *ALK* rearrangement testing (category 1) in all patients with metastatic nonsquamous NSCLC or NSCLC NOS based on data showing the efficacy of several agents for patients with *ALK* rearrangements (see *Principles of Molecular and Biomarker Analysis* in the NCCN Guidelines for NSCLC). *ALK* testing can be considered in patients with metastatic squamous cell carcinoma. The NCCN NSCLC Panel recommends five agents for patients with *ALK*-positive metastatic NSCLC—alectinib, brigatinib, ceritinib, crizotinib, and lorlatinib—based on clinical trial data and FDA approvals, which are described in the following sections.

Alectinib

Alectinib is a second-generation oral TKI that inhibits *ALK* rearrangements.⁸²⁹

First-Line Therapy

ALEX, a phase 3 randomized trial, assessed first-line therapy with alectinib versus crizotinib in 303 patients with *ALK*-positive advanced NSCLC including those with asymptomatic CNS disease.²²⁰ Disease progression or death occurred in fewer patients receiving alectinib (41% [62/152]; median follow-up of 18.6 months) when compared with crizotinib (68% [102/151]; median follow-up of 17.6 months). The HR was 0.47 (95% CI, 0.34–0.65; $P < .001$) for disease progression or death. PFS was significantly increased with alectinib (68.4%; 95% CI, 61.0%–75.9%) versus crizotinib (48.7%; 95% CI, 40.4%–56.9%). The median PFS was not reached for alectinib (95% CI, 17.7–not reached) when compared with crizotinib at 11.1 months (95% CI, 9.1–13.1). Fewer patients receiving alectinib had CNS progression (12% [18/152]) versus crizotinib (45% [68/151]). Response rates were 83% (126/152) in the alectinib group versus 75% (114/151) in the crizotinib group ($P = .09$). Patients receiving

alectinib had fewer grade 3 to 5 adverse events when compared with crizotinib (41% [63/152] vs. 50% [75/151], respectively) even though patients received alectinib for a longer duration than crizotinib (median, 17.9 vs. 10.7 months). Fewer deaths were reported with alectinib (3.3% [5/152]) versus crizotinib (4.6% [7/151]); two treatment-related deaths were reported in the crizotinib arm and none in the alectinib arm.

J-ALEX, a phase 3 randomized trial, assessed first-line therapy with alectinib versus crizotinib in 207 Japanese patients with *ALK*-positive advanced NSCLC.⁸⁴¹ Median PFS was not reached with alectinib (95% CI, 20.3 months–not reached) versus 10.2 months (95% CI, 8.2–12.0) with crizotinib (HR, 0.34; 99.7% CI, 0.17–0.71; stratified log-rank $P < .0001$). Grade 3 or 4 adverse events were less frequent with alectinib (26% [27/103]) when compared with crizotinib (52% [54/104]); adverse events did not lead to death in either group. Fewer patients stopped taking alectinib (9%) because of an adverse event when compared with crizotinib (20%).

The NCCN NSCLC Panel recommends alectinib as a first-line therapy option for patients with *ALK*-positive metastatic NSCLC based on clinical trial data and FDA approval.^{220,841,842} Panel members voted that alectinib is a preferred first-line therapy option for patients with *ALK*-positive metastatic NSCLC based on these trials. Alectinib is a category 1 (preferred) option if an *ALK* rearrangement is discovered before giving first-line systemic therapy; alectinib is a preferred monotherapy option if an *ALK* rearrangement is discovered during first-line systemic therapy (eg, carboplatin plus paclitaxel). Brigatinib, ceritinib, crizotinib, and lorlatinib are also recommended as first-line therapy options in patients with *ALK*-positive NSCLC. The NCCN NSCLC Panel has preference stratified the first-line therapy regimens and decided that alectinib, brigatinib, and lorlatinib are all preferred first-line therapy options for patients with



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

ALK-positive metastatic NSCLC. Ceritinib is an “other recommended” option, whereas crizotinib is useful in certain circumstances.

Subsequent Therapy

Phase 2 trials assessed alectinib in patients with *ALK*-positive metastatic NSCLC and disease progression on crizotinib; overall response rates were 48% to 50%.^{182,843} In the larger trial (138 patients), patients on alectinib had a response rate of 50% (95% CI, 41%–59%), and median duration of response of 11.2 months (95% CI, 9.6–not reached).¹⁸² For CNS disease, the control rate was 83% (95% CI, 74%–91%) and the median duration of response was 10.3 months (95% CI, 7.6–11.2). Of 84 patients with baseline CNS metastases, 23 (27%) had a complete CNS response to alectinib. Of 23 patients with baseline CNS metastases and no previous brain RT, 10 (43%) had a complete CNS response to alectinib. Most adverse events were only grade 1 to 2 (constipation, fatigue, and peripheral edema); 4 patients (3%) had grade 3 dyspnea. One death due to intestinal perforation may have been related to alectinib. The NCCN NSCLC Panel recommends alectinib as a subsequent therapy option for patients with *ALK*-positive metastatic NSCLC and disease progression after crizotinib based on these trials and FDA approval.^{182,842,843} Brigatinib, ceritinib, and lorlatinib are also recommended as subsequent therapy options in patients with *ALK*-positive NSCLC and disease progression after crizotinib. Patients who do not tolerate crizotinib may be switched to alectinib, brigatinib, ceritinib, or lorlatinib unless an adverse side effect requiring discontinuation has occurred (eg, pneumonitis).

Brigatinib

Brigatinib is a second-generation oral TKI that inhibits *ALK* rearrangements.

First-Line Therapy

ALTA-1L, a phase 3 randomized trial, assessed brigatinib versus crizotinib as first-line therapy for patients with *ALK*-positive metastatic NSCLC.²²³ At the first interim analysis, PFS was increased in patients receiving brigatinib (67%; 95% CI, 56%–75%) versus those receiving crizotinib (43%; 95% CI, 32%–53%) (HR for disease progression or death, 0.49; 95% CI, 0.33–0.74; $P < .001$). Intracranial response was also increased with brigatinib (78%; 95% CI, 52%–94%) versus crizotinib (29%; 95% CI, 11%–52%). At the second interim analysis (24.9 months of follow-up), brigatinib continued to show improved PFS when compared with crizotinib (HR 0.43; 95% CI, 0.31–0.61; median, 29.4 vs. 9.2 months).⁸⁴⁴ After 3 years, the PFS was 43% for brigatinib versus 19% for crizotinib (HR, 0.48; 95% CI, 0.35–0.66). The median overall survival was not reached in either arm.⁸⁴⁵

The NCCN NSCLC Panel recommends brigatinib as a first-line therapy option for patients with *ALK*-positive NSCLC based on clinical trial data and FDA approval.²²³ Brigatinib is a category 1 (preferred) option if an *ALK* rearrangement is discovered before giving first-line systemic therapy. Brigatinib is a preferred option if an *ALK* rearrangement is discovered during first-line systemic therapy (eg, carboplatin plus paclitaxel). Other first-line therapy options include alectinib, ceritinib, crizotinib, and lorlatinib for patients with *ALK*-positive NSCLC. The NCCN NSCLC Panel has preference stratified the first-line therapy regimens and decided that alectinib, brigatinib, and lorlatinib are all preferred options for patients with *ALK*-positive metastatic NSCLC; ceritinib is an “other recommended” option and crizotinib is useful in certain circumstances.⁸⁴⁴

Subsequent Therapy

ALTA, a phase 2 study, assessed two different doses of brigatinib—90 mg (arm A) or 180 mg (arm B) every day—in patients with *ALK*-positive metastatic NSCLC and disease progression on, or who were intolerant to, crizotinib.^{846,847} The overall response rates were 45% (97% CI, 34%–56%)



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

and 54% (97% CI, 43%–65%) in arms A and B, respectively. Many patients had brain metastases (71% and 67%, respectively). The intracranial overall response rates were 42% (11/26) and 67% (12/18), respectively, in patients with measurable brain metastases. The median PFS was 9.2 months (95% CI, 7.4–15.6) and 12.9 months (95% CI, 11.1–not reached), respectively. Grade 3 or higher adverse events included hypertension (6% and 6%, respectively) and pneumonia (3% and 5%, respectively).

The NCCN NSCLC Panel recommends brigatinib as a subsequent therapy option for patients with *ALK*-positive NSCLC and disease progression after crizotinib based on clinical trial data and FDA approval.^{846,847} Patients receiving brigatinib should be carefully monitored for respiratory symptoms, especially during the first week of treatment. Alectinib, ceritinib, and lorlatinib are also recommended as subsequent therapy options in patients with *ALK*-positive NSCLC and disease progression after crizotinib. Patients who do not tolerate crizotinib may be switched to alectinib, brigatinib, ceritinib, or lorlatinib unless an adverse side effect requiring discontinuation has occurred (eg, pneumonitis).

Ceritinib

Ceritinib is a second-generation oral TKI that inhibits *ALK* and *ROS1* rearrangements.⁸⁴⁸

First-Line Therapy

ASCEND-4, a phase 3 randomized trial, assessed ceritinib versus platinum-based chemotherapy as first-line therapy for patients with *ALK*-positive metastatic NSCLC.²²² The median PFS was 16.6 months (95% CI, 12.6–27.2) for ceritinib and 8.1 months (95% CI, 5.8–11.1) for chemotherapy (HR, 0.55; 95% CI, 0.42–0.73; $P < .00001$). For ceritinib, common adverse events included diarrhea (85% [160/189] of patients), nausea (69% [130/189]), vomiting (66% [125/189]), and an increase in

ALT (60% [114/189]). For chemotherapy, common adverse events included nausea (55% [97/175 patients]), vomiting (36% [63/175]), and anemia (35% [62/175]).

The NCCN NSCLC Panel recommends ceritinib as a first-line therapy option for patients with *ALK*-positive metastatic NSCLC based on clinical trial data and FDA approval. Ceritinib is a category 1 (other recommended) option if an *ALK* rearrangement is discovered before giving first-line systemic therapy; ceritinib is a treatment if an *ALK* rearrangement is discovered during first-line systemic therapy (eg, carboplatin plus paclitaxel). Alectinib, brigatinib, crizotinib, and lorlatinib are also recommended as first-line therapy options in patients with *ALK*-positive NSCLC. The NCCN NSCLC Panel has preference stratified the first-line therapy regimens and decided that alectinib, brigatinib, and lorlatinib are all preferred options for patients with *ALK*-positive metastatic NSCLC; ceritinib is an “other recommended” option and crizotinib is useful in certain circumstances.

Subsequent Therapy

ASCEND-5, a phase 3 randomized trial, assessed subsequent therapy with ceritinib versus chemotherapy (with pemetrexed or docetaxel) in patients with advanced *ALK*-positive NSCLC who had previously received at least two or more treatments (including chemotherapy and crizotinib) and had disease progression.⁸⁴⁹ The median PFS for ceritinib was 5.4 months (95% CI, 4.1–6.9) versus a median PFS of 1.6 months for chemotherapy (95% CI, 1.4–2.8; HR, 0.49; 95% CI, 0.36–0.67; $P < .0001$). Serious adverse events were reported in 43% (49/115) of patients receiving ceritinib versus 32% (36/113) of those receiving chemotherapy. ASCEND-2, a phase 2 study, assessed ceritinib in patients who had previously received at least two or more treatments, with disease progression on crizotinib, and with brain metastases.⁸⁵⁰ The overall response rate was 38%; the duration of response was 9.7 months (95%



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

CI, 7.1–11.1).⁸⁵⁰ The intracranial overall response rate was 45.0% (95% CI, 23.1%–68.5%).

The NCCN NSCLC Panel recommends ceritinib as a subsequent therapy option for patients with *ALK*-positive metastatic NSCLC and disease progression after crizotinib based on clinical trial data and FDA approval.⁸⁴⁹ Alectinib, brigatinib, and lorlatinib are also recommended as subsequent therapy options in patients with *ALK*-positive NSCLC and disease progression after crizotinib. Patients who do not tolerate crizotinib may be switched to alectinib, brigatinib, ceritinib, or lorlatinib unless an adverse side effect requiring discontinuation has occurred (eg, pneumonitis).

Crizotinib

Crizotinib is a first-generation oral TKI that inhibits *ALK* rearrangements, *ROS1* rearrangements, and some *MET* tyrosine kinases (high-level *MET* amplification or *MET*ex14 skipping mutation).^{192,221,320,851-855}

First-Line Therapy

Randomized phase 3 trials have compared crizotinib with first-line chemotherapy (PROFILE 1014) and with subsequent chemotherapy (PROFILE 1007) for patients with *ALK*-positive metastatic NSCLC.^{10,221,856}

First-line therapy with crizotinib improved PFS, response rate (74% vs. 45%; $P < .001$), lung cancer symptoms, and quality of life when compared with chemotherapy (pemetrexed with either cisplatin or carboplatin).²²¹

Crizotinib yields high response rates (>60%) when used in patients with advanced NSCLC who have *ALK* rearrangements, including those with brain metastases.^{123,221,857-859} Patients whose disease responds to crizotinib may have rapid improvement in symptoms; median time to progression on crizotinib is about 7 months to 1 year.^{860,861} Crizotinib has relatively few side effects (eg, eye disorders, edema, transient changes in

renal function).^{858,862,863} However, some patients have had pneumonitis; crizotinib should be discontinued in these patients.⁸⁵³ Patients who do not tolerate crizotinib may be switched to alectinib, brigatinib, ceritinib, or lorlatinib unless an adverse side effect requiring discontinuation has occurred (eg, pneumonitis).

The NCCN NSCLC Panel recommends crizotinib as a first-line treatment option for patients with *ALK* rearrangement-positive metastatic NSCLC based on clinical trial data and FDA approval.²²¹ Crizotinib is a category 1 (useful in certain circumstances) option if an *ALK* rearrangement is discovered before giving first-line systemic therapy; crizotinib is a treatment option if an *ALK* rearrangement is discovered during first-line systemic therapy (eg, carboplatin plus paclitaxel). Alectinib, brigatinib, ceritinib, and lorlatinib are also recommended as first-line therapy options in patients with *ALK*-positive NSCLC. The NCCN NSCLC Panel has preference stratified the first-line therapy regimens and decided that alectinib, brigatinib, and lorlatinib are all preferred options for patients with *ALK*-positive metastatic NSCLC, whereas ceritinib is an “other recommended” option and crizotinib is useful in certain circumstances. Patients who do not tolerate crizotinib may be switched to alectinib, brigatinib, ceritinib, or lorlatinib unless an adverse side effect requiring discontinuation has occurred (eg, pneumonitis).

Subsequent Therapy

Subsequent therapy with crizotinib improved PFS (7.7 vs. 3.0 months; $P < .001$) and response rate (65% vs. 20%; $P < .001$) when compared with either docetaxel or pemetrexed in patients with *ALK*-positive NSCLC and disease progression after first-line chemotherapy who had not previously received *ALK* inhibitors.⁸⁵² Therefore, crizotinib may also be continued for patients with *ALK* rearrangements and disease progression on crizotinib, depending on the type of progression.⁸⁵² The NCCN NSCLC Panel does not recommend continuing crizotinib for patients with brain metastases



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

and disease progression after first-line therapy with crizotinib; the other ALK inhibitors are recommended options in this setting because they have better CNS response rates (ie, alectinib, brigatinib, ceritinib, lorlatinib).^{849,850,864,865}

Lorlatinib

Lorlatinib is an oral third-generation TKI that targets ALK and ROS1 tyrosine kinases and has good CNS penetration; it inhibits a broad range of *ALK* resistance mutations that develop after treatment with first- and second-generation ALK inhibitors.^{358,831-833}

First-Line Therapy

CROWN, a phase 3 randomized trial, assessed lorlatinib versus crizotinib as first-line therapy for patients with *ALK*-positive metastatic NSCLC.²¹⁹ At 12 months, 78% (95% CI, 70%–84%) of patients were alive without disease progression in the lorlatinib group versus 39% (95% CI, 30%–48%) in the crizotinib group (HR for disease progression or death, 0.28; 95% CI, 0.19–0.41; $P < .001$). The objective response rate was 76% (95% CI, 68%–83%) for patients receiving lorlatinib and 58% (95% CI, 49%–66%) for those receiving crizotinib. For patients with measurable brain metastases, 82% (95% CI, 57%–96%) of those receiving lorlatinib had an intracranial response and 23% (95% CI, 5%–54%) of those receiving crizotinib responded. Of patients who received lorlatinib, 71% had a complete intracranial response. Hyperlipidemia, edema, increased weight, peripheral neuropathy, and cognitive effects were the most common adverse events seen with lorlatinib. More grade 3 or 4 adverse events (mainly altered lipid levels) occurred with lorlatinib than with crizotinib (72% vs. 56%, respectively). Discontinuation of treatment because of adverse events was similar in both groups (7% for lorlatinib and 9% for crizotinib). Updated data from the CROWN study show that in patients with brain metastases at baseline, the 12-month cumulative incidence of CNS progression was 7% with lorlatinib compared with 72% with

crizotinib; the 12-month PFS rates were 78% versus 22%, respectively.⁸⁶⁶ In patients without brain metastases at baseline, the 12-month cumulative incidence of CNS progression was 1% with lorlatinib compared with 18% with crizotinib; the 12-month PFS rates were 78% versus 45%, respectively.⁸⁶⁶

The NCCN NSCLC Panel recommends lorlatinib as a first-line therapy option for patients with *ALK*-positive metastatic NSCLC based on clinical trial data and FDA approval.^{219,866} Panel members voted that lorlatinib is a preferred first-line therapy option for patients with *ALK*-positive metastatic NSCLC based on trial data.²¹⁹ Lorlatinib is a category 1 (preferred) option if an *ALK* rearrangement is discovered before giving first-line systemic therapy; lorlatinib is a preferred option if an *ALK* rearrangement is discovered during first-line systemic therapy (eg, carboplatin plus paclitaxel). Alectinib, brigatinib, ceritinib, and crizotinib are also recommended as first-line therapy options in patients with *ALK*-positive metastatic NSCLC. The NCCN NSCLC Panel has preference stratified the first-line therapy regimens and decided that alectinib, brigatinib, and lorlatinib are all preferred options for patients with *ALK*-positive metastatic NSCLC; ceritinib is an “other recommended” option and crizotinib is useful in certain circumstances.

Subsequent Therapy

Data show that lorlatinib is effective in select patients with disease progression after treatment with ALK inhibitors, including those with CNS metastases.^{831,832} A phase 2 trial assessed lorlatinib in patients with *ALK*-positive or *ROS1*-positive metastatic NSCLC and disease progression after ALK inhibitor therapy; many patients had asymptomatic CNS metastases.⁸³¹ In patients who had received at least one previous ALK inhibitor, objective responses were achieved in 47% of patients (93/198; 95% CI, 39.9%–54.2%); there were 4 complete responses and 89 partial responses. In those with measurable baseline CNS lesions, an



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

objective intracranial response was observed in 63% of patients (51/81; 95% CI, 51.5%–73.4%). Lorlatinib was effective in patients who had received up to three previous ALK inhibitors. Grade 3 to 4 adverse events included hypercholesterolemia and hypertriglyceridemia (43/275 [16%] for both). Serious treatment-related adverse events occurred in 7% of patients (19/275) including cognitive effects in 1% (2/275); the cognitive effects resulted in permanent discontinuation of lorlatinib. No treatment-related deaths were reported. Data from this phase 2 trial also show that lorlatinib is effective as subsequent therapy in patients with the resistance mutation *ALK* G1202R, which is often detected after progression on second-generation ALK TKIs such as brigatinib, alectinib, or ceritinib.⁸³³ The objective response rate with lorlatinib was 62% when using plasma ctDNA (and 69% when using tissue) for patients with *ALK* resistance mutations and disease progression on second-generation ALK TKIs compared with 32% (plasma) and 27% (tissue) in patients without *ALK* mutations.

The NCCN NSCLC Panel recommends lorlatinib as a subsequent therapy option for select patients with *ALK*-positive metastatic NSCLC and disease progression after treatment with ALK inhibitors based on clinical trial data and FDA approval.^{832,833} Lorlatinib is recommended as a subsequent therapy option for select patients with *ALK* G1202R-positive metastatic NSCLC after progression on alectinib, brigatinib, or ceritinib (depending on the type of progression) based on clinical trial data.⁸³³ At progression, the panel recommends considering plasma and/or tissue-based testing using broad molecular profiling for genomic resistance mechanisms.⁸³³ Lorlatinib is recommended as a subsequent therapy option for select patients with *ALK*-positive metastatic NSCLC after progression on crizotinib. If not previously given, lorlatinib is also recommended as a subsequent therapy option for *ALK*-positive metastatic NSCLC after progression on crizotinib followed by progression on either alectinib, brigatinib, or ceritinib.

Oral Agents that Inhibit *BRAF* Mutations

The *BRAF* p.V600E mutation occurs in 1% to 2% of patients with lung adenocarcinoma; it is the most common of the *BRAF* point mutations when considered across all tumor types. Rare *BRAF* mutations include p.V600K and p.V600D. The NCCN NSCLC Panel recommends *BRAF* mutation testing in all patients with metastatic nonsquamous NSCLC or NSCLC NOS based on data showing the efficacy of several agents for patients with *BRAF* mutations (see *Principles of Molecular and Biomarker Analysis* in the NCCN Guidelines for NSCLC).⁸⁶⁷⁻⁸⁷⁰ *BRAF* mutation testing can be considered in patients with metastatic squamous cell carcinoma.

Dabrafenib and Trametinib

Dabrafenib and trametinib inhibit kinases in the RAS/RAF/MEK/ERK pathway.^{190,191} Dabrafenib inhibits *BRAF* p.V600E mutations; trametinib inhibits MEK 1/2, which is downstream of *BRAF* signaling.

First-Line Therapy

A phase 2 trial assessed first-line combination therapy with dabrafenib plus trametinib for 36 patients with metastatic NSCLC and *BRAF* p.V600E mutations.^{18,867} The overall response rate was 64% (23/36; 95% CI, 46%–79%); there were 2 complete responses. The median PFS was 10.9 months (95% CI, 7.0–16.6). Many patients (69% [25/36]) had one or more grade 3 or 4 adverse events. Serious adverse events included increased alanine aminotransferase (ALT) (14% [5/36]), increased aspartate transaminase (AST) (8% [3/36]), pyrexia (11% [4/36]), and decreased ejection fraction (8% [3/36]). An updated analysis reported that patients receiving dabrafenib plus trametinib had a median overall survival of 17.3 months (95% CI, 12.3–40.2).¹⁸ After 5 years, the overall survival rate was 22%.



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

The NCCN NSCLC Panel recommends combination therapy with dabrafenib plus trametinib as a preferred first-line therapy option for patients with metastatic NSCLC and *BRAF* p.V600E mutations based on these trials and FDA approval.^{867,868,870} Single-agent therapy with dabrafenib or vemurafenib is also an option for patients with *BRAF* p.V600E mutations who do not tolerate combination therapy with dabrafenib plus trametinib.^{191,868,871} Other systemic therapy regimens are also recommended options for patients with *BRAF* p.V600E mutations; the same initial systemic regimens used for patients with metastatic NSCLC may be used (eg, carboplatin plus paclitaxel). The NCCN NSCLC Panel has preference stratified the first-line therapy options for patients with

BRAF p.V600E mutation-positive metastatic NSCLC and decided that: 1) dabrafenib plus trametinib is the preferred option; 2) other systemic therapy regimens (eg, carboplatin plus paclitaxel) are other recommended options; and 3) single-agent dabrafenib or vemurafenib are useful in certain circumstances. For the 2023 update (Version 1), the NCCN Panel revised the preference stratification for the other systemic therapy options to “other recommended options” from “useful in certain circumstances.”

Subsequent Therapy

A phase 2 study assessed the dabrafenib plus trametinib regimen as subsequent therapy in 57 patients with advanced NSCLC and *BRAF* p.V600E mutations and disease progression on chemotherapy.^{18,190,869} The overall response rate was 68% (23/36; 95% CI, 55%–80%); PFS was 9.7 months (6.9–19.6). Serious adverse events occurred in 56% (32/57) of patients, including pyrexia, anemia, confused state, hemoptysis, hypercalcemia, and cutaneous squamous cell carcinoma. Grade 3 to 4 adverse events included neutropenia in 9% of patients (5/57), hyponatremia in 7% (4/57), and anemia in 5% (3/57). Four patients died during the study, but these deaths were not felt to be related to treatment (deaths were due to retroperitoneal hemorrhage, subarachnoid

hemorrhage, respiratory distress, or severe disease progression). An updated analysis reported that the median overall survival was 18.2 months (95% CI, 14.3–28.6).¹⁸ After 5 years, the overall survival rate was 19%.

The NCCN NSCLC Panel recommends dabrafenib plus trametinib as a subsequent therapy option if patients with *BRAF* p.V600E mutations have disease progression after first-line systemic therapy regimens (eg, carboplatin plus paclitaxel) and have not received *BRAF* p.V600E inhibitors as first-line therapy.^{190,869} Single-agent therapy with dabrafenib or vemurafenib is also an option for patients with *BRAF* p.V600E mutations who do not tolerate combination therapy with dabrafenib plus trametinib.^{191,868,871} Clinicians should be aware of common adverse events that may occur with dabrafenib plus trametinib including pyrexia, vomiting, and nausea along with less frequent and unique adverse events, such as cutaneous, ocular, and hemorrhagic events.⁸⁷²

Agents that Inhibit *EGFR* Mutations

Oral TKIs that Inhibit *EGFR* Exon 19 Deletions and Exon 21 L858R Mutations

The NCCN NSCLC Panel recommends *EGFR* mutation testing (category 1) in all patients with metastatic nonsquamous NSCLC or NSCLC NOS based on data showing the efficacy of several agents for patients with *EGFR* mutations.^{252,258} *EGFR* mutation testing can be considered in patients with metastatic squamous cell carcinoma. The NCCN NSCLC Panel recommends afatinib, dacomitinib, erlotinib (± bevacizumab or ramucirumab), gefitinib, and osimertinib for patients with metastatic NSCLC and *EGFR* exon 19 deletions or *EGFR* exon 21 L858R mutations based on clinical trial data and FDA approvals, which are described in the following sections.

Afatinib



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

Afatinib is a second-generation oral TKI that irreversibly inhibits the ErbB/HER family of receptors including EGFR and ERBB2.^{873,874} LUX-Lung 3, a phase 3 randomized trial, reported that first-line therapy with afatinib improved PFS when compared with cisplatin plus pemetrexed in patients with metastatic adenocarcinoma who had *EGFR* mutations (11.1 vs. 6.9 months, $P = .001$).²⁵¹ The NCCN NSCLC Panel recommends afatinib as a first-line therapy option in patients with metastatic NSCLC and *EGFR* exon 19 deletions or *EGFR* exon 21 L858R mutations based on the clinical trial and FDA approval.^{251,873,875-877} Afatinib is a category 1 (other recommended) option if an *EGFR* exon 19 deletion or *EGFR* exon 21 L858R mutation is discovered before giving first-line systemic therapy. Afatinib is an option if an *EGFR* exon 19 deletion or *EGFR* exon 21 L858R mutation is discovered during first-line systemic therapy (eg, carboplatin plus paclitaxel).

The NCCN NSCLC Panel has preference stratified the systemic therapy regimens for patients with metastatic NSCLC and *EGFR* exon 19 deletions or *EGFR* exon 21 L858R mutations and decided that afatinib is an “other recommended” option; osimertinib is the preferred option in this setting. Afatinib may also be continued in patients with disease progression if they do not have multiple systemic symptomatic lesions (see *Continuation of Targeted Therapy After Progression on Initial Therapy* in this Discussion).²⁴⁶ However, afatinib is not recommended as subsequent therapy in patients with metastatic squamous cell NSCLC but without *EGFR* mutations based on a phase 3 randomized trial showing low response rates (11%); it is less efficacious and safe compared to other available options [see *Second-Line and Beyond (Subsequent) Systemic Therapy* in this Discussion].⁸⁷⁸ Afatinib is also recommended for eligible patients with metastatic NSCLC and less common *EGFR* mutations including *EGFR* S768I, L861Q, and/or G719X mutations (see *Oral TKIs that Inhibit EGFR S768I, L861Q, and G719X Alterations* in this Discussion).⁸⁷⁹

A phase 2B trial assessed afatinib compared with gefitinib for first-line therapy in patients with metastatic adenocarcinoma and common *EGFR* mutations.⁸⁸⁰ The median PFS was 11.0 months (95% CI, 10.6–12.9) with afatinib versus 10.9 months (95% CI, 9.1–11.5) with gefitinib (HR, 0.73; 95% CI, 0.57–0.95; $P = .017$). These slight PFS differences are not clinically relevant. Updated results indicate that overall survival was not significantly different between afatinib and gefitinib (27.9 vs. 24.5 months; HR, 0.86; 95% CI, 0.66–1.12; $P = .2580$).⁸⁸¹ Patients receiving afatinib had more serious treatment-related side effects when compared with those receiving gefitinib (11% [17/160] for afatinib vs. 4% [7/159] for gefitinib). One patient receiving gefitinib died from treatment-related hepatic and renal failure; other deaths were not considered to be related to treatment (9% vs. 6% [15/160 vs. 10/159]). More patients receiving afatinib had diarrhea (13% vs. 1%), whereas more patients receiving gefitinib had elevations in liver enzyme levels (0% vs. 9%). The NCCN Guidelines do not state that afatinib is more efficacious than gefitinib (see the NCCN Guidelines with Evidence Blocks™ for NSCLC, available at www.NCCN.org).⁸⁸² Afatinib is rated as slightly less safe than erlotinib or gefitinib (ie, a rating of 3 for afatinib vs. 4 for erlotinib and gefitinib).

Erlotinib and Gefitinib

Erlotinib and gefitinib are first-generation oral TKIs that inhibit *EGFR* exon 19 deletions and *EGFR* exon 21 L858R mutations (both are common mutations) as well as *EGFR* G719X, L861Q, and/or S768I (less common mutations) (see *Oral TKIs that Inhibit EGFR G719X, L861Q, and S768I Alterations* in this Discussion).

Clinical Trial Data

IPASS, a phase 3 randomized trial, assessed first-line therapy with gefitinib alone versus carboplatin plus paclitaxel in Asian patients with *EGFR*-positive metastatic NSCLC.²⁵⁸ Patients with *EGFR* mutations who received gefitinib had longer PFS (24.9% vs. 6.7%), increased response



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

rate (71.2% vs. 47.3%), and improved quality of life with fewer side effects (eg, neutropenia) compared with carboplatin plus paclitaxel.²⁵⁸ Updated results from the IPASS trial showed that overall survival was similar in patients receiving gefitinib or chemotherapy regardless of *EGFR* mutation status.⁸⁸³ These results probably occurred because patients who had been assigned to first-line chemotherapy were able to receive TKIs as subsequent therapy if they were found to have *EGFR* mutations.

EURTAC, a phase 3 randomized trial, assessed first-line therapy with erlotinib versus chemotherapy in European patients with metastatic NSCLC and *EGFR* mutations.²⁵² For erlotinib, the median PFS was 9.7 months (95% CI, 8.4–12.3) compared with 5.2 months (95% CI, 4.5–5.8) for chemotherapy (HR, 0.37; 95% CI, 0.25–0.54; $P < .0001$). Fewer patients receiving erlotinib had severe adverse events or died when compared with those receiving chemotherapy. Previously, erlotinib was often used in the United States in patients with common *EGFR* mutations because of restrictions on the use of gefitinib. However, gefitinib was re-approved by the FDA based on a phase 4 study and is available in the United States.^{183,884}

CALGB 30406, a phase 3 randomized trial, compared first-line erlotinib monotherapy versus erlotinib plus carboplatin plus paclitaxel in patients (mainly Caucasian) with advanced NSCLC and *EGFR* mutations.⁸⁸⁵ Erlotinib monotherapy was associated with fewer side effects in patients with *EGFR* mutations compared with erlotinib plus chemotherapy. Thus, it is appropriate to interrupt or complete planned chemotherapy and switch to *EGFR* TKI therapy in patients found to have *EGFR* mutations during first-line chemotherapy.⁸⁸⁶ The NCCN Guidelines do not recommend adding *EGFR* TKIs to current chemotherapy based on this CALGB study.⁸⁸⁵ *EGFR* TKIs may be continued in patients with disease progression if they do not have multiple systemic symptomatic lesions

(see *Continuation of Targeted Therapy After Progression on Initial Therapy* in this Discussion).

WJOG 5108L, a phase 3 randomized trial, assessed gefitinib versus erlotinib for patients with advanced lung cancer who had been previously treated with chemotherapy; most patients (72%) were positive for *EGFR* mutations.⁸⁸² The median PFS was 8.3 months for gefitinib versus 10.0 months for erlotinib in patients positive for *EGFR* mutations (HR, 1.093; 95% CI, 0.879–1.358; $P = .424$). The main grade 3 or 4 toxicities included rash (gefitinib: 2.2% vs. erlotinib: 18.1%) and increases in ALT/AST levels (gefitinib: 6.1%/13.0% vs. erlotinib: 2.2%/3.3%).

An analysis of 5 clinical trials in patients ($n = 223$) with NSCLC (stage IIIB or IV), mainly from the Western hemisphere, found that those with *EGFR* mutations who received TKIs had a 67% response rate and an overall survival of about 24 months.⁸⁸⁷ The TORCH trial suggested that *EGFR* mutation testing should be done in patients with advanced nonsquamous NSCLC.⁸⁸⁸ Survival was longer in patients with wild-type *EGFR* who received first-line chemotherapy compared with those who received erlotinib first followed by subsequent chemotherapy (11.6 vs. 8.7 months). The OPTIMAL trial reported that PFS was increased in patients with *EGFR* mutations who received erlotinib.^{255,256} *EGFR* TKIs are recommended in patients with metastatic NSCLC and *EGFR* mutations, because quality of life is improved when compared with chemotherapy. Erlotinib and gefitinib are orally active TKIs that are very well tolerated by most patients.^{889,890}

RELAY, a phase 3 randomized trial, compared first-line therapy with erlotinib plus ramucirumab versus erlotinib alone in patients with advanced NSCLC and *EGFR* mutations.⁸⁹¹ PFS was 19.4 months (95% CI, 15.4–21.6) with erlotinib plus ramucirumab versus 12.4 months (95% CI, 11.0–13.5) with erlotinib alone (HR, 0.59; 95% CI, 0.46–0.76; $P < .0001$). The overall response rate was similar (erlotinib plus ramucirumab: 76% vs.



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

erlotinib alone: 75%). Serious adverse events (grade 3–4) occurred in 72% (159/221) of patients receiving erlotinib plus ramucirumab (including hypertension) versus 54% (121/225) in those receiving erlotinib alone (including increased ALT). One treatment-related death occurred in a patient receiving erlotinib plus ramucirumab.

NEJ026, a phase 3 randomized trial, compared first-line erlotinib plus bevacizumab versus erlotinib alone in 228 patients with *EGFR*-positive advanced nonsquamous NSCLC.^{892,893} At interim analysis, PFS was 16.9 months (95% CI, 14.2–21.0) for erlotinib plus bevacizumab versus 13.3 months (95% CI, 11.1–15.3) for erlotinib alone (HR, 0.605; 95% CI, 0.417–0.877; $P = .016$).⁸⁹³ Grade 3 or worse events occurred in 88% (98/112) of patients receiving erlotinib plus bevacizumab versus 46% (53/114) of those receiving erlotinib alone. Grade 4 adverse events occurred in 8% (9/112) of patients receiving erlotinib plus bevacizumab (including neutropenia, hepatic dysfunction) versus 4% (5/114) of patients receiving erlotinib alone (hepatic dysfunction); no treatment-related deaths were reported.⁸⁹³ Updated data showed that median overall survival was 50.7 months (95% CI, 37.3–not estimable) in those receiving erlotinib plus bevacizumab versus 46.2 months (38.2–not estimable) in those receiving erlotinib alone (HR, 1.007; 95% CI, 0.681–1.490; $P = .97$).⁸⁹²

NCCN Recommendations

The NCCN NSCLC Panel recommends erlotinib and gefitinib as first-line therapy options in patients with metastatic nonsquamous NSCLC and *EGFR* exon 19 deletions or *EGFR* exon 21 L858R mutations (regardless of their PS) based on these trials and FDA approvals.^{121,258,894,895} Erlotinib and gefitinib are category 1 (other recommended) options if an *EGFR* exon 19 deletion or *EGFR* exon 21 L858R mutation is discovered before giving first-line systemic therapy; erlotinib and gefitinib are also treatment options if an *EGFR* exon 19 deletion or *EGFR* exon 21 L858R mutation is discovered during first-line systemic therapy (eg, carboplatin plus paclitaxel).

The NCCN NSCLC Panel recommends erlotinib plus ramucirumab as a first-line therapy option for patients with metastatic NSCLC and *EGFR* exon 19 deletions or *EGFR* exon 21 L858R mutations (other recommended intervention) based on clinical data.⁸⁹¹ The panel also recommends erlotinib plus bevacizumab as a first-line therapy option for eligible patients with metastatic nonsquamous NSCLC who have common *EGFR* mutations and no contraindications to bevacizumab (other recommended) based on clinical data.^{892,893} The NCCN NSCLC Panel recommends that bevacizumab biosimilars may be used in any of the regimens containing bevacizumab used for eligible patients with metastatic NSCLC.⁸⁹⁶⁻⁹⁰⁰ Erlotinib (± bevacizumab or ramucirumab) or gefitinib may also be continued in patients with disease progression if they do not have multiple systemic symptomatic lesions [see *Second-Line and Beyond (Subsequent) Systemic Therapy* in this Discussion].²⁴⁶ However, erlotinib is not recommended as subsequent therapy in patients with metastatic squamous cell NSCLC but without *EGFR* mutations based on a phase 3 randomized trial showing low response rates (3%); it is less efficacious and safe compared to other available options.⁸⁷⁸

The NCCN NSCLC Panel has preference stratified the systemic therapy regimens and decided that erlotinib (± bevacizumab or ramucirumab) and gefitinib are “other recommended” first-line options for patients with metastatic NSCLC and *EGFR* exon 19 deletions or *EGFR* exon 21 L858R mutations; osimertinib is the preferred option in this setting. Erlotinib and gefitinib are also recommended for eligible patients with metastatic NSCLC and less common *EGFR* mutations, such as *EGFR* S768I, L861Q, and/or G719X mutations (see *Oral TKIs that Inhibit EGFR S768I, L861Q, and G719X Alterations* in this Discussion). Flare phenomenon may occur in some patients who discontinue *EGFR* TKIs. If disease flare occurs, then the *EGFR* TKIs should be restarted.

Dacomitinib



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

Dacomitinib is a second-generation oral TKI that irreversibly inhibits ErbB/HER receptors including EGFR, HER1, HER2, and HER4.

Clinical Trial Data

ARCHER 1050, a phase 3 randomized trial, compared dacomitinib versus gefitinib as first-line therapy for patients with *EGFR*-positive metastatic NSCLC.^{901,902} Patients with brain metastases were not eligible for enrollment. PFS was increased in patients receiving dacomitinib (14.7 months; 95% CI, 11.1–16.6) compared with those receiving gefitinib (9.2 months; 95% CI, 9.1–11.0). Serious adverse events related to treatment were reported in 21 (9%) patients given dacomitinib and in 10 (4%) patients given gefitinib. Treatment-related deaths included two patients in the dacomitinib group (one related to untreated diarrhea and one to untreated cholelithiasis/liver disease) and one patient in the gefitinib group (related to sigmoid colon diverticulitis/rupture complicated by pneumonia). An updated analysis reported that the median overall survival was 34.1 months (95% CI, 29.5–39.8) in patients receiving dacomitinib compared with 27.0 months (95% CI, 24.4–31.6) in those receiving gefitinib (HR, 0.748; 95% CI, 0.591–0.947; two-sided $P = .0155$).^{901,903}

NCCN Recommendations

The NCCN NSCLC Panel recommends dacomitinib as a first-line treatment option for patients with metastatic NSCLC and *EGFR* exon 19 deletions or *EGFR* exon 21 L858R mutations based on clinical trial data and FDA approval.^{901,904} Dacomitinib is a category 1 (other recommended) option if an *EGFR* exon 19 deletion or *EGFR* exon 21 L858R mutation is discovered before giving first-line systemic therapy; dacomitinib is an option if an *EGFR* exon 19 deletion or *EGFR* exon 21 L858R mutation is discovered during first-line systemic therapy (eg, carboplatin plus paclitaxel). The NCCN NSCLC Panel has preference stratified the systemic therapy regimens and decided that dacomitinib is an “other recommended” option; osimertinib is the preferred option in this setting. Dacomitinib is also recommended for eligible patients with metastatic

NSCLC and less common *EGFR* mutations, such as *EGFR* S768I, L861Q, and/or G719X mutations (see *Oral TKIs that Inhibit EGFR S768I, L861Q, and G719X Alterations* in this Discussion).

Osimertinib

Osimertinib is a third-generation oral TKI that inhibits the common and uncommon *EGFR* mutations and T790M. Common *EGFR* mutations include exon 19 deletion and exon 21 L858R; less common *EGFR* mutations include S768I, L861Q, and/or G719X (see *EGFR Mutations* in this Discussion). Patients with these mutations are sensitive to small-molecule oral EGFR TKIs, such as afatinib, dacomitinib, erlotinib, gefitinib, and osimertinib.²⁴⁶ *EGFR* T790M is a mutation associated with acquired resistance to first-line therapy with EGFR TKIs and has been reported in about 60% of patients with disease progression after initial response to first-line EGFR TKIs, such as erlotinib or gefitinib.^{212,267-273} Most patients with *EGFR* mutations and metastatic NSCLC typically have disease progression after about 9.7 to 13 months of therapy with erlotinib, gefitinib, or afatinib.^{252,258,266,267} Data show that patients receiving osimertinib as first-line therapy have PFS of about 19 months.^{373,905}

First-Line Therapy

Clinical Trial Data

FLAURA, a phase 3 randomized trial, assessed first-line therapy with osimertinib compared with either erlotinib or gefitinib in patients with metastatic NSCLC and *EGFR* mutations regardless of T790M status.^{21,373,905,906} PFS was longer with osimertinib (18.9 months; 95% CI, 15.2–21.4) compared with either erlotinib or gefitinib (10.2 months; 95% CI, 9.6–11.1; HR, 0.46; 95% CI, 0.37–0.57; $P < .001$). The median duration of response was longer with osimertinib compared with erlotinib or gefitinib (median response, 17.2 vs. 8.5 months). Only 6% (17/279) of patients receiving osimertinib had CNS progression events when compared with 15% (42/277) of those receiving erlotinib or gefitinib. Grade



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

3 or higher adverse events were reported in 34% (94/279) of patients receiving osimertinib and 45% (124/277) of patients receiving erlotinib or gefitinib. An updated analysis showed that median overall survival was 38.6 months with osimertinib (95% CI, 34.5–41.8) compared with 31.8 months (95% CI, 26.6–36.0) for either erlotinib or gefitinib (HR, 0.8; 95% CI, 0.64–1.0; $P = .046$).²¹

NCCN Recommendations

The NCCN NSCLC Panel recommends osimertinib as a preferred first-line therapy option for patients with metastatic NSCLC and *EGFR* exon 19 deletions or *EGFR* exon 21 L858R mutations based on clinical data and FDA approval.^{21,373} Osimertinib is a category 1 (preferred) recommended option if an *EGFR* exon 19 deletion or *EGFR* exon 21 L858R mutation is discovered before giving first-line systemic therapy, and osimertinib is a preferred option if an *EGFR* mutation is discovered during first-line systemic therapy (eg, carboplatin plus paclitaxel).²¹ For patients receiving first-line ICIs with or without chemotherapy, oncologists should be aware of the long half-life of the ICIs and potential adverse effects when using osimertinib in combination with or following ICIs.^{907,908} If patients require an urgent start to first-line therapy, but molecular testing results are pending, then clinicians should consider holding immunotherapy regimens and only using chemotherapy regimens for the first cycle. Flare phenomenon may occur in some patients who discontinue *EGFR* TKIs. If disease flare occurs, then the *EGFR* TKIs should be restarted.⁸³⁶⁻⁸³⁹

Osimertinib is also recommended as an adjuvant therapy option for eligible patients with completely resected stage IB to IIIA or stage IIIB (only T3,N2) NSCLC and *EGFR* exon 19 deletions or *EGFR* exon 21 L858R mutations who have previously received adjuvant chemotherapy or are ineligible to receive platinum-based chemotherapy (see *Surgery Followed by Adjuvant Therapy: Trial Data and NCCN Recommendations* in this Discussion).⁷⁴⁹ In addition, osimertinib is recommended for eligible

patients with metastatic NSCLC and less common *EGFR* mutations, such as *EGFR* S768I, L861Q, and/or G719X mutations (see *Oral TKIs that Inhibit EGFR S768I, L861Q, and G719X Alterations* in this Discussion).²⁹³

Subsequent Therapy

Clinical Trial Data

AURA3, a phase 3 randomized trial, assessed osimertinib versus platinum-pemetrexed chemotherapy in patients with *EGFR* T790M-positive metastatic NSCLC and disease progression on first-line erlotinib, gefitinib, or afatinib. PFS was longer with osimertinib compared with chemotherapy (10.1 vs. 4.4 months; HR, 0.30; 95% CI, 0.23–0.41; $P < .001$).²⁶⁶ PFS was also longer in patients with CNS metastases who received osimertinib versus chemotherapy (8.5 vs. 4.2 months; HR, 0.32; 95% CI, 0.21–0.49). In addition, the objective response rate was increased with osimertinib (71%; 95% CI, 65%–76%) compared with chemotherapy (31%; 95% CI, 24%–40%) (odds ratio for objective response, 5.39; 95% CI, 3.47–8.48; $P < .001$). The disease control rate was about 93% with osimertinib (95% CI, 90%–96%) and about 74% with chemotherapy (95% CI, 66%–81%). Patients receiving osimertinib had fewer grade 3 or higher adverse events compared with those receiving chemotherapy (23% vs. 47% [63/279 vs. 64/136]). There were four fatal events with osimertinib (respiratory failure [2 patients], pneumonitis, and ischemic stroke) and one with chemotherapy (hypovolemic shock).

The BLOOM study ($n = 41$) assessed osimertinib (160 mg once daily) in patients with *EGFR* mutations (regardless of T790M status) who had progressive leptomeningeal disease.^{909,910} Most patients were women (71%) and all were Asian. Most patients had previously received gefitinib (76%). The median overall survival was 11 months (95% CI, 8.0–18).⁹⁰⁹ The overall response rate was 41% (95% CI, 26%–58%). TC clearance in the cerebrospinal fluid was confirmed in 28% of patients (11/40; 95% CI, 15%–44%). Neurologic function improved in 57% (12/21) of patients with



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

abnormal baseline assessments. Grade 3 or higher adverse events were reported in 66% (27/41) of patients. There were seven deaths due to adverse events, but they were not considered to be related to osimertinib.

NCCN Recommendations

The NCCN NSCLC Panel recommends broad molecular profiling for T790M (category 1) and other genomic resistance mutations for eligible patients with *EGFR* mutation-positive NSCLC and disease progression on certain *EGFR* TKIs based on the efficacy of osimertinib. The NCCN NSCLC Panel recommends osimertinib (category 1) as a subsequent therapy option for patients with metastatic *EGFR* T790M–positive NSCLC and disease progression on certain *EGFR* TKIs (including erlotinib [± ramucirumab or bevacizumab]) based on clinical trial data and FDA approval [see *Second-Line and Beyond (Subsequent) Systemic Therapy* in this Discussion].²⁶⁶

For patients with *EGFR* mutations who have disease progression during or after first-line therapy with osimertinib, recommended subsequent therapy depends on whether the progression is asymptomatic or symptomatic and includes: 1) considering local therapy (eg, SABR or surgery); 2) continuing osimertinib; or 3) a first-line systemic therapy regimen for metastatic NSCLC (such as carboplatin plus paclitaxel). There are no data to support using erlotinib (with or without ramucirumab or bevacizumab), gefitinib, afatinib, or dacomitinib after progression on first-line therapy with osimertinib. T790M can be assessed using an FDA-approved test or another validated laboratory test done in a CLIA-approved laboratory. Data suggest that plasma genotyping (also known as plasma ctDNA testing or liquid biopsy) may be considered at progression instead of tissue biopsy to detect whether patients have T790M; however, if plasma ctDNA testing is negative, then tissue biopsy is recommended.⁹¹¹⁻⁹¹³

The NCCN NSCLC Panel also recommends osimertinib (category 1) for patients with T790M who have symptomatic brain metastases after

progression on erlotinib (with or without ramucirumab or bevacizumab), gefitinib, afatinib, or dacomitinib based on data showing an improvement.^{266,904,914-916} The NCCN Panel recommends that osimertinib (regardless of T790M status) can be considered for patients with *EGFR* mutations who have progressive CNS disease or leptomeningeal disease based on clinical trial data.^{909,910}

Oral TKIs that Inhibit *EGFR* S768I, L861Q, and G719X Alterations

EGFR exon 18 G719X, exon 21 L861Q, and exon 20 S768I are less common *EGFR* mutations (10%) that are also sensitive to first-, second-, and third-generation *EGFR* TKIs, such as afatinib, dacomitinib, erlotinib, gefitinib, and osimertinib.^{248,249,293-295,879} *EGFR* exon 18 G719X mutations occur more frequently than exon 21 L861Q; exon 20 S768I mutations are the least frequent. The NCCN NSCLC Panel recommends testing for *EGFR* S768I, L861Q, and G719X mutations in all patients with metastatic nonsquamous NSCLC or NSCLC NOS based on data showing the efficacy of *EGFR* TKIs as first-line therapy options for patients with these mutations.^{293,295,879} *EGFR* S768I, L861Q, and G719X mutation testing can be considered in patients with metastatic squamous cell carcinoma.

Clinical Trial Data

KCSG-LU15-09, a phase 2 trial, assessed first-line therapy with osimertinib in 37 patients with metastatic NSCLC and less common *EGFR* mutations, including S768I, L861Q, and G719X.²⁹³ The median PFS was 8.2 months (95% CI, 5.9–10.5 months). The objective response rate was 50% (18/36; 95% CI, 33%–67%). Manageable adverse events included rash, pruritis, decreased appetite, diarrhea, and dyspnea.²⁹³

A post-hoc analysis of several LUX-Lung trials (LUX-Lung 2, 3, and 6) assessed afatinib in a few patients with *EGFR* L861Q, G719X, and S768I mutation-positive metastatic NSCLC.⁸⁷⁹ Median overall survival was 19.4 months (95% CI, 16.4–26.9). Of those with *EGFR* S768I mutations,



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

100% (95% CI, 63.1%–100%) responded to afatinib. Of patients with *EGFR* G719X mutations, 77.8% (95% CI, 52.4%–93.6%) had an objective response to afatinib. Of those with *EGFR* L861Q, 56% (95% CI, 29.9%–80.2%) responded to afatinib.

NCCN Recommendations

The NCCN NSCLC Panel recommends testing for *EGFR* S768I, L861Q, and G719X mutations in eligible patients with metastatic NSCLC based on data showing the efficacy of EGFR TKIs as first-line therapy options for patients with these mutations.^{293,295,879} The panel recommends afatinib or osimertinib as first-line therapy options (preferred) for patients with metastatic NSCLC who have these less common *EGFR* mutations based on clinical trial data.^{293,879} The NCCN NSCLC Panel has preference stratified the systemic therapy regimens and decided that afatinib or osimertinib are preferred options for patients with metastatic NSCLC and *EGFR* L861Q, G719X, and S768I mutations; other recommended options include erlotinib, gefitinib, or dacomitinib.

Agents that Inhibit *EGFR* Exon 20 Insertion Mutations

EGFR exon 20 mutations are a heterogeneous group; most variants do not respond to first-, second- or third-generation EGFR TKIs. However, some *EGFR* exon 20 alterations, such as p.A763_Y764insFQEA, are sensitive to EGFR TKIs; p.A763_Y764insLQEA may be sensitive to first and third-generation EGFR TKIs. First-line platinum-based chemotherapy is typically recommended for most patients with *EGFR* exon 20 insertion-positive metastatic NSCLC (eg, carboplatin plus paclitaxel).^{299,301} The response rates (0%–25%) to immunotherapy regimens vary, depending on the specific *EGFR* exon 20 insertion mutation.^{181,304} Response rates to classic subsequent therapy regimens, such as docetaxel, are low (14%) in patients with metastatic NSCLC and disease progression after first-line therapy.

The NCCN NSCLC Panel recommends testing for *EGFR* exon 20 insertion mutations in all patients with metastatic nonsquamous NSCLC or NSCLC NOS based on data showing the efficacy of several agents as subsequent therapy options for patients with *EGFR* exon 20 insertion-positive metastatic NSCLC.^{180,181} *EGFR* exon 20 insertion mutation testing can be considered in patients with metastatic squamous cell carcinoma.

Amivantamab

Amivantamab-vmjw is a bispecific human antibody to EGFR and MET receptors that bypasses resistance to EGFR TKIs and has immune-cell directing activity.¹⁸⁰

Subsequent Therapy

CHRYSLIS, a phase 1 study, assessed subsequent therapy with amivantamab-vmjw in 81 patients with *EGFR* exon 20 insertion-positive metastatic NSCLC who had received one or more previous lines of therapy.¹⁸⁰ The reported overall response rate was 40% (95% CI, 29%–51%) with 3 complete responses. The median PFS was 8.3 months (95% CI, 6.5–10.9). Common treatment-related adverse events included cutaneous reactions, infusion-related reactions, and paronychia. The most common grade 3 to 4 adverse events included hypokalemia (5% [6/114]) as well as pulmonary embolism, neutropenia, diarrhea, and rash (4% for each [4/114]). Eight deaths were reported in the safety assessment (7% [8/114]).

The NCCN NSCLC Panel recommends amivantamab as a subsequent therapy option for patients with *EGFR* exon 20 insertion mutation-positive metastatic NSCLC and disease progression on or after systemic therapy options based on clinical trial data and FDA approval.¹⁸⁰ If patients have disease progression on amivantamab, then the panel recommends either mobocertinib or subsequent systemic therapy options (eg, albumin-bound paclitaxel). If the patient has previously received amivantamab, then they



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

can receive mobocertinib as third-line or beyond therapy and vice versa because these agents have different mechanisms of action. For the 2023 update (Version 1), the NCCN Panel added a caveat that amivantamab or mobocertinib can be used as third-line or beyond therapy if the patient has previously received only one, or none, of these agents.

Mobocertinib

Mobocertinib is an oral TKI that selectively inhibits diverse *EGFR* and *ERBB2* (*HER2*) exon 20 insertion mutations.^{181,297}

Subsequent Therapy

Phase 1/2 trials (n =114) assessed subsequent therapy with mobocertinib for patients with *EGFR* exon 20 insertion mutation-positive metastatic NSCLC who had received first-line platinum-based chemotherapy.^{181,297} The objective response rate was 28% (95% CI, 20%–37%). Median overall survival was 24 months (95% CI, 14.6–28.8). Patients with brain metastases had a lower overall response rate (18%; 95% CI, 7%–33%) to mobocertinib compared to those without brain metastases (34%; 95% CI, 23%–46%). Diarrhea and rash were the most common treatment-related adverse events. Grade 3 or 4 treatment-related adverse events were reported in 47% of patients (54/114); diarrhea was the most common grade 3 or 4 adverse event (21% [24/114]).²⁹⁷ One death (cardiac failure) was reported to be related to treatment with mobocertinib.

The NCCN NSCLC Panel recommends mobocertinib as a subsequent therapy option for patients with *EGFR* exon 20 insertion-positive metastatic NSCLC and disease progression on or after initial systemic therapy options based on clinical trial data and FDA approval.¹⁸¹ If patients have disease progression on mobocertinib, then the panel recommends either amivantamab or subsequent systemic therapy options (eg, albumin-bound paclitaxel). If the patient has previously received mobocertinib, then they can receive amivantamab as third-line or beyond therapy and vice versa because these agents have different mechanisms

of action. For the 2023 update (Version 1), the NCCN Panel added a caveat that mobocertinib or amivantamab can be used as third-line or beyond therapy if the patient has previously received only one, or none, of these agents.

Monoclonal Antibody that Inhibits *EGFR*

Cetuximab

Cetuximab is a monoclonal antibody that targets *EGFR*. Data suggest that an afatinib plus cetuximab regimen may be useful for patients who have progressed after receiving erlotinib, afatinib, or gefitinib and after chemotherapy.⁹¹⁷ Patients with T790M-positive and T790M-negative tumors had a similar response rate to an afatinib plus cetuximab regimen (32% vs. 25%; $P = .341$). The NCCN NSCLC Panel recommends considering afatinib plus cetuximab as an option for patients with disease progression after receiving afatinib, dacomitinib, erlotinib (\pm bevacizumab or ramucirumab), or gefitinib and after chemotherapy based on these data.

FLEX, a large phase 3 randomized trial, assessed cisplatin plus vinorelbine with (or without) cetuximab for patients with advanced NSCLC; most patients had stage IV disease.⁹¹⁸ Adding cetuximab slightly increased overall survival (11.3 vs. 10.1 months; HR for death, 0.87; 95% CI, 0.762–0.996; $P = .044$). Patients receiving cetuximab had increased grade 4 events versus control (62% vs. 52%, $P < .01$); cetuximab was also associated with grade 2 acne-like rash.

The NCCN NSCLC Panel does not recommend the cetuximab plus cisplatin plus vinorelbine regimen based on the clinical data.⁹¹⁸ The benefits of this cetuximab-based regimen are very slight, it is a difficult regimen to administer, and patients have poorer tolerance for this regimen when compared with other regimens; for example, almost 40% of patients



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

have grade 4 neutropenia.⁷¹⁴ Patients may also have comorbid conditions that prevent them from receiving cisplatin such as poor kidney function. Cisplatin plus vinorelbine with (or without) cetuximab is generally not used in the United States because of concerns about toxicity.^{714,732,918} Although the FLEX trial results were reported to be statistically significant, panel members feel they were not clinically significant.⁷¹⁴ The NCCN NSCLC Panel does not recommend the cisplatin plus vinorelbine or the carboplatin plus vinorelbine regimens for patients with metastatic NSCLC.

Monoclonal Antibody-Drug Conjugates that Inhibit ERBB2 (HER2) Mutations

Ado-Trastuzumab Emtansine

Ado-trastuzumab emtansine (also known as T-DM1) is a humanized antibody-drug conjugate consisting of trastuzumab, which is an antibody (immunoglobulin G1) targeting HER2, and emtansine, which is a microtubule inhibitor.¹⁷⁰ A phase 2 basket trial assessed ado-trastuzumab emtansine in patients with metastatic NSCLC and *ERBB2* (*HER2*) mutations.^{170,919} The partial response rate was 44% (95% CI, 22%–69%). The median PFS was 5 months (95% CI, 3–9). Minor toxicities (grade 1–2) included infusion reactions, thrombocytopenia, and transaminitis; no treatment-related deaths were reported. Patients (n = 18) were mostly women (72%), did not smoke cigarettes, and all had adenocarcinoma histology. Another study assessed ado-trastuzumab emtansine in 22 patients with metastatic NSCLC and *ERBB2* (*HER2*) exon 20 mutations.⁹²⁰ The overall response rate with ado-trastuzumab emtansine was 38% (95% CI, 23%–56%). The median overall survival was 8.1 months.

Fam-Trastuzumab Deruxtecan-nxki

Fam-trastuzumab deruxtecan-nxki is a humanized monoclonal antibody-drug conjugate consisting of trastuzumab linked to deruxtecan, which is a topoisomerase I inhibitor; the agent remains stable until it is cleaved by peptides in cancer cells.¹⁷¹ DESTINY-Lung01, a phase 2

study, assessed fam-trastuzumab deruxtecan-nxki in 91 patients with metastatic nonsquamous NSCLC and *ERBB2* mutations.^{171,921,922} Most patients had *ERBB2* (*HER2*) exon 20 insertion mutations (86%); 66% were women, 57% had never smoked cigarettes, 36% had CNS metastases at baseline, and all had nonsquamous NSCLC. Updated results show that the overall response rate with fam-trastuzumab deruxtecan-nxki was 55% (95% CI, 44%–65%); most patients had received prior treatment.¹⁷¹ However, responses were also noted in patients who did not have *ERBB2* (*HER2*) mutations. Median overall survival was 17.8 months (95% CI, 13.8–22.1).¹⁷¹ Grade 3 or higher adverse events occurred in 46% of patients including neutropenia (19%). Two patients died from drug-related interstitial lung disease.

NCCN Recommendations

The NCCN NSCLC Panel recommends fam-trastuzumab deruxtecan-nxki as a subsequent therapy option for patients with metastatic NSCLC and *ERBB2* (*HER2*) mutations based on clinical trial data and FDA approval.^{171,921,922} The NCCN NSCLC Panel also recommends ado-trastuzumab emtansine as a subsequent therapy option for patients with *ERBB2* (*HER2*) mutation–positive metastatic NSCLC.¹⁷⁰ The panel preference stratified the recommended regimens for patients with *ERBB2* (*HER2*) mutation–positive metastatic NSCLC and voted that fam-trastuzumab deruxtecan-nxki is a preferred subsequent therapy option; ado-trastuzumab emtansine is an other recommended subsequent therapy option. The NCCN NSCLC Panel does not recommend single-agent therapy with trastuzumab or afatinib for patients with *ERBB2* (*HER2*) mutations, because response rates are lower and treatment is less effective with these agents.^{923,924}

The NCCN NSCLC Panel recommends testing for *ERBB2* (*HER2*) mutations in all patients with metastatic nonsquamous NSCLC or NSCLC NOS based on clinical trial data and FDA approval of fam-trastuzumab



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

deruxtecan-nxki. Testing for these biomarkers can be considered in patients with metastatic squamous cell carcinoma. Previously, *ERBB2* (*HER2*) mutations were listed as emerging biomarkers, but they are now included in the standard list because of the FDA approval. Resources are available to assess whether the *ERBB2* (*HER2*) mutations are oncogenic or likely to be oncogenic (see oncoKB.org). The panel recommends first-line therapy with platinum-based chemotherapy with or without immunotherapy for patients with metastatic NSCLC and *ERBB2* (*HER2*) mutations. Data suggest that patients with *ERBB2* mutations respond to first-line immunotherapy regimens.³⁰⁸ The response to immunotherapy ranges from 7% to 27% in patients with *ERBB2* (*HER2*) exon 20 insertion mutations.^{186,925}

Oral Agents that Inhibit KRAS Mutations

KRAS is a G-protein with GTPase activity that is part of the MAP/ERK pathway; point mutations in KRAS most commonly occur at codon 12. Data suggest that approximately 25% of patients with adenocarcinomas in a North American population have *KRAS* mutations; *KRAS* is the most common mutation in this population.^{121,177,213,309,310} *KRAS* mutation prevalence is associated with cigarette smoking, unlike many of the other actionable mutations (eg, *EGFR* mutations, *ALK* rearrangements).³¹¹ The NCCN NSCLC Panel recommends testing for *KRAS* mutations in all patients with metastatic nonsquamous NSCLC or NSCLC NOS based on data showing the efficacy of adagrasib and sotorasib as subsequent therapy options for patients with *KRAS* p.G12C mutation-positive metastatic NSCLC.³¹⁷⁻³¹⁹ *KRAS* mutation testing can be considered in patients with metastatic squamous cell carcinoma.

Adagrasib

Adagrasib is an oral RAS GTPase family inhibitor that inhibits *KRAS* p.G12C mutations in patients with metastatic NSCLC who have been previously treated with combination chemotherapy regimens (±

immunotherapy). Adagrasib has a long half-life of 23 hours. A phase 2 study assessed adagrasib as subsequent therapy in 116 patients with *KRAS* p.G12C mutation-positive advanced NSCLC who had previously received platinum-based chemotherapy (± immunotherapy).³¹⁷ Most patients had metastatic NSCLC (89%); a few had locally advanced NSCLC (11%). Adenocarcinoma was the most common histology (97%). Most patients had previously smoked (86%), although a few currently smoked (9.5%). The median overall survival was 12.6 months (95% CI, 9.2–19.2). The objective response rate was 42.9% (95% CI, 33.5%–52.6%). The intracranial response rate was 33.3% (95% CI, 18.0%–51.8%). Grade 3 or higher adverse events occurred in 44.8% of patients (52/116), including anemia, pneumonia, dyspnea, acute kidney injury, hyponatremia, fatigue, nausea, and increased ALT and AST levels. There were two deaths (one from cardiac failure, one from pulmonary hemorrhage).

For the 2023 update (Version 1), the NCCN NSCLC Panel recommends adagrasib as a subsequent therapy option for select patients with metastatic NSCLC and *KRAS* p.G12C mutations who have disease progression after treatment with platinum-based chemotherapy (± immunotherapy) based on clinical trial data and FDA approval.³¹⁷ However, responsiveness to adagrasib has not been assessed for mutations other than *KRAS* G12C. The NCCN Panel clarified that adagrasib may be used as third-line or beyond therapy if the patient has not previously received *KRAS* p.G12C targeted therapy. Because adagrasib and sotorasib have similar mechanisms of action, switching to sotorasib is not recommended after progression on adagrasib.

Sotorasib

Sotorasib is an oral RAS GTPase family inhibitor that inhibits *KRAS* p.G12C mutations in patients with metastatic NSCLC who have been previously treated with combination chemotherapy regimens (±



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

immunotherapy). Sotorasib has a half-life of 5 hours. CodeBreak 200, a randomized phase 3 trial, assessed subsequent therapy with sotorasib versus docetaxel in 345 patients with *KRAS* p.G12C mutation-positive advanced NSCLC who had previously received platinum-based chemotherapy plus immunotherapy.³¹⁸ Median PFS was 5.6 months (95% CI, 4.3–7.8) in patients receiving sotorasib compared with 4.5 months (95% CI, 3.0–5.7) in those receiving docetaxel (HR, 0.66; 95% CI, 0.51–0.86; *P* = .0017). A phase 2 study assessed sotorasib as subsequent therapy in 126 patients with *KRAS* p.G12C mutation-positive advanced NSCLC who had previously received platinum-based chemotherapy (± immunotherapy).³¹⁹ Adenocarcinoma was the most common histology (95%). Most patients had previously smoked (81%), although a few currently smoked (12%). The median overall survival was 12.5 months (95% CI, 10.0–could not be evaluated). The response rate was 37.1% (95% CI, 28.6%–46.2%). Treatment-related grade 3 adverse events occurred in 19.8% of patients (25/126), including diarrhea, and increased ALT and AST levels; one grade 4 event occurred (dyspnea and pneumonitis).

The NCCN NSCLC Panel recommends sotorasib as a subsequent therapy option for select patients with metastatic NSCLC and *KRAS* p.G12C mutations who have disease progression after treatment with platinum-based chemotherapy (± immunotherapy) based on clinical trial data and FDA approval.³¹⁹ However, responsiveness to sotorasib has not been assessed for mutations other than *KRAS* G12C. For the 2023 update (Version 1), the NCCN Panel clarified that sotorasib is recommended as third-line or beyond therapy if the patient has not previously received targeted therapy for *KRAS* p.G12C mutations. Because sotorasib and adagrasib have similar mechanisms of action, switching to adagrasib is not recommended after progression on sotorasib.

Oral TKIs that Inhibit *MET*ex14 Skipping Mutations

Oncogenic driver genomic alterations in *MET* include *MET*ex14 skipping mutations, *MET* GCN gain or amplification, and *MET* protein overexpression (see *MET* Genomic Alterations in this Discussion). The NCCN Panel recommends testing for *MET*ex14 skipping mutations in eligible patients with metastatic NSCLC based on clinical trial data and FDA approvals for several agents.^{172,327,835} Although high-level *MET* amplification is an emerging biomarker in the NCCN Guidelines, the panel feels that optimal biomarker testing will include this biomarker based on clinical trial data.^{167–169,172} Note that the definition of high-level *MET* amplification is evolving and may differ depending on which assay is used. When using NGS, a copy number greater than 10 is consistent with high-level *MET* amplification.

Capmatinib

Capmatinib is an oral TKI that selectively inhibits *MET* genomic alterations. Capmatinib has been assessed in phase 1 and 2 studies of patients with advanced NSCLC.^{172,326,926,927} GEOMETRY, a phase 2 study, assessed capmatinib in different cohorts of patients with *MET* genomic alterations, including those with *MET*ex14 skipping mutations; patients had stage IIIB or IV NSCLC and were wild-type for *EGFR* and *ALK* genomic alterations.^{172,326} Updated results from GEOMETRY show that first-line therapy with capmatinib yielded an overall response rate of 68% (95% CI, 48%–84%) in 28 patients with *MET*ex14 skipping mutations; the median PFS was 9.13 months (5.52–13.9 months) for first-line therapy.¹⁷² Subsequent therapy with capmatinib yielded an overall response rate of 41% (95% CI, 29%–53%) in 69 patients with *MET*ex14 skipping mutations; the median PFS was 5.42 months (95% CI, 4.17–6.97 months) for subsequent therapy.¹⁷² Updated results from GEOMETRY suggest that capmatinib is effective for patients with brain metastases.^{172,926} Of patients with brain metastases, 54% (7/13) responded to capmatinib; 4 patients had a complete response in the brain. However, 43% (3/7) of patients



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

whose disease responded had previously received RT.¹⁷² Common adverse events for patients with *MET*ex14 skipping mutations across all cohorts included peripheral edema (65%), nausea (46%), and vomiting (26%), but most of these events were grades 1 to 2.¹⁷² Grade 3 to 4 adverse events occurred in 75% of patients. One treatment-related death occurred. There were fewer adverse gastrointestinal events when capmatinib was administered without fasting.

The NCCN NSCLC Panel recommends capmatinib as either a first-line therapy or subsequent therapy option (preferred) for patients with metastatic NSCLC and *MET*ex14 skipping mutations based on clinical trial data and FDA approval.^{172,326,926} Capmatinib may be used as a subsequent therapy option if it, tepotinib, or crizotinib were not previously given as first-line therapy. The panel preference stratified the recommended regimens for patients with *MET*ex14 skipping mutation–positive metastatic NSCLC and decided that capmatinib and tepotinib are preferred first-line therapy or subsequent therapy options. The panel preference stratified crizotinib as useful in certain circumstances for either a first-line therapy or subsequent therapy option.³²⁷ Systemic therapy regimens were preference stratified as useful in certain circumstances as first-line therapy options (eg, carboplatin plus paclitaxel). These platinum doublets may be used as subsequent therapy for patients with disease progression on capmatinib, tepotinib, or crizotinib.

GEOMETRY also assessed capmatinib in patients with metastatic NSCLC and high-level *MET* amplification.¹⁷² For first-line capmatinib, response rates were 40% (95% CI, 16%–68%); for second-line capmatinib, response rates were 29% (95% CI, 19%–41%). Adverse events for patients with high-level *MET* amplification across all cohorts included peripheral edema (49%), vomiting (30%), and nausea (48%); most of these events were grade 1 or 2. Grade 3 to 4 adverse events occurred in 68% of patients. The NCCN Panel recommends capmatinib, crizotinib,

and tepotinib for patients with metastatic NSCLC and high-level *MET* amplification (GCN of ≥ 10) based on clinical trial data.^{167-169,172} As previously mentioned, high-level *MET* amplification is an emerging biomarker in the NCCN Guidelines.

Crizotinib

Crizotinib is an oral TKI that inhibits some *MET* tyrosine kinases (high-level *MET* amplification or *MET*ex14 skipping mutation), *ALK* rearrangements, and *ROS1* rearrangements; it is approved by the FDA for patients with metastatic NSCLC who have *ALK* or *ROS1* rearrangements. A phase 2 study assessed crizotinib in 69 patients with advanced NSCLC who were positive for *MET*ex14 skipping mutations.³²⁷ The objective response rate was 32% (95% CI, 21%–45%). Median PFS was 7.3 months (95% CI, 5.4–9.1 months).

The NCCN NSCLC Panel recommends crizotinib as a first-line therapy or subsequent therapy option (useful in certain circumstances) for patients with metastatic NSCLC and *MET*ex14 skipping mutations based on these data.³²⁷ Crizotinib may be used as a subsequent therapy option if it, tepotinib, or capmatinib were not previously given as first-line therapy for *MET*ex14 skipping mutation–positive metastatic NSCLC. The panel preference stratified the recommended regimens for patients with *MET*ex14 skipping mutation–positive metastatic NSCLC and decided that capmatinib and tepotinib are preferred first-line therapy or subsequent therapy options based on clinical trial data.³²⁶ The panel decided that crizotinib is useful in certain circumstances as either a first-line therapy or subsequent therapy option.³²⁷ Systemic therapy regimens are also recommended as useful in certain circumstances for first-line therapy (eg, carboplatin plus paclitaxel). These platinum doublets may be used as subsequent therapy options for patients with disease progression on capmatinib, tepotinib, or crizotinib. The NCCN Panel also recommends capmatinib, crizotinib, and tepotinib for patients with metastatic NSCLC



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

who are positive for high-level *MET* amplification (GCN of ≥ 10) based on clinical trial data.^{167-169,172}

Tepotinib

Tepotinib is an oral TKI that selectively inhibits *MET*ex14 skipping mutations and high-level *MET* amplification (see *MET Genomic Alterations* in this Discussion). VISION, a phase 2 study, assessed tepotinib in patients with *MET*ex14 skipping mutations; patients mainly had stage IV NSCLC and were wild-type (negative) for *EGFR* and *ALK* genomic alteration.^{169,835,928} The response rate to tepotinib was 46% (95% CI, 36%–57%); PFS was 8.5 months (95% CI, 6.7–11) in the combined biopsy group (tissue biopsy plus plasma ctDNA). Grade 3 or higher adverse events occurred in 28% of patients receiving tepotinib, such as peripheral edema (7%); 11% of patients had to permanently discontinue tepotinib because of peripheral edema, pleural effusion, or dyspnea. One treatment-related death occurred. Another cohort of the VISION trial assessed tepotinib in 24 patients with advanced NSCLC and *MET* amplification but without *MET*ex14 skipping mutations.¹⁶⁹ Preliminary data suggest the overall response rate is about 42% (10/24).

The NCCN NSCLC Panel recommends tepotinib as either a first-line or subsequent therapy option (preferred) for eligible patients with metastatic NSCLC who are positive for *MET*ex14 skipping mutations based on clinical trial data and FDA approval.⁸³⁵ Tepotinib may be used as a subsequent therapy option for *MET*ex14 skipping mutation–positive metastatic NSCLC if tepotinib, capmatinib, or crizotinib were not previously given as first-line therapy. The NCCN NSCLC Panel preference stratified the recommended regimens for patients with *MET*ex14 skipping mutation–positive metastatic NSCLC and decided that tepotinib and capmatinib are preferred first-line therapy or subsequent therapy options based on clinical trial data.^{172,835} The NCCN NSCLC Panel recommends tepotinib for patients with advanced NSCLC and high-level *MET* amplification based on

preliminary data; capmatinib and crizotinib are also recommended options in this setting.^{167-169,172} As previously mentioned, high-level *MET* amplification is an emerging biomarker in the NCCN Guidelines.

Oral TKIs that Inhibit NTRK1/2/3 Gene Fusions

NTRK1/2/3 gene fusions encode *TRK* fusion proteins that act as oncogenic drivers for various solid tumors, including lung, salivary gland, thyroid, and sarcoma (see *NTRK1/2/3 Gene Fusions* in this Discussion).³³⁰ The NCCN Panel recommends testing for *NTRK1/2/3* gene fusions in all patients with metastatic nonsquamous NSCLC or NSCLC NOS based on clinical trial data. *NTRK1/2/3* gene fusion testing can be considered in patients with metastatic squamous cell carcinoma.

Entrectinib

Entrectinib is an oral TKI that inhibits *TRK* fusion proteins across a range of solid tumors in younger and older patients with unresectable or metastatic disease; thus, entrectinib is an age- and tumor-agnostic therapy. Entrectinib has been assessed in several phase 1 and 2 trials in patients with *NTRK* gene fusion–positive metastatic NSCLC (phase 2 STARTRK-2 trial, phase 1 STARTRK-1 trial, and phase 1 ALKA-372-001 trial).^{333,350,929} Pooled data from these 3 trials in 10 patients with *NTRK* gene fusion–positive NSCLC showed that entrectinib yielded an overall response rate of 70% (95% CI, 35%–93%; 7/10: 7/7 adenocarcinoma NSCLC, 0/3 squamous cell carcinoma, unclassified, or undifferentiated NSCLC); there was one complete response.³³³ Most patients (70%) with *NTRK* gene fusion–positive NSCLC had received one or more lines of previous therapy. In 6 patients with CNS disease, entrectinib yielded an intracranial response rate of 67% (4/6; 2 complete responses and 2 partial responses). Grade 3 adverse events with entrectinib across a range of solid tumors included anemia and increased weight. Grade 4 adverse events occurred in 3 patients (ie, increased AST, increased ALT, blood uric acid, hyperuricemia). Nervous system disorders were the most



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

common serious treatment-related adverse event (4% [3/68] and 3% [10/355]). No treatment-related deaths were reported.

The NCCN NSCLC Panel recommends entrectinib as either a first-line or subsequent therapy option for *NTRK1/2/3* gene fusion–positive metastatic NSCLC based on these data.³³³ Entrectinib may be used as a subsequent therapy option if larotrectinib or entrectinib were not previously given as first-line therapy. The NCCN NSCLC Panel has preference stratified the recommended regimens for patients with *NTRK1/2/3* gene fusion–positive metastatic NSCLC and decided that entrectinib and larotrectinib are preferred first-line therapy options. Systemic therapy regimens are also recommended (useful in certain circumstances) for patients with *NTRK1/2/3* gene fusions; the same initial systemic regimens used for patients with metastatic NSCLC may be used (eg, carboplatin plus paclitaxel).

Larotrectinib

Larotrectinib is an oral TKI that inhibits *TRK* fusion proteins across a diverse range of solid tumors in younger and older patients with unresectable or metastatic disease; thus, larotrectinib is an age- and tumor-agnostic therapy.³³⁰ A study in 55 patients with *NTRK* gene fusion–positive disease across a range of solid tumors showed that larotrectinib yielded an overall response rate of 75% (95% CI, 61%–85%).³³⁰ An updated analysis showed that 90% of patients were still alive after 1 year, 18% of patients had a complete response, 69% of patients were still responding, and 58% of patients did not have disease progression.⁸³⁴ An additional 35 patients with *NTRK* gene fusion–positive disease had an overall response rate of 74%.⁸³⁴ Fewer than 3% of patients had adverse events of grade 3 to 4. A combined analysis of pediatric and adult patients reported an overall response rate of 79% (95% CI, 72%–85%).

The NCCN NSCLC Panel recommends larotrectinib as either a first-line or subsequent therapy option for patients with *NTRK1/2/3* gene fusion–

positive metastatic NSCLC based on these data.^{330,834} Larotrectinib may be used as a subsequent therapy option for patients with *NTRK1/2/3* gene fusion–positive metastatic NSCLC if larotrectinib or entrectinib were not previously given as first-line therapy. The panel has preference stratified the systemic therapy options for patients with *NTRK1/2/3* gene fusion–positive metastatic NSCLC and decided that larotrectinib and entrectinib are preferred first-line therapy options. Other systemic therapy regimens (eg, carboplatin plus paclitaxel) are also recommended as either first-line or subsequent therapy options and categorized as useful in certain circumstances.

Oral TKIs that Inhibit RET Rearrangements

Previously, the panel deleted vandetanib as an option for *RET* rearrangement–positive metastatic NSCLC because there are better therapy options.^{343,344}

Cabozantinib

Cabozantinib is an oral TKI that inhibits *RET* rearrangements but also inhibits other kinases. A phase 2 study assessed cabozantinib in 26 patients.^{341,342,930} The overall response rate was 28% (95% CI, 12%–49%). Many patients (19 [73%]) needed dose reductions because of adverse events. The most common grade 3 adverse events included lipase elevation (4 patients [15%]), increased ALT (2 [8%]), decreased platelet count (2 [8%]), and hypophosphatemia (2 [8%]). The NCCN NSCLC Panel recommends cabozantinib as a first-line or subsequent therapy option (useful in certain circumstances) for *RET* rearrangement–positive metastatic NSCLC based on these data.^{341,342} Cabozantinib may be used as a subsequent therapy option if pralsetinib, selpercatinib, or cabozantinib were not previously given as first-line therapy for *RET* rearrangement–positive metastatic NSCLC.

Pralsetinib



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

Pralsetinib is an oral TKI that selectively inhibits *RET* rearrangements. ARROW, a phase 1/2 study, assessed pralsetinib in patients with metastatic NSCLC and *RET* rearrangements.³³⁹ First-line therapy with pralsetinib yielded an overall response rate of 70% (19/27; 95% CI, 50%–86%); 3 patients (11%) had a complete response. Second-line therapy with pralsetinib yielded an overall response rate of 61% (53/87; 95% CI, 50%–71%); 5 patients (6%) had a complete response. Nine patients had measurable brain metastases, and 56% of them responded to pralsetinib; 3 patients had an intracranial complete response. Grade 3 or more adverse events with pralsetinib include anemia (10%), neutropenia (18%), and hypertension (11%). Common adverse events with pralsetinib included increased AST levels (31%), increased ALT levels (21%), anemia (22%), hypertension (20%), constipation (21%), and neutropenia (19%). Only 4% of patients (5/132) had to stop taking pralsetinib because of side effects. No treatment-related deaths were reported. The NCCN NSCLC Panel recommends pralsetinib as a first-line or subsequent therapy option (preferred) for patients with metastatic NSCLC who are positive for *RET* rearrangements based on clinical trial data and FDA approval.³³⁹ Pralsetinib may be used as a subsequent therapy option for patients with *RET* rearrangement–positive metastatic NSCLC if pralsetinib, selpercatinib, or cabozantinib were not previously given as first-line therapy.

Selpercatinib

Selpercatinib is an oral TKI that selectively inhibits *RET* rearrangements. Libretto-001, a phase 1/2 study, assessed selpercatinib in patients with NSCLC and *RET* rearrangements.^{340,931} Updated results from Libretto-001 show that first-line therapy with selpercatinib yielded an overall response rate of 85% (33/39; 95% CI, 70%–94%).³⁴⁰ Second-line therapy with selpercatinib yielded an overall response rate of 64% (67/105; 95% CI, 54%–73%); the median PFS was 18.4 months (95% CI, 16.4–24.8). Of patients with brain metastases, 91% (10/11) responded to selpercatinib.

Common grade 3 or more adverse events with selpercatinib included hypertension (14%), increased liver enzyme levels (12%), hyponatremia (6%), and lymphopenia (6%). Only 2% of patients (12/531) had to stop taking selpercatinib because of side effects.

The NCCN NSCLC Panel recommends selpercatinib as a first-line or subsequent therapy option (preferred) for patients with metastatic NSCLC who are positive for *RET* rearrangements based on clinical trial data and FDA approval.^{926,931} Selpercatinib may be used as a subsequent therapy option for patients with for *RET* rearrangement–positive metastatic NSCLC if selpercatinib, pralsetinib, or cabozantinib were not previously given as first-line therapy.

Preference Stratification

The NCCN NSCLC Panel preference stratified the recommended regimens for *RET* rearrangement–positive metastatic NSCLC and decided that selpercatinib and pralsetinib are preferred first-line or subsequent therapy options based on clinical trial data.^{339,340} The panel decided that cabozantinib is useful in certain circumstances.^{341,342,930} Selpercatinib, pralsetinib, or cabozantinib may be used as subsequent therapy options if they were not previously given as first-line therapy for *RET* rearrangement–positive metastatic NSCLC. Systemic therapy regimens (other recommended regimens) are also recommended as first-line therapy options for patients with metastatic NSCLC who are positive for *RET* rearrangements; these systemic regimens include platinum doublets (eg, carboplatin plus paclitaxel). These platinum doublets may be used as subsequent therapy options for patients with disease progression on pralsetinib, selpercatinib, or cabozantinib.

Oral TKIs that Inhibit ROS1 Rearrangements

Although ROS1 is a distinct receptor tyrosine kinase, it is very similar to ALK and members of the insulin receptor family.^{345,346} It is estimated that



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

ROS1 gene rearrangements occur in about 1% to 2% of patients with NSCLC.³⁴⁶⁻³⁴⁹ The NCCN Panel recommends testing for *ROS1* rearrangements in all patients with metastatic nonsquamous NSCLC or NSCLC NOS based on clinical trial data and FDA approvals of several agents.^{330,332,350,357,834,929} *ROS1* rearrangement testing can be considered in patients with metastatic squamous cell carcinoma. The panel recommends four agents for patients with *ROS1*-positive metastatic NSCLC—ceritinib, crizotinib, entrectinib, and lorlatinib—based on clinical trial data and FDA approvals, which are described in the following sections.

Ceritinib

Ceritinib is a second-generation oral TKI that inhibits *ALK* and *ROS1* rearrangements.⁸⁴⁸ A phase 2 trial assessed ceritinib as first-line therapy in patients (n = 28 evaluable) with NSCLC and *ROS1* rearrangements.⁸⁴⁸ One complete response and 19 partial responses (overall response rate, 62% [95% CI, 45%–77%]) were reported in patients receiving ceritinib. PFS was 19.3 months (95% CI, 1–37) for crizotinib-naïve patients and 9.3 months (95% CI, 0–22) for all patients. The median overall survival was 24 months (95% CI, 5–43).

The NCCN NSCLC Panel recommends ceritinib as a first-line therapy option for patients with *ROS1*-positive metastatic NSCLC based on clinical trial data.⁸⁴⁸ Ceritinib is an option if a *ROS1* rearrangement is discovered before giving, or during, first-line systemic therapy (eg, carboplatin plus paclitaxel). The NCCN NSCLC Panel has preference stratified the first-line therapy options for patients with *ROS1*-positive metastatic NSCLC. The panel decided that crizotinib and entrectinib are preferred first-line therapy options for patients with *ROS1*-positive metastatic NSCLC because they are better tolerated, have been assessed in more patients, and are approved by the FDA. Ceritinib is an “other recommended” option for patients with *ROS1*-positive metastatic NSCLC. Lorlatinib is

recommended as a subsequent therapy option in certain patients with *ROS1*-positive metastatic NSCLC whose disease becomes resistant to ceritinib, crizotinib, or entrectinib.³⁵⁸ However, entrectinib is recommended as a subsequent therapy option for patients with CNS progression after crizotinib or ceritinib.

Crizotinib

Crizotinib is a first-generation oral TKI that inhibits *ALK* rearrangements, *ROS1* rearrangements, and some *MET* tyrosine kinases (high-level *MET* amplification or *MET*ex14 skipping mutation).^{192,221,320,851-855} Crizotinib is very effective for patients with *ROS1* rearrangements with response rates of about 70% to 80% including complete responses.^{192,346,351,932,933} A phase 2 trial assessed crizotinib in 127 East Asian patients with *ROS1*-positive advanced NSCLC who had received 3 or fewer lines of therapy. The overall response rate was 72% (95% CI, 63%–79%) with 17 complete responses; the median duration of response was 19.7 months (95% CI, 14.1–not reached). The median PFS was 15.9 months (95% CI, 12.9–24.0).⁹³³

PROFILE 1001, a phase 2 study, assessed crizotinib in 50 patients with advanced NSCLC and *ROS1* rearrangements.³⁴⁶ Crizotinib yielded an objective response rate of 72% (95% CI, 58%–84%); there were 3 complete responses and 33 partial responses.³⁴⁶ The median duration of response was 17.6 months (95% CI, 14.5–not reached), and the median PFS was 19.2 months (95% CI, 14.4–not reached). Updated results from PROFILE 1001 reported an overall response rate of 72% (95% CI, 58%–83%) with crizotinib including 6 confirmed complete responses in 53 patients with *ROS1*-positive advanced NSCLC.²⁵ The median overall survival was 51.4 months (95% CI, 29.3–not reached). No grade 4 or higher treatment-related adverse events were reported.



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

The EUCROSS study reported that crizotinib yielded an overall response rate of 70% (21/30; 95% CI, 51%–85%) in 30 patients with *ROS1*-positive advanced NSCLC.⁹³² Adverse events related to treatment occurred in 97% (33/34) of patients. A retrospective European study in patients (n = 30 evaluable) with stage IV NSCLC and *ROS1* rearrangements also

assessed crizotinib.¹⁹² There were 5 complete responses (overall response rate, 80%; disease control rate, 86.7%). The median PFS was 9.1 months. Many patients (n = 26) received pemetrexed (either alone or in combination with platinum and either before or after crizotinib) and had a response rate of 57.7% and a median PFS of 7.2 months.

The NCCN NSCLC Panel recommends *ROS1* testing in certain patients with metastatic NSCLC based on data showing the efficacy of several agents for patients with *ROS1* rearrangements and on FDA approvals (see *Principles of Molecular and Biomarker Analysis* in the NCCN Guidelines for NSCLC).^{192,346,351} The panel recommends crizotinib as a first-line treatment option for patients with *ROS1*-positive metastatic NSCLC based on clinical trial data and FDA approval.^{25,346,932,933} Crizotinib is a preferred option if a *ROS1* rearrangement is discovered before giving, or during, first-line systemic therapy (eg, carboplatin plus paclitaxel). The NCCN NSCLC Panel decided that crizotinib and entrectinib are preferred agents for first-line therapy in patients with *ROS1*-positive metastatic NSCLC, compared with ceritinib, because they are better tolerated, have been assessed in more patients, and are approved by the FDA. Lorlatinib is recommended as a subsequent therapy option in patients with *ROS1*-positive metastatic NSCLC whose disease becomes resistant to crizotinib, ceritinib, or entrectinib.³⁵⁸ However, entrectinib is recommended as a subsequent therapy option for patients with CNS progression after crizotinib or ceritinib.³⁵⁹

Crizotinib is a first-generation oral TKI that inhibits *ALK* rearrangements, *ROS1* rearrangements, and some *MET* tyrosine kinases (high-level *MET*

amplification or *MET*ex14 skipping mutation).^{192,221,320,851-855} Crizotinib is very effective for patients with *ROS1* rearrangements with response rates of about 70% to 80% including complete responses.^{192,346,351,932,933} A phase 2 trial assessed crizotinib in 127 East Asian patients with *ROS1*-positive advanced NSCLC who had received 3 or fewer lines of therapy. The overall response rate was 72% (95% CI, 63%–79%) with 17 complete responses; the median duration of response was 19.7 months (95% CI, 14.1–not reached). The median PFS was 15.9 months (95% CI, 12.9–24.0).⁹³³

PROFILE 1001, a phase 2 study, assessed crizotinib in 50 patients with advanced NSCLC and *ROS1* rearrangements.³⁴⁶ Crizotinib yielded an objective response rate of 72% (95% CI, 58%–84%); there were 3 complete responses and 33 partial responses.³⁴⁶ The median duration of response was 17.6 months (95% CI, 14.5–not reached), and the median PFS was 19.2 months (95% CI, 14.4–not reached). Updated results from PROFILE 1001 reported an overall response rate of 72% (95% CI, 58%–83%) with crizotinib including 6 confirmed complete responses in 53 patients with *ROS1*-positive advanced NSCLC.²⁵ The median overall survival was 51.4 months (95% CI, 29.3–not reached). No grade 4 or higher treatment-related adverse events were reported.

The EUCROSS study reported that crizotinib yielded an overall response rate of 70% (21/30; 95% CI, 51%–85%) in 30 patients with *ROS1*-positive advanced NSCLC.⁹³² Adverse events related to treatment occurred in 97% (33/34) of patients. A retrospective European study in patients (n = 30 evaluable) with stage IV NSCLC and *ROS1* rearrangements also assessed crizotinib.¹⁹² There were 5 complete responses (overall response rate, 80%; disease control rate, 86.7%). The median PFS was 9.1 months. Many patients (n = 26) received pemetrexed (either alone or in combination with platinum and either before or after crizotinib) and had a response rate of 57.7% and a median PFS of 7.2 months.



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

The NCCN NSCLC Panel recommends *ROS1* testing in certain patients with metastatic NSCLC based on data showing the efficacy of several agents for patients with *ROS1* rearrangements and on FDA approvals (see *Principles of Molecular and Biomarker Analysis* in the NCCN Guidelines for NSCLC).^{192,346,351} The panel recommends crizotinib as a first-line treatment option for patients with *ROS1*-positive metastatic NSCLC based on clinical trial data and FDA approval.^{25,346,932,933} Crizotinib is a preferred option if a *ROS1* rearrangement is discovered before giving, or during, first-line systemic therapy (eg, carboplatin plus paclitaxel). The NCCN NSCLC Panel decided that crizotinib and entrectinib are preferred agents for first-line therapy in patients with *ROS1*-positive metastatic NSCLC, compared with ceritinib, because they are better tolerated, have been assessed in more patients, and are approved by the FDA. Lorlatinib is recommended as a subsequent therapy option in patients with *ROS1*-positive metastatic NSCLC whose disease becomes resistant to crizotinib, ceritinib, or entrectinib.³⁵⁸ However, entrectinib is recommended as a subsequent therapy option for patients with CNS progression after crizotinib or ceritinib.³⁵⁹

Entrectinib

Entrectinib is an oral TKI that inhibits several tyrosine kinases including *ROS1* and *TRK* (see *ROS1 Rearrangements* and *NTRK1/2/3 Gene Rearrangements* in this Discussion).^{332,934} Entrectinib has been assessed in several phase 1 and 2 trials in patients with *ROS1*-positive metastatic NSCLC (ie, phase 2 STARTRK-2 trial, phase 1 STARTRK-1 trial, phase 1 ALKA-372-001 trial).^{350,929} Pooled data from these 3 trials in 53 patients with *ROS1*-positive metastatic NSCLC receiving first-line entrectinib showed an overall response rate of 77% (41/53; 95% CI, 64%–88%; 3 complete responses).^{350,359} The intracranial overall response rate was 55% (95% CI, 32%–77%; 4 complete responses, 7 partial responses).^{350,929} In the larger *ROS1* population (n = 134), grade 3 to 4 adverse events were seen in 34% of patients. Fifteen patients had serious adverse events such

as nervous system disorders (4 patients [3%]) and cardiac disorders (3 patients [2%]). No treatment-related deaths were reported. Although entrectinib has better CNS penetration than crizotinib, it is more toxic.³⁵⁹

The NCCN NSCLC Panel recommends entrectinib as a first-line therapy option (preferred) for patients with *ROS1*-positive metastatic NSCLC based on these data and FDA approval.^{332,350,357,929} Entrectinib is recommended as a subsequent therapy option for patients with *ROS1*-positive metastatic NSCLC and disease progression with brain metastases on crizotinib or ceritinib.³⁵⁹ Subsequent therapy with lorlatinib is also recommended for select patients with *ROS1*-positive metastatic NSCLC and disease progression after treatment with crizotinib, ceritinib, or entrectinib. The panel has preference stratified the recommended regimens for patients with *ROS1*-positive metastatic NSCLC and decided that entrectinib and crizotinib are preferred first-line therapy options; ceritinib is stratified as an “other recommended” option.

Lorlatinib

Lorlatinib is an oral third-generation TKI that targets *ALK* and *ROS1* tyrosine kinases and has good CNS penetration; it inhibits a broad range of *ALK* resistance mutations that develop after treatment with first- and second-generation *ALK* inhibitors.^{358,831-833} A phase 1 to 2 trial assessed lorlatinib in patients with *ROS1*-positive metastatic NSCLC.³⁵⁸ Many patients (58% [40/69]) had previously received crizotinib; some patients were TKI naïve (30% [21/69]). Objective responses were achieved in 35% (14/40) of patients who had previously received crizotinib and 62% (13/21) of TKI-naïve patients. An intracranial response was observed in 50% (12/24) of patients who had previously received crizotinib and 64% (7/11) of TKI-naïve patients. Serious treatment-related adverse events occurred in 7% (5/69) of patients; no treatment-related deaths were reported.

The NCCN NSCLC Panel recommends lorlatinib as a subsequent therapy option for select patients with *ROS1*-positive metastatic NSCLC and



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

disease progression after treatment with ceritinib, crizotinib, or entrectinib, depending on the type of progression.³⁵⁸ However, entrectinib is recommended as a subsequent therapy option for patients with CNS progression after crizotinib or ceritinib. For the 2023 update (Version 1), the NCCN Panel now recommends lorlatinib as a subsequent therapy option, if not previously given, for patients with *ROS1*-positive metastatic NSCLC and symptomatic brain metastases.^{935,936}

Agents that Inhibit VEGF or VEGF Receptors

Bevacizumab

Bevacizumab is a recombinant monoclonal antibody that targets VEGF. ECOG 4599, a phase 3 randomized trial, assessed bevacizumab added to paclitaxel plus carboplatin versus chemotherapy alone in patients with recurrent or advanced nonsquamous NSCLC (stage IIIB–IV).⁸⁰⁶ In the bevacizumab plus chemotherapy group, median survival was 12.3 months versus 10.3 months with chemotherapy alone (HR for death, 0.79; $P=.003$). Clinically significant bleeding occurred more often with bevacizumab plus chemotherapy versus chemotherapy alone (4.4% vs. 0.7%, respectively; $P < .001$). Fifteen treatment-related deaths were reported with bevacizumab plus chemotherapy.

Bevacizumab may be added to the following regimens: carboplatin plus paclitaxel (category 1), carboplatin plus pemetrexed, or cisplatin plus pemetrexed. The NCCN NSCLC Panel has preference stratified the systemic therapy regimens and decided that these specific bevacizumab plus chemotherapy options are “useful in certain circumstances” for eligible patients with metastatic NSCLC.^{806,937} These bevacizumab plus chemotherapy regimens are first-line therapy options for patients with PS 0 to 1, nonsquamous NSCLC or NSCLC NOS, negative test results for actionable driver mutations, PD-L1 expression less than 1%, and/or contraindications to PD-1 or PD-L1 inhibitors. Bevacizumab plus chemotherapy regimens may be used as first-line or subsequent therapy

options, although they are not preferred options, in eligible patients with actionable driver mutations as shown in the algorithm. Targeted therapy is a preferred first-line therapy option for patients with actionable driver mutations, except for *EGFR* exon 20 mutations or *KRAS* mutations.

Bevacizumab in combination with a PD-L1 inhibitor plus chemotherapy (eg, ABCP) is a first-line therapy option (category 1, other recommended) regardless of PD-L1 expression for patients with PS 0 to 1, nonsquamous NSCLC or NSCLC NOS, negative test results for actionable driver mutations, and no contraindications to PD-1 or PD-L1 inhibitors or bevacizumab (see *Immune Checkpoint Inhibitors* in this Discussion).

Combination therapy with the ABCP regimen may be used as a first-line or subsequent therapy option, although it is not a preferred option, in eligible patients with actionable driver mutations as shown in the algorithm.

Targeted therapy is a preferred first-line therapy option for patients with actionable driver mutations, except for *EGFR* exon 20 mutations or *KRAS* mutations.

To receive treatment with bevacizumab and chemotherapy, patients must meet the following criteria: nonsquamous NSCLC and no recent history of hemoptysis. Any regimen with a high risk for thrombocytopenia—and, therefore, possible bleeding—should be used with caution when combined with bevacizumab. Bevacizumab is not recommended for patients with squamous cell NSCLC. The NCCN NSCLC Panel recommends that bevacizumab biosimilars may be used in any of the recommended regimens containing bevacizumab that are used for eligible patients with metastatic NSCLC.⁸⁹⁶⁻⁹⁰⁰

NEJ026, a phase 3 randomized trial, compared first-line bevacizumab plus erlotinib versus erlotinib alone in 228 patients with *EGFR*-positive advanced nonsquamous NSCLC.^{892,893} At interim analysis, PFS was 16.9 months (95% CI, 14.2–21.0) for erlotinib plus bevacizumab versus 13.3 months (95% CI, 11.1–15.3) for erlotinib alone (HR, 0.605; 95% CI, 0.417–



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

0.877; $P = .016$).⁸⁹³ Grade 3 or worse events occurred in 88% (98/112) of patients receiving erlotinib/bevacizumab versus 46% (53/114) of those receiving erlotinib alone. Grade 4 adverse events occurred in 8% (9/112) of patients receiving erlotinib plus bevacizumab (including neutropenia, hepatic dysfunction) versus 4% (5/114) of patients receiving erlotinib alone (hepatic dysfunction). No treatment-related deaths were reported.^{892,893} Updated data show that median overall survival was 50.7 months (95% CI, 37.3–not estimable) in those receiving erlotinib plus bevacizumab versus 46.2 months (95% CI, 38.2–not estimable) in those receiving erlotinib alone (HR, 1.007; 95% CI, 0.681–1.490; $P = .97$).⁸⁹² The NCCN NSCLC Panel recommends erlotinib plus bevacizumab as a first-line therapy option for eligible patients with metastatic nonsquamous NSCLC, *EGFR* exon 19 deletions or *EGFR* exon 21 L858R mutations, and no contraindications to bevacizumab (other recommended) based on clinical trial data.^{892,893} The panel recommends osimertinib as a preferred first-line therapy option for patients with *EGFR* positive metastatic NSCLC.

Ramucirumab

Ramucirumab is a recombinant monoclonal antibody that targets VEGF receptors.

First-Line Therapy

RELAY, a phase 3 randomized trial, compared first-line therapy with ramucirumab plus erlotinib versus erlotinib alone in patients with advanced NSCLC and the common *EGFR* mutations.⁸⁹¹ PFS was 19.4 months (95% CI, 15.4–21.6) with ramucirumab plus erlotinib versus 12.4 months (95% CI, 11.0–13.5) with erlotinib alone (HR, 0.59; 95% CI, 0.46–0.76; $P < .0001$). Serious adverse events (grade 3–4) occurred in 72% (159/221) of patients receiving erlotinib plus ramucirumab (including hypertension) versus 54% (121/225) in those receiving erlotinib alone (including increased ALT). One treatment-related death (hemothorax) occurred in a patient receiving erlotinib plus ramucirumab. The NCCN NSCLC Panel

recommends erlotinib plus ramucirumab as a first-line therapy option for patients with metastatic NSCLC and *EGFR* exon 19 deletions or *EGFR* exon 21 L858R mutations (other recommended intervention) based on clinical trial data.⁸⁹¹ The panel recommends osimertinib as a preferred first-line therapy option for patients with *EGFR*-positive metastatic NSCLC.

Subsequent Therapy

REVEL, a phase 3 randomized trial, assessed ramucirumab plus docetaxel versus docetaxel alone in patients with metastatic NSCLC and disease progression.⁹³⁸ The median overall survival was 10.5 months for ramucirumab plus docetaxel versus 9.1 months for docetaxel alone (HR, 0.86; 95% CI, 0.75–0.98; $P < .023$). More than 70% of patients had grade 3 or higher adverse events in both groups (79% for ramucirumab plus docetaxel vs. 71% for docetaxel alone). Adverse events of special concern with ramucirumab plus docetaxel therapy include risk for severe hemorrhage, grade 3 to 4 gastrointestinal bleeding, gastrointestinal perforation or fistula, impaired wound healing, and poorly controlled hypertension. There were 16 deaths from grade 3 or worse pulmonary hemorrhage and other adverse events in the REVEL trial: there were 8 deaths in the ramucirumab plus docetaxel arm and 8 deaths in the docetaxel alone arm. The NCCN NSCLC Panel recommends ramucirumab plus docetaxel as a subsequent therapy option for patients with metastatic NSCLC, regardless of histology, based on clinical data and FDA approval.^{938,939}

Agents that Inhibit High-Level *MET* Amplifications

High-level *MET* amplifications are designated as emerging biomarkers in the NCCN Guidelines because there is less evidence for using targeted agents for these biomarkers. In addition, the recommended agents have not been FDA approved for these mutations in patients with NSCLC, although they are approved for other cancers. The panel decided that



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

optimal biomarker testing should include testing for high-level *MET* amplifications in all patients with metastatic nonsquamous NSCLC or NSCLC NOS based on clinical trial data.^{167-169,172,940,941} Testing for high-level *MET* amplifications can be considered in patients with squamous cell carcinoma. The definition of high-level *MET* amplification is evolving and may differ depending on which assay is used. When using NGS, a copy number greater than 10 is consistent with high-level *MET* amplification.¹⁷²

Capmatinib

Capmatinib is an oral TKI that selectively inhibits *MET* genomic alterations. GEOMETRY assessed capmatinib in patients with metastatic NSCLC and high-level *MET* amplification.¹⁷² For first-line capmatinib, response rates were 40% (95% CI, 16%–68%); for second-line capmatinib, response rates were 29% (95% CI, 19%–41%). Adverse events for patients with high-level *MET* amplification across all cohorts included peripheral edema (49%), vomiting (30%), and nausea (48%); most of these events were grade 1 or 2. Grade 3 to 4 adverse events occurred in 68% of patients. Another trial assessed capmatinib in patients with *MET* alterations. The median overall survival was 21.5 months (95% CI, 20.8–not applicable) for 10 patients with high-level *MET* amplification who received capmatinib.⁹⁴⁰

The NCCN NSCLC Panel recommends capmatinib for patients with advanced NSCLC and high-level *MET* amplification based on clinical trial data; tepotinib and crizotinib are also recommended options in this setting.^{167-169,172,941}

Crizotinib

Crizotinib is an oral TKI that inhibits some *MET* tyrosine kinases (high-level *MET* amplification or *MET*ex14 skipping mutation), *ALK* rearrangements, and *ROS1* rearrangements. A subgroup analysis of

PROFILE 1001 assessed crizotinib in patients with advanced NSCLC and different levels of *MET* amplification; 40% (6/15) of patients responded to crizotinib if they had *MET* amplification with a GCN of 6 or more by NGS.⁹⁴¹ Two of these patients whose disease responded had concurrent *MET*ex14 skipping mutations. The overall response rate to crizotinib was 29% in patients with a GCN of 10 or more. Patients who had concurrent *KRAS*, *BRAF*, or *EGFR* mutations did not respond to crizotinib. Most patients had adenocarcinoma and had received at least one line of therapy. The median overall survival was 11.4 months (95% CI, 7.2–19.3) in the group with higher *MET* amplification. There was one treatment-related death.

A patient with stage IV moderately differentiated adenocarcinoma had a partial response on first-line therapy with carboplatin plus gemcitabine plus bevacizumab; she received maintenance therapy with bevacizumab.¹⁶⁷ Her tumor was found to have high-level *MET* amplification (*MET*/*CEPT* ratio >5.0) and was negative for *ALK* rearrangements. The patient had a rapid and durable response to single-agent therapy with crizotinib (54.8% reduction in aggregate tumor measurement). She had mild, grade 1 side effects including asymptomatic sinus bradycardia and transient visual disturbances.

The NCCN NSCLC Panel recommends crizotinib for patients with advanced NSCLC and high-level *MET* amplification based on clinical trial data; tepotinib and capmatinib are also recommended options in this setting.^{167-169,172,941}

Tepotinib

Tepotinib is an oral TKI that selectively inhibits *MET*ex14 skipping mutations and high-level *MET* amplification; it is approved by the FDA for patients with metastatic NSCLC who have *MET*ex14 skipping mutations. Another cohort of the VISION trial assessed tepotinib in 24 patients with advanced NSCLC and *MET* amplification but without *MET*ex14 skipping



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

mutations.¹⁶⁹ Preliminary data suggest the overall response rate is about 42% (10/24). The NCCN NSCLC Panel recommends tepotinib for patients with advanced NSCLC and high-level *MET* amplification based on preliminary data; capmatinib and crizotinib are also recommended options in this setting.^{167-169,172}

Immune Checkpoint Inhibitors

Human ICI antibodies inhibit the PD-1 receptor or PD-L1, which improves antitumor immunity; PD-1 receptors are expressed on activated cytotoxic T cells.³⁶⁰⁻³⁶² Cemiplimab-rwlc, nivolumab, and pembrolizumab inhibit PD-1 receptors;^{363,364,141} atezolizumab and durvalumab inhibit PD-L1.^{365,366} The single-agent immunotherapy or combination immunotherapy plus chemotherapy regimens are not recommended if patients have contraindications to immunotherapy, which may include active or previously documented autoimmune disease and/or current use of immunosuppressive agents. PD-L1 inhibitors are less beneficial in patients with some oncogenic drivers (ie, *EGFR* exon 19 deletions, *EGFR* exon 21 L858R mutations, *ALK* rearrangements).¹⁸⁶ Monitoring is recommended during initial therapy with response assessment with CT, with or without contrast, of known or high-risk sites of disease after 2 cycles and then every 2 to 4 cycles. During maintenance or subsequent therapy, monitoring is also recommended with CT, with or without contrast, every 6 to 12 weeks. ICIs (also known as immunotherapy or IO agents) are associated with a delay in benefit when compared with targeted therapy or cytotoxic chemotherapy.

The NCCN NSCLC Panel recommends IHC testing for PD-L1 expression (category 1) before first-line treatment in all patients with metastatic NSCLC, regardless of histology, based on the efficacy of pembrolizumab with or without chemotherapy.³⁶⁷ Ideally, PD-L1 expression levels are assessed before first-line therapy in patients with metastatic NSCLC, if clinically feasible. Every effort also needs to be made to assess for

oncogenic driver variants for which targeted therapies are available (eg, *EGFR* mutations, *ALK* rearrangements). Plasma-based ctDNA testing can be used to evaluate for genomic alterations if the assay has been validated and if the patient is unfit for invasive tissue sampling or if there is insufficient tissue for molecular analyses; however, plasma-based ctDNA testing is less sensitive than tissue assays.

It is important to note that targeted therapies are recommended for patients with metastatic NSCLC and specific oncogenic drivers, independent of PD-L1 levels. Patients with metastatic NSCLC and PD-L1 expression levels of 1% or more—but who also have a targetable driver oncogene molecular variant (eg, *EGFR*, *ALK*, *ROS1*)—should receive first-line targeted therapy for that oncogene and not first-line ICIs, because targeted therapies yield higher response rates (eg, osimertinib, 80%) than ICIs (lower response rates) in the first-line setting, targeted therapy is better tolerated, and these patients are less likely to respond to single-agent ICIs.^{186,373-375,840} For patients receiving first-line ICIs with or without chemotherapy, oncologists should be aware of the long half-life of the ICIs and potential adverse effects when using osimertinib in combination with or following ICIs.^{907,908} A few case reports suggest that patients treated with ICIs who have acquired resistance may have transformation to SCLC.⁹⁴²

In 2020, the NCCN Panel removed TMB as an emerging immune biomarker for patients with metastatic NSCLC based on clinical trial data, concerns about variable TMB measurements, and other issues as previously described (see *TMB* in this Discussion).^{173,377,381} The NCCN Guidelines do not recommend measurement of TMB levels before deciding whether to use nivolumab plus ipilimumab regimens or other ICIs, such as pembrolizumab.¹⁷³

The following content briefly summarizes the use of ICIs as first-line or subsequent therapy options in eligible patients with metastatic NSCLC;



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

detailed information, including clinical trial data, is provided in subsequent sections.⁹⁴³ ICIs may also be used in other settings; clinical trial data and appropriate use are described in greater detail elsewhere. For example, adjuvant atezolizumab or pembrolizumab are recommended following adjuvant chemotherapy for eligible patients with completely resected early-stage NSCLC (see *Surgery Followed by Adjuvant Therapy: Trial Data and NCCN Recommendations* in this Discussion).^{32,751} Durvalumab is recommended as a consolidation immunotherapy option for eligible patients with unresectable stage II or III NSCLC and without disease progression after treatment with definitive concurrent chemoradiation (see *Chemoradiation: Trial Data and NCCN Recommendations* in this Discussion).³⁶⁵

Single-agent pembrolizumab, atezolizumab, or cemiplimab-rwlc are recommended (category 1; preferred) as first-line therapy options for eligible patients with metastatic NSCLC regardless of histology, PD-L1 expression levels of 50% or more, and negative test results for actionable driver mutations that have recommended first-line targeted therapies (ie, *ALK*, *BRAF* p.V600E, *EGFR*, *MET*ex14 skipping, *NTRK1/2/3*, *RET*, *ROS1*). The NCCN NSCLC Panel recommends single-agent pembrolizumab as a first-line therapy option in eligible patients with metastatic NSCLC regardless of histology, PD-L1 levels of 1% to 49% (category 2B; useful in certain circumstances), and negative test results for actionable driver mutations.⁹⁴⁴

Combination therapy with pembrolizumab plus chemotherapy is recommended (category 1; preferred) as a first-line therapy option in eligible patients with metastatic NSCLC and negative test results for actionable driver mutations, regardless of PD-L1 expression levels. Combination therapy with the ABCP regimen is recommended (category 1; other recommended intervention) as a first-line therapy option for eligible patients with metastatic NSCLC and negative test results for

actionable driver mutations, regardless of PD-L1 expression levels. Combination therapy with pembrolizumab plus chemotherapy or with ABCP may be used as first-line or subsequent therapy options, although they are not preferred options, in eligible patients with actionable driver mutations as shown in the algorithm. Targeted therapy is a preferred first-line therapy option for patients with actionable driver mutations, except for *EGFR* exon 20 mutations or *KRAS* mutations. Continuation maintenance immunotherapy is recommended for 2 years, if tolerated, for all first-line immunotherapy (± chemotherapy) regimens.

If patients have disease progression on PD-1/PD-L1 inhibitor therapy (± chemotherapy), then using a PD-1/PD-L1 inhibitor is not recommended for subsequent therapy.⁸⁹⁷ Single-agent pembrolizumab is recommended (category 1; preferred) as a subsequent therapy option for select patients with metastatic NSCLC and PD-L1 levels greater than 1%; nivolumab or atezolizumab is recommended (category 1; preferred) as a subsequent monotherapy option for select patients with metastatic NSCLC regardless of PD-L1 levels. Based on data in the second-line setting, PD-1 or PD-L1 inhibitor monotherapy appears to be less effective in patients with *EGFR* exon 19 deletions, *EGFR* exon 21 L858R mutations, or *ALK* rearrangements, regardless of PD-L1 expression levels.^{360,364,840,945,946} A small study suggests that single-agent pembrolizumab is not effective as first-line therapy in patients with metastatic NSCLC and *EGFR* mutations, even those with PD-L1 levels more than 50%.³⁷⁴ Patients with *ALK*-positive NSCLC and very high PD-L1 expression levels do not respond to pembrolizumab.^{186,840} In the trials assessing the efficacy of first-line therapy with pembrolizumab with (or without) chemotherapy, most of the patients were wild type for *EGFR* or *ALK* variants. Maintenance immunotherapy is recommended, if tolerated, until progression for all the subsequent therapy regimens.



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

ICIs are associated with unique immune-mediated adverse events, such as endocrine disorders, that are not seen with traditional cytotoxic chemotherapy; therefore, health care providers should be aware of the spectrum of potential immune-mediated adverse events, know how to manage the adverse events, and educate their patients about possible side effects (see the NCCN Guidelines for the Management of Immunotherapy-Related Toxicities, available at www.NCCN.org).^{947,948}

Atezolizumab, cemiplimab-rwlc, durvalumab, nivolumab, or pembrolizumab should be discontinued for patients with severe or life-threatening pneumonitis or myocarditis and should be withheld or discontinued for other severe or life-threatening immune-mediated adverse events when indicated (see prescribing information).

Pseudoprogression has been reported; therefore, traditional RECIST criteria may not be applicable.⁹⁴⁹⁻⁹⁵¹

Atezolizumab

Atezolizumab is a humanized monoclonal immunoglobulin G1 antibody that inhibits PD-L1, which improves antitumor immune response.³⁶⁶

Immune-mediated adverse events may occur with ICIs, including atezolizumab.^{945,952} For patients with immune-mediated adverse events, intravenous high-dose corticosteroids should be administered based on the severity of the reaction (see the NCCN Guidelines for the Management of Immunotherapy-Related Toxicities, available at www.NCCN.org).

Atezolizumab should also be permanently discontinued for patients with severe or life-threatening pneumonitis or myocarditis and should be discontinued for other severe or life-threatening immune-mediated adverse events when indicated (see prescribing information). The following sections describe the use of atezolizumab in eligible patients with metastatic NSCLC. The use of adjuvant atezolizumab following adjuvant chemotherapy for eligible patients with completely resected early-stage NSCLC is described in a different section (see *Surgery Followed by*

Adjuvant Therapy: Trial Data and NCCN Recommendations in this Discussion).⁷⁵¹

First-Line Therapy

IMpower110, a phase 3 randomized trial, compared first-line therapy with single-agent atezolizumab versus platinum-based chemotherapy in three different subgroups of patients with metastatic NSCLC, including those with high PD-L1 expression (PD-L1 stained $\geq 50\%$ of TC [TC $\geq 50\%$] or PD-L1 stained tumor-infiltrating cells [IC] covering $\geq 10\%$ of the tumor area [IC $\geq 10\%$]); patients were wild type for *EGFR* or *ALK* variants and most either currently or previously smoked cigarettes.^{953,954} Patients receiving first-line atezolizumab monotherapy also received maintenance therapy with atezolizumab. Chemotherapy regimens for patients with nonsquamous NSCLC included cisplatin (or carboplatin) plus pemetrexed and maintenance therapy with pemetrexed; patients with squamous cell NSCLC received cisplatin plus gemcitabine and best supportive care as maintenance therapy. Atezolizumab monotherapy was associated with fatal adverse reactions in 3.8% of all patients (11/286, all 3 groups), including aspiration, COPD, pulmonary embolism, acute myocardial infarction, cardiac arrest, mechanical ileus, sepsis, cerebral infarction, and device occlusion; 4.2% of patients (11/263) receiving chemotherapy also died. Grade 3 to 4 treatment-related adverse events occurred in 12.9% of patients receiving atezolizumab monotherapy versus 44.1% with chemotherapy. The most frequent serious adverse reactions with atezolizumab monotherapy were pneumonia (2.8%), COPD (2.1%), and pneumonitis (2.1%); 28% of patients had serious adverse reactions.

It is important to note that a different IHC assay was used to test for PD-L1 levels in IMpower110 (SP142 PD-L1 IHC assay) compared with IHC



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

assays used for pembrolizumab monotherapy in KEYNOTE-024 (PD-L1 IHC 22C3 pharmDx assay); however, the results were similar regardless of which PD-L1 IHC assay was used.^{953,955} Data suggest that different methods of testing for PD-L1 levels are not equivalent.^{370,371} Based on an

analysis using the SP142 PD-L1 IHC assay, median OS was 20.2 months (95% CI, 16.5–not estimable) with atezolizumab monotherapy (n = 107) versus 13.1 months (95% CI, 7.4–16.5 months) with chemotherapy (n = 98) (HR, 0.59; 95% CI, 0.40–0.89; $P = .0106$) in patients with high PD-L1 expression. Based on an analysis using the 22C3 pharmDx assay, median OS was 20.2 months with atezolizumab monotherapy (n = 134) versus 11.0 months with chemotherapy (n = 126) (HR, 0.60; 95% CI, 0.41–0.86).^{953,955} There was no survival advantage in the other two subgroups of patients with lower PD-L1 expression (ie, TC \geq 5% or IC \geq 5%; TC \geq 1% or IC \geq 1%).

The NCCN NSCLC Panel recommends atezolizumab monotherapy (category 1; preferred) as a first-line therapy option for eligible patients with metastatic NSCLC based on clinical trial data and FDA approval.

^{953,954} Atezolizumab monotherapy is recommended as a first-line therapy option for patients with metastatic NSCLC, PD-L1 levels of 50% or more, and negative test results for actionable driver mutations, regardless of histology; maintenance therapy with atezolizumab is also recommended in this setting. Cemiplimab-rwlc and pembrolizumab are also recommended first-line therapy options in this setting (category 1; preferred). The NCCN NSCLC Panel has preference stratified the ICI regimens and decided that atezolizumab, cemiplimab-rwlc, and pembrolizumab (all are category 1) are preferred first-line therapy options for eligible patients with metastatic NSCLC based on clinical trial data.^{141,363,944,953}

IMpower150, a phase 3 randomized trial, compared first-line therapy with the ABCP regimen versus bevacizumab plus chemotherapy for patients with metastatic nonsquamous NSCLC.⁹⁵² Median overall survival was 19.2

months (95% CI, 17.0–23.8) in the ABCP arm versus 14.7 months (95% CI, 13.3–16.9) in the carboplatin plus paclitaxel plus bevacizumab arm; the HR for death was 0.78 (95% CI, 0.64–0.96; $P = .02$). PFS was longer in the ABCP arm versus chemotherapy plus bevacizumab (8.3 vs. 6.8 months; HR, 0.62; 95% CI, 0.52–0.74; $P < .001$). Some patients with *EGFR* mutations or *ALK* rearrangements (n = 108) and disease progression on (or were intolerant of) prior TKI were enrolled in this trial, although most patients (87%) did not have these genetic variants. In these patients with *EGFR* mutations or *ALK* rearrangements, PFS was also increased with ABCP compared with chemotherapy plus bevacizumab (9.7 vs. 6.1 months; HR, 0.59; 95% CI, 0.37–0.94). A subgroup analysis of IMpower150 reported that subsequent therapy with the ABCP regimen increased median overall survival in a few patients with *EGFR* mutation–positive metastatic NSCLC (n = 34) compared with those receiving carboplatin plus paclitaxel plus bevacizumab (n = 45).⁹⁵⁶ Therefore, the ABCP regimen may be an option for patients with *EGFR* mutations or *ALK* rearrangements and disease progression after initial therapy with TKIs.

The NCCN NSCLC Panel recommends the ABCP regimen (category 1; other recommended intervention) as a first-line therapy option for eligible patients with metastatic nonsquamous NSCLC (including adenocarcinoma) based on clinical trial data and FDA approval.⁹⁵² The ABCP regimen (also known as the quadruplicate regimen) is recommended as a first-line therapy option for patients with negative test results for actionable driver mutations (ie, *ALK*, *BRAF* p.V600E, *EGFR*, *MET*ex14 skipping, *NTRK1/2/3*, *RET*, and *ROS1*), regardless of PD-L1 expression levels. Combination therapy with the ABCP regimen may be used as a first-line or subsequent therapy option, although it is not a preferred option, in eligible patients with actionable driver mutations as shown in the algorithm. Targeted therapy is a preferred first-line therapy option for patients with actionable driver mutations, except for *EGFR* exon 20 mutations or *KRAS* mutations. Maintenance therapy with atezolizumab



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

and bevacizumab is also recommended in this setting (category 1) (see *Maintenance Therapy* in this Discussion). The NCCN NSCLC Panel has preference stratified the systemic therapy regimens and decided that the ABCP regimen is an other recommended intervention, because the NCCN NSCLC Panel prefers the pembrolizumab plus chemotherapy regimens based on tolerability and experience with these regimens. The NCCN NSCLC Panel recommends that bevacizumab biosimilars may be used in any of the systemic therapy regimens containing bevacizumab, such as ABCP, that are used for eligible patients with metastatic NSCLC based on clinical data and FDA approvals.⁸⁹⁶⁻⁹⁰⁰

IMpower130, a phase 3 randomized trial, compared atezolizumab plus carboplatin plus albumin-bound paclitaxel versus chemotherapy alone as first-line therapy in patients with metastatic nonsquamous NSCLC with no *EGFR* mutations or *ALK* rearrangements.⁹⁵⁷ Median overall survival was 18.6 months (95% CI, 16.0–21.2) in the atezolizumab plus chemotherapy arm versus 13.9 months (95% CI, 12.0–18.7) in carboplatin plus albumin-bound paclitaxel arm (HR, 0.79; 95% CI, 0.64–0.98; *P* = .033). Treatment-related deaths were reported in 2% (8/473) of patients in the atezolizumab plus chemotherapy arm and in less than 1% (1/232) of patients in the chemotherapy-only arm.

The NCCN NSCLC Panel recommends atezolizumab plus carboplatin plus albumin-bound paclitaxel (other recommended intervention) as a first-line therapy option for eligible patients with metastatic nonsquamous NSCLC based on clinical trial data.⁹⁵⁷ Atezolizumab plus carboplatin plus albumin-bound paclitaxel is recommended as a first-line therapy option for patients with metastatic NSCLC and negative test results for actionable driver mutations, regardless of PD-L1 levels. The atezolizumab plus carboplatin plus albumin-bound paclitaxel regimen may be used as a first-line or subsequent therapy option, although it is not a preferred option, in eligible patients with actionable driver mutations as shown in the

algorithm. Maintenance therapy with atezolizumab is also recommended in these settings. Targeted therapy is a preferred first-line therapy option for patients with actionable driver mutations, except for *EGFR* exon 20 mutations or *KRAS* mutations.

Subsequent Therapy

OAK, a phase 3 randomized trial, compared atezolizumab versus docetaxel in patients with metastatic NSCLC and disease progression during or after systemic therapy.^{945,958} Most patients either currently or previously smoked cigarettes and had received platinum-based chemotherapy; 10% of patients were not reported because they had *EGFR* mutations and *ALK* rearrangements.^{945,958} Patients with nonsquamous NSCLC who received atezolizumab had longer overall survival (15.6 months; 95% CI, 13.3–17.6) when compared with those receiving docetaxel (11.2 months; 95% CI, 9.3–12.6; HR, 0.73; 0.6–0.89; *P* = .0015). In patients with squamous cell NSCLC, overall survival was 8.9 months (95% CI, 7.4–12.8) in patients receiving atezolizumab versus 7.7 months (95% CI, 6.3–8.9) with docetaxel (HR, 0.73; 0.54–0.98; *P* = .038). Fewer patients were in the squamous group compared with the nonsquamous group (222 vs. 628). Fewer treatment-related severe adverse events (grades 3–4) were reported for atezolizumab versus docetaxel (15% vs. 43% [90/609 vs. 247/578]).

If patients have not previously received a PD-1/PD-L1 inhibitor, the NCCN NSCLC Panel recommends atezolizumab (category 1; preferred) as a subsequent therapy option for patients with metastatic nonsquamous or squamous cell NSCLC based on clinical trial data and FDA approval.^{366,945,958} Nivolumab and pembrolizumab are also recommended subsequent therapy options in this setting (category 1; preferred). Testing for PD-L1 expression levels is not required for prescribing subsequent therapy with atezolizumab or nivolumab but may provide useful



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

information. If patients have progressed on a PD-1/PD-L1 inhibitor, then using a PD-1/PD-L1 inhibitor is not recommended.

Cemiplimab-rwlc

Cemiplimab-rwlc is a recombinant human immunoglobulin G4 monoclonal antibody that inhibits PD-1 receptors, which improves antitumor immune response.³⁶³ Immune-mediated adverse events may occur with ICIs, including cemiplimab-rwlc.³⁶³ For patients with immune-mediated adverse events, intravenous high-dose corticosteroids should be administered based on the severity of the reaction. Cemiplimab-rwlc should also be permanently discontinued for patients with grades 2, 3, or 4 myocarditis and should be discontinued for other severe or life-threatening immune-mediated adverse events when indicated (see prescribing information).

EMPOWER-Lung 1, a phase 3 randomized trial, compared first-line therapy with single-agent cemiplimab versus platinum-based chemotherapy for patients with metastatic squamous or nonsquamous NSCLC, PD-L1 levels of 50% or more, and negative test results for *EGFR* mutations, *ALK* rearrangements, or *ROS1* rearrangements.³⁶³ Patients receiving first-line cemiplimab also received maintenance therapy with cemiplimab. Median overall survival was not reached for patients receiving cemiplimab (95% CI, 17.9–not evaluable) versus 14.2 months (95% CI, 11.2–17.5) for chemotherapy (HR, 0.57; 95% CI, 0.42–0.77; $P=0.0002$). The response rate was 39% (95% CI, 34%–45%) for those receiving cemiplimab versus 20% (95% CI, 16%–26%) for chemotherapy. Of patients treated with cemiplimab, 28% (98/355) had grade 3 to 4 adverse events compared with 39% (135/342) for those treated with chemotherapy. Treated-related deaths occurred in 2.5% (9/355) of patients receiving cemiplimab versus 2.0% (7/342) with chemotherapy. The cemiplimab-related deaths were due to autoimmune myocarditis, cardiac failure, cardiopulmonary failure, cardiorespiratory arrest, nephritis,

respiratory failure, septic shock, tumor hyperprogression, and unknown reasons.

The NCCN NSCLC Panel recommends cemiplimab-rwlc monotherapy (category 1; preferred) as a first-line therapy option for eligible patients with metastatic NSCLC regardless of histology, PD-L1 levels of 50% or more, and negative test results for actionable driver mutations based on clinical trial data and FDA approval; maintenance therapy with cemiplimab-rwlc is also recommended.³⁶³ Atezolizumab and pembrolizumab are also recommended first-line therapy options in this setting (category 1; preferred). The NCCN NSCLC Panel has preference stratified the ICI regimens and decided that atezolizumab, cemiplimab-rwlc, and pembrolizumab (all are category 1) are preferred first-line therapy options for eligible patients with metastatic NSCLC based on clinical trial data.^{141,363,944,953}

EMPOWER-Lung 3, a randomized phase 3 trial, assessed first-line therapy with cemiplimab plus platinum-based chemotherapy versus platinum-based chemotherapy alone in 466 patients with advanced NSCLC, regardless of PD-L1 levels or histology. The chemotherapy regimens included 1) cemiplimab plus paclitaxel plus either cisplatin or carboplatin for either nonsquamous NSCLC or squamous cell histology; or 2) cemiplimab plus pemetrexed plus either cisplatin or carboplatin for nonsquamous NSCLC.⁹⁵⁹ Patients received maintenance therapy if indicated. Most patients (85%) had stage IV NSCLC and 57% had nonsquamous NSCLC. Most patients either currently smoked cigarettes (55%) or had previously smoked (31%). Patients did not have *EGFR*, *ALK*, or *ROS1* mutations. The median overall survival was 21.9 months (95% CI, 15.5–not evaluable) with cemiplimab plus chemotherapy versus 13.0 months (95% CI, 11.9–16.1) with chemotherapy alone (HR, 0.71; 95% CI, 0.53–0.93; $P=0.01$). Subgroup analysis showed that overall survival was similar between the groups for PD-L1 levels less than 1%, nonsquamous



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

NSCLC, and other subgroups; however, the trial was not powered to detect differences in efficacy within the subgroups. Grade 3 or higher treatment-related adverse events occurred in 44% (136/312) of patients receiving cemiplimab plus chemotherapy, and 31% (48/153) of patients receiving chemotherapy alone. Immune-related adverse events occurred in 19% of patients receiving cemiplimab plus chemotherapy. Death occurred in 6.1% (19/312) of patients receiving cemiplimab plus chemotherapy versus 7.8% (12/153) of patients receiving chemotherapy alone. One patient died from immune-mediated pneumonitis.

The NCCN NSCLC Panel recommends cemiplimab-rwlc plus platinum-based chemotherapy regimens as first-line therapy options (category 1; other recommended) for eligible patients with metastatic NSCLC—regardless of histology or PD-L1 levels and with negative test results for actionable driver mutations—based on clinical trial data and FDA approval; maintenance therapy with cemiplimab-rwlc with or without pemetrexed is also recommended.⁹⁵⁹ Combination therapy with cemiplimab-rwlc plus platinum-based chemotherapy may be used as a first-line or subsequent therapy option, although it is not a preferred option, in eligible patients with actionable driver mutations as shown in the algorithm. Targeted therapy is a preferred first-line therapy option for patients with actionable driver mutations, except for *EGFR* exon 20 mutations or *KRAS* mutations. The NCCN Panel has preference stratified the cemiplimab-rwlc plus chemotherapy regimens and decided that they are other recommended first-line therapy options for eligible patients with metastatic NSCLC based on clinical trial data.⁹⁵⁹

Nivolumab with or Without Ipilimumab

Nivolumab and ipilimumab are ICIs that have complementary mechanisms of action on T cells; nivolumab is used either with or without ipilimumab, depending on the setting. Nivolumab is a human immunoglobulin G4 monoclonal antibody that inhibits PD-1 receptors, which improves

antitumor immune response.^{360,364,141} PD-1 receptors are expressed on activated cytotoxic T cells.³⁶⁰⁻³⁶² Ipilimumab is a human cytotoxic T-lymphocyte antigen 4 (CTLA-4)–blocking antibody that binds to CTLA-4 and prevents the interactions with CD80/CD86, which induces de novo T-cell responses against tumors; CTLA-4 inhibits T-cell activation.⁹⁶⁰ Immune-mediated adverse events may occur with ICIs, including nivolumab or nivolumab plus ipilimumab.³⁸¹ For patients with immune-mediated adverse events, intravenous high-dose corticosteroids should be administered based on the severity of the reaction (see the NCCN Guidelines for the Management of Immunotherapy-Related Toxicities, available at www.NCCN.org). Nivolumab either with or without ipilimumab should also be permanently discontinued for patients with severe or life-threatening pneumonitis or myocarditis and should be discontinued for other severe or life-threatening immune-mediated adverse events when indicated (see prescribing information). If patients are receiving nivolumab plus ipilimumab and have treatment-related adverse events, it may be reasonable to discontinue ipilimumab and continue nivolumab.³⁸¹

First-Line Therapy

CheckMate 227, a phase 3 randomized trial with a complex design, compared first-line therapy with nivolumab plus ipilimumab, nivolumab monotherapy, or chemotherapy for patients with metastatic nonsquamous or squamous NSCLC who had PD-L1 expression levels of 1% or more, PS 0 to 1, and no *EGFR* mutations or *ALK* rearrangements. First-line therapy with nivolumab plus ipilimumab, nivolumab plus chemotherapy, or chemotherapy alone was also compared for patients with PD-L1 expression levels less than 1%. In addition, first-line therapy with nivolumab plus ipilimumab or chemotherapy was compared as one of the co-primary analyses in patients who had high TMB levels (≥ 10 mutations/megabase).³⁸⁰ Preliminary data for PFS from CHECKMATE 227 suggested that TMB might be a useful immune biomarker for deciding



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

whether to use immunotherapy in patients with metastatic NSCLC.³⁸⁰ However, updated data from CHECKMATE 227 showed that overall survival was improved with nivolumab plus ipilimumab regardless of TMB or PD-L1 expression levels.³⁸¹

The PFS rate at 1 year was 42.6% for nivolumab plus ipilimumab versus 13.2% for chemotherapy alone. The median PFS for nivolumab plus ipilimumab was 7.2 months (95% CI, 5.5–13.2) compared with 5.5 months for chemotherapy alone (95% CI, 4.4–5.8) (HR for disease progression or death, 0.58; 97.5% CI, 0.41–0.81; $P < .001$). The objective response rate for nivolumab plus ipilimumab was 45.3% versus 26.9% with chemotherapy alone; nivolumab plus ipilimumab was beneficial regardless of PD-L1 expression levels or histology. The rate of grade 3 or 4 adverse events was similar for nivolumab plus ipilimumab versus chemotherapy alone (31% vs. 36%). The median PFS was not significantly different when comparing nivolumab monotherapy (N = 71) (4.2 months; 95% CI, 2.7–8.3) versus chemotherapy (N = 79) (5.6 months; 95% CI, 4.5–7.0). Updated results from CheckMate 227 for patients with PD-L1 expression of 1% or more, reported that the median overall survival was 17.1 months (95% CI, 15.0–20.1) for nivolumab plus ipilimumab versus 14.9 months (95% CI, 12.7–16.7) for chemotherapy (HR, 0.79; 95% CI, 0.65–0.96; $P = .007$).³⁸¹ For patients with PD-L1 levels of 1% or more, the 4-year overall survival rate was 29% with nivolumab plus ipilimumab versus 18% for those receiving chemotherapy (HR, 0.76; 95% CI, 0.65–0.90).⁹⁶¹ For patients with PD-L1 levels less than 1%, the 4-year overall survival rate was 24% with nivolumab plus ipilimumab versus 10% for those receiving chemotherapy (HR, 0.64; 95% CI, 0.51–0.81).

The NCCN NSCLC Panel recommends nivolumab plus ipilimumab as a first-line therapy option for eligible patients with metastatic NSCLC based on clinical trial data and FDA approval.^{380,381,383} Nivolumab plus ipilimumab is recommended for patients with metastatic NSCLC, regardless of PD-L1

levels or histology, negative test results for actionable driver mutations (ie, *ALK*, *BRAF* p.V600E, *EGFR*, *MET*ex14 skipping, *NTRK1/2/3*, *RET*, *ROS1*), and no contraindications to immunotherapy. For patients with PD-L1 levels of 1% or more, the NCCN recommendation is category 1 for nivolumab plus ipilimumab. Nivolumab plus ipilimumab is also a recommended option for patients with PD-L1 levels less than 1%. Maintenance therapy with nivolumab plus ipilimumab is also recommended. Combination therapy with nivolumab plus ipilimumab may be used as a first-line or subsequent therapy option, although it is not a preferred option, in eligible patients with actionable driver mutations as shown in the algorithm. Targeted therapy is a preferred first-line therapy option for patients with actionable driver mutations, except for *EGFR* exon 20 mutations or *KRAS* mutations.

The NCCN NSCLC Panel has preference stratified the systemic therapy regimens and decided that first-line therapy with nivolumab plus ipilimumab is “useful in certain circumstances” (eg, renal impairment) for patients with PD-L1 levels of 50% or more and is an “other recommended” first-line therapy option for patients with PD-L1 levels less than 50%. For the 2023 update (Version 1), the NCCN Panel revised the preference stratification for first-line therapy with nivolumab plus ipilimumab to an “other recommended” option from useful in certain circumstances in patients with PD-L1 levels of 1% to 49%. Previously, the NCCN Panel deleted TMB as an emerging immune biomarker based on clinical trial data and other issues (see *TMB* in this Discussion).¹⁷³ The NCCN Guidelines do not recommend measurement of TMB levels before deciding whether to use nivolumab plus ipilimumab regimens or to use other ICIs, such as pembrolizumab.¹⁷³

CheckMate 9LA, a phase 3 randomized trial, compared first-line nivolumab plus ipilimumab and 2 cycles of platinum-doublet chemotherapy versus 4 cycles of chemotherapy alone in patients with metastatic



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

nonsquamous or squamous NSCLC, regardless of PD-L1 expression levels, who had PS 0 to 1 and no *EGFR* mutations or *ALK* rearrangements.⁹⁶² For metastatic nonsquamous NSCLC, the chemotherapy was pemetrexed with either cisplatin or carboplatin; for metastatic squamous NSCLC, the chemotherapy was paclitaxel with carboplatin. Updated data show that the median overall survival with nivolumab plus ipilimumab plus chemotherapy was 15.8 months (95% CI, 13.9–19.7) versus 11.0 months (95% CI, 9.5–12.7) with chemotherapy alone regardless of histology or PD-L1 expression levels (HR, 0.72; 95% CI, 0.61–0.86).⁹⁶³ In patients receiving nivolumab plus ipilimumab plus chemotherapy, the 2-year overall survival rate was 38% versus 26% in those receiving chemotherapy alone. Based on histology or PD-1 levels, overall survival was also significantly different in patients receiving nivolumab plus chemotherapy compared with chemotherapy alone. The overall response rate was 38% with nivolumab plus ipilimumab plus chemotherapy versus 25% with chemotherapy alone.

Serious grade 3 or 4 adverse events occurred in 25.4% of patients receiving nivolumab plus ipilimumab plus chemotherapy versus 15% in those receiving chemotherapy alone. The death rate was 2% in each arm (nivolumab plus ipilimumab plus chemotherapy: 7/358; chemotherapy alone: 6/349). In the nivolumab plus ipilimumab plus chemotherapy arm, treatment-related deaths were from acute renal failure due to chemotherapy, thrombocytopenia, pneumonitis, hepatic toxicity, hepatitis, diarrhea, sepsis, and acute renal insufficiency; treatment-related deaths in the chemotherapy arm were from anemia, pancytopenia, febrile neutropenia, respiratory failure, pulmonary sepsis, and sepsis. The most common treatment-related adverse events (≥15%) were nausea, anemia, asthenia, and diarrhea.

The NCCN NSCLC Panel recommends nivolumab plus ipilimumab plus chemotherapy (category 1; other recommended) as first-line therapy

options for eligible patients with metastatic NSCLC based on clinical trial data and FDA approval.^{962,964} For metastatic nonsquamous NSCLC, the recommended chemotherapy is pemetrexed with either cisplatin or carboplatin; for metastatic squamous NSCLC, the recommended chemotherapy is paclitaxel with carboplatin. Nivolumab plus ipilimumab plus chemotherapy is recommended for patients with metastatic NSCLC, regardless of PD-L1 levels; negative test results for actionable driver mutations; and no contraindications to PD-1/PD-L1 inhibitors. Maintenance therapy with nivolumab plus ipilimumab is also recommended. Combination therapy with nivolumab plus ipilimumab may be used as a first-line or subsequent therapy option, although it is not a preferred option, in eligible patients with actionable driver mutations as shown in the algorithm. Targeted therapy is a preferred first-line therapy option for patients with actionable driver mutations, except for *EGFR* exon 20 mutations or *KRAS* mutations. The panel has preference stratified the systemic therapy regimens and voted that first-line therapy with nivolumab plus ipilimumab plus chemotherapy is an “other recommended” first-line therapy option for eligible patients with metastatic NSCLC regardless of PD-L1 expression levels or histology.

Subsequent Therapy

CheckMate-057, a phase 3 randomized trial, compared nivolumab versus docetaxel as subsequent therapy for patients with metastatic nonsquamous NSCLC and disease progression on or after first-line chemotherapy.³⁶⁰ Median overall survival was 12.2 months (95% CI, 9.7–15.0) for patients receiving nivolumab compared with 9.4 months (95% CI, 8.1–10.7) for docetaxel (HR, 0.73; 95% CI, 0.59–0.89; $P = .002$).³⁶⁰ The median duration of response was 17.2 months with nivolumab compared with 5.6 months for docetaxel. At 18 months, the overall survival rate was 39% (95% CI, 34%–45%) with nivolumab compared with 23% (95% CI, 19%–28%) for docetaxel. Fewer grade 3 to 5 adverse events were reported for nivolumab (10%) when compared with docetaxel (54%).



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

Although many patients with metastatic nonsquamous NSCLC benefit from nivolumab, those whose tumors have PD-L1 staining of 1% to 10% or more have an overall survival of 17 to 19 months compared with 8 to 9 months for docetaxel. For patients who did not have PD-L1 expression, there was no difference in overall survival for nivolumab versus docetaxel; however, nivolumab was associated with a longer duration of response and fewer side effects.

CheckMate-017, a phase 3 randomized trial, compared nivolumab versus docetaxel as subsequent therapy for patients with metastatic NSCLC squamous cell and disease progression on or after first-line chemotherapy.³⁶⁴ Median overall survival was 9.2 months (95% CI, 7.3–13.3) for nivolumab compared with 6.0 months (95% CI, 5.1–7.3) for docetaxel (HR, 0.59; 95% CI, 0.44–0.79; $P < .001$).³⁶⁴ Patients had a response rate of 20% with nivolumab compared with 9% for docetaxel ($P = .008$). PD-L1 expression was not associated with response to nivolumab in patients with squamous cell NSCLC. Fewer grade 3 to 4 adverse events were reported with nivolumab (7%) compared with docetaxel (55%). No patients died in the nivolumab arm versus three deaths in the docetaxel arm.

In a long-term analysis of CheckMate-057 and CheckMate-017, 2-year survival and durable responses were increased in patients with advanced NSCLC receiving nivolumab when compared with docetaxel.⁹⁶⁵ For patients with nonsquamous NSCLC, 2-year survival was 29% (95% CI, 24%–34%) with nivolumab versus 16% (95% CI, 12%–20%) with docetaxel. For those with squamous NSCLC, 2-year survival was 23% (95% CI, 16%–30%) with nivolumab versus 8% (95% CI, 4%–13%) with docetaxel. Fewer severe treatment-related adverse events were reported with nivolumab compared with docetaxel (grade 3–4, 10% vs. 55%). At 5 years, overall survival was 13.4% for patients receiving nivolumab versus 2.6% for those receiving docetaxel.²⁰

If patients have not previously received a PD-1/PD-L1 inhibitor, the NCCN NSCLC Panel recommends single-agent nivolumab (category 1; preferred) as a subsequent therapy option for patients with metastatic nonsquamous or squamous NSCLC and disease progression on or after first-line chemotherapy based on clinical trial data and FDA approval.^{360,364,965,966} Testing for PD-L1 is not required for prescribing nivolumab but may provide useful information.³⁷² The NCCN NSCLC Panel recommends nivolumab, atezolizumab, or pembrolizumab as preferred subsequent therapy options (category 1 for all) based on improved overall survival rates, longer duration of response, and fewer adverse events when compared with cytotoxic chemotherapy.^{360,364,946,967} If patients have progressed on a PD-1/PD-L1 inhibitor, then using a PD-1/PD-L1 inhibitor is not recommended.

Immune-related adverse events, such as pneumonitis, may occur with nivolumab.^{362,968-974} Intravenous high-dose corticosteroids should be administered for patients with immune-mediated adverse events based on the severity of the reaction (see the NCCN Guidelines for the Management of Immunotherapy-Related Toxicities, available at www.NCCN.org). Nivolumab should be discontinued for patients with severe or life-threatening pneumonitis or myocarditis and should be withheld or discontinued for other severe or life-threatening immune-mediated adverse events when indicated (see prescribing information).

Pembrolizumab

Pembrolizumab is a humanized immunoglobulin G4 monoclonal antibody that inhibits PD-1 receptors, which improves antitumor immune response.^{364,141} The FDA has approved a companion diagnostic biomarker test for assessing PD-L1 expression and determining which patients are eligible for pembrolizumab therapy. Although it is not an optimal biomarker, PD-L1 expression is currently the best available biomarker to assess whether patients are candidates for



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

pembrolizumab.^{368,369} PD-L1 expression is continuously variable and dynamic; thus, a cutoff value for a positive result is artificial. Patients with PD-L1 expression levels just below and just above 50% will probably have similar responses.³⁶⁸ Unique anti-PD-L1 IHC assays have been developed for each one of the different ICIs currently available.^{368,372} The definition of a positive PD-L1 test result varies depending on which biomarker assay is used.³⁷²

Immune-mediated adverse events may occur with ICIs, including pembrolizumab.^{968,970,975} For patients with immune-mediated adverse events, intravenous high-dose corticosteroids should be administered based on the severity of the reaction (see the NCCN Guidelines for the Management of Immunotherapy-Related Toxicities, available at www.NCCN.org). Pembrolizumab should also be discontinued for patients with severe or life-threatening pneumonitis or myocarditis and should be withheld or discontinued for other severe or life-threatening immune-mediated adverse events when indicated (see prescribing information). The following sections describe the use of pembrolizumab in eligible patients with metastatic NSCLC. The use of adjuvant pembrolizumab following adjuvant chemotherapy in eligible patients with completely resected NSCLC is described in a different section (see *Surgery Followed by Adjuvant Therapy: Trial Data and NCCN Recommendations* in this Discussion).³²

First-Line Monotherapy

KEYNOTE-024, a phase 3 randomized trial, compared single-agent pembrolizumab versus platinum-based chemotherapy as first-line therapy for patients with advanced nonsquamous or squamous NSCLC and PD-L1 expression levels of 50% or more, but without *EGFR* mutations or *ALK* rearrangements.^{12,15,141} At 6 months, the rate of overall survival was 80.2% with pembrolizumab monotherapy versus 72.4% with chemotherapy (HR for death, 0.60; 95% CI, 0.41–0.89; *P* = .005). Responses were higher for

pembrolizumab than for chemotherapy (44.8% vs. 27.8%).¹⁴¹ An updated analysis of KEYNOTE-024 showed that median overall survival was 26.3 months (95% CI, 18.3 months–40.4) with pembrolizumab monotherapy compared with 13.4 months (95% CI, 9.4–18.3 months) with chemotherapy (HR, 0.62; 95% CI, 0.48–0.81).¹⁵ The 5-year overall survival rate was 31.9% for pembrolizumab and 16.3% for chemotherapy. Fewer severe treatment-related adverse events (grades 3–5) were reported in patients receiving pembrolizumab monotherapy compared with those receiving chemotherapy (31.2% vs. 53.3%).¹² Treatment-related deaths occurred in 1.3% (2/154) of patients receiving pembrolizumab monotherapy versus 2% (3/150) of patients receiving chemotherapy alone.

KEYNOTE-042, a phase 3 randomized trial, compared single-agent pembrolizumab versus platinum-based chemotherapy as first-line therapy for patients with advanced nonsquamous or squamous NSCLC and PD-L1 expression levels of 1% or more, but without *EGFR* mutations or *ALK* rearrangements.⁹⁴⁴ Overall survival was 20.0 months (95% CI, 15.4–24.9) in patients with PD-L1 levels of 50% or more who received single-agent pembrolizumab compared with 12.2 months (95% CI, 10.4–14.2) with chemotherapy (HR, 0.69; 95% CI, 0.56–0.85; *P* = .0003). In a subgroup analysis, overall survival was similar in patients with PD-L1 levels of 1% to 49% who received single-agent pembrolizumab (13.4 months; 95% CI, 10.7–18.2) compared with chemotherapy (12.1 months; 95% CI, 11.0–14.0) (HR, 0.92; 95% CI, 0.77–1.11).

Long-term data from KEYNOTE-001 show that 5-year survival for patients with metastatic NSCLC is approximately 23% for patients who received first-line pembrolizumab monotherapy; for patients with PD-L1 levels of 50% or more, 5-year overall survival is about 29.6%.²² Median overall survival was 22.3 months (95% CI, 17.1–32.3) for



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

treatment-naïve patients. For patients with metastatic NSCLC receiving chemotherapy alone, 5-year overall survival is approximately 6%.²²

The NCCN NSCLC Panel recommends IHC testing for PD-L1 expression (category 1) before first-line treatment in all patients with metastatic NSCLC based on the efficacy of pembrolizumab (see *Principles of Molecular and Biomarker Analysis* in the NCCN Guidelines for NSCLC).³⁶⁷

The panel recommends single-agent pembrolizumab (category 1; preferred) as a first-line therapy option for eligible patients with advanced nonsquamous or squamous NSCLC, PD-L1 expression levels of 50% or more, no contraindications to PD-1 or PD-L1 inhibitors, and negative test results for actionable driver mutations based on clinical trial data and FDA approval.^{141,944,976} Maintenance therapy with pembrolizumab is also a recommended option in this setting (category 1). The NCCN NSCLC Panel has preference stratified the ICI regimens and decided that pembrolizumab, atezolizumab, and cemiplimab-rwlc (all are category 1) are all preferred single-agent options for eligible patients with metastatic NSCLC.^{141,363,944,953} For patients who have disease progression on first-line therapy with single-agent pembrolizumab, subsequent therapy with initial cytotoxic systemic therapy regimens (eg, carboplatin plus paclitaxel) is recommended by the NCCN NSCLC Panel.

The NCCN NSCLC Panel also recommends single-agent pembrolizumab as a first-line therapy option (category 2B; useful in certain circumstances) for eligible patients with metastatic NSCLC, PD-L1 expression levels of 1% to 49%, no contraindications to PD-1 or PD-L1 inhibitors, and negative test results for actionable driver mutations based on clinical trial data and FDA approval.^{944,976} The NCCN NSCLC Panel decided that single-agent pembrolizumab is a useful intervention in patients with PD-L1 levels of 1% to 49% when there are contradictions to combination chemotherapy (category 2B; useful in certain circumstances). In patients with PD-L1 levels of 1% to 49%, the HR of

0.92 is not statistically or clinically significant for pembrolizumab monotherapy versus chemotherapy; therefore, pembrolizumab plus chemotherapy is recommended (category 1; preferred) if patients can tolerate the therapy.

First-Line Combination Therapy

KEYNOTE-189, a phase 3 randomized trial, compared pembrolizumab added to carboplatin (or cisplatin) plus pemetrexed versus chemotherapy in patients with metastatic nonsquamous NSCLC.⁷¹² Most patients received pembrolizumab plus carboplatin plus pemetrexed (72% [445/616]) in this trial, but some received pembrolizumab plus cisplatin plus pemetrexed (28% [171/616]); patients did not have *EGFR* mutations or *ALK* rearrangements. The estimated rate of overall survival at one year was 69.2% (95% CI, 64.1%–73.8%) in patients receiving pembrolizumab plus chemotherapy versus 49.4% (95% CI, 42.1%–56.2%) for chemotherapy alone (HR for death, 0.49; 95% CI, 0.38–0.64; $P < .001$) after a median follow-up of 10.5 months. Overall survival was improved regardless of PD-L1 expression levels; TMB did not predict for response.⁹⁷⁷ For the pembrolizumab plus chemotherapy group, median PFS was 8.8 months (95% CI, 7.6–9.2) compared with 4.9 months (95% CI, 4.7–5.5) for chemotherapy alone (HR for disease progression or death, 0.52; 95% CI, 0.43–0.64; $P < .001$). Grade 3 or higher adverse events occurred at a similar rate in both arms (pembrolizumab plus chemotherapy, 67.2% vs. chemotherapy, 65.8%).

The NCCN NSCLC Panel recommends pembrolizumab plus pemetrexed and either carboplatin or cisplatin (category 1; preferred) as a first-line therapy option for eligible patients with metastatic nonsquamous NSCLC (ie, adenocarcinoma, large cell carcinoma) or NSCLC NOS based on



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

clinical trial data and FDA approval.^{712,713} The NCCN NSCLC Panel has preference stratified the systemic therapy regimens and decided that these pembrolizumab plus chemotherapy regimens are preferred first-line options for eligible patients with metastatic nonsquamous NSCLC. These pembrolizumab plus chemotherapy regimens are recommended (category

1; preferred) as first-line therapy options for patients with metastatic nonsquamous NSCLC, no contraindications to PD-1 or PD-L1 inhibitors, and negative test results for actionable driver mutations, regardless of their PD-L1 expression levels. Maintenance therapy with pembrolizumab plus pemetrexed is also a recommended option (category 1) in this setting. Combination therapy with pembrolizumab plus pemetrexed and either carboplatin or cisplatin may be used as a first-line or subsequent therapy option in eligible patients with actionable driver mutations as shown in the algorithm. However, targeted therapy is a preferred first-line therapy option for patients with actionable driver mutations, except for *EGFR* exon 20 mutations or *KRAS* mutations. For patients with metastatic NSCLC and disease progression on combination therapy with PD-1/PD-L1 inhibitors plus chemotherapy, subsequent therapy with docetaxel (with or without ramucirumab), pemetrexed (nonsquamous only), albumin-bound paclitaxel, or gemcitabine is recommended if not previously given.

KEYNOTE-407, a phase 3 randomized trial, compared pembrolizumab added to carboplatin and either paclitaxel or albumin-bound paclitaxel in patients with metastatic squamous cell NSCLC; 32% of patients received albumin-bound paclitaxel (also known as nab-paclitaxel).⁹⁷⁸ Median overall survival was 15.9 months (95% CI, 13.2–not reached) with pembrolizumab plus chemotherapy versus 11.3 months (95% CI, 9.5–14.8) with chemotherapy alone (HR for death, 0.64; 95% CI, 0.49–0.85; $P < .001$). Patients receiving pembrolizumab plus chemotherapy had an overall response rate of 57.9% compared to 38.4% for those receiving

chemotherapy alone. Only 38% of patients had a PD-L1 tumor proportion score (TPS) less than 1%. Grade 3 or higher adverse events were similar in both groups (pembrolizumab plus chemotherapy, 69.8% vs. chemotherapy alone, 68.2%). Because of adverse events, more patients discontinued treatment with pembrolizumab plus chemotherapy than with chemotherapy (13.3% vs. 6.4%, respectively). A pooled analysis of three randomized trials (ie, KEYNOTE-189, KEYNOTE-407, KEYNOTE-021) in patients with metastatic NSCLC and PD-L1 levels less than 1% showed that overall survival was improved in those receiving pembrolizumab plus chemotherapy versus chemotherapy alone (HR, 0.63; 95% CI, 0.50–0.79).⁹⁷⁹

The NCCN NSCLC Panel recommends pembrolizumab plus carboplatin and either paclitaxel or albumin-bound paclitaxel (category 1; preferred) as a first-line therapy option for patients with metastatic squamous cell NSCLC based on clinical trial data and FDA approval.^{978,980} Maintenance therapy with pembrolizumab is also a recommended option in this setting (category 1). The NCCN NSCLC Panel has preference stratified the systemic therapy regimens and decided that these pembrolizumab plus chemotherapy regimens are preferred for eligible patients with metastatic squamous cell NSCLC. These pembrolizumab plus chemotherapy regimens are recommended (category 1; preferred) as first-line therapy options for patients with metastatic squamous cell NSCLC, no contraindications to PD-1 or PD-L1 inhibitors, and negative test results for actionable driver mutations, regardless of their PD-L1 expression levels. The panel does not recommend pembrolizumab plus cisplatin with either paclitaxel or albumin-bound paclitaxel, because there are fewer data for this regimen. Combination therapy with pembrolizumab plus carboplatin and either paclitaxel or albumin-bound paclitaxel may be used as a first-line or subsequent therapy option in eligible patients with actionable driver mutations as shown in the algorithm. However, targeted therapy is



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

a preferred first-line therapy option for patients with actionable driver mutations, except for *EGFR* exon 20 mutations or *KRAS* mutations.

Subsequent Therapy

KEYNOTE-010, a phase 3 randomized trial, assessed single-agent pembrolizumab in patients with previously treated advanced nonsquamous and squamous NSCLC who were PD-L1 positive ($\geq 1\%$); most patients either currently or previously smoked cigarettes.⁹⁴⁶ There were three arms in this trial: pembrolizumab at 2 mg/kg, pembrolizumab at 10 mg/kg, and docetaxel at 75 mg/m² every 3 weeks. The median overall survival was 10.4 months for the lower dose of pembrolizumab, 12.7 months for the higher dose, and 8.5 months for docetaxel. Overall survival was significantly longer for both doses of pembrolizumab versus docetaxel (pembrolizumab 2 mg/kg: HR, 0.71; 95% CI, 0.58–0.88; $P = .0008$) (pembrolizumab 10 mg/kg: HR, 0.61; CI, 0.49–0.75; $P < .0001$). For those patients with at least 50% PD-L1 expression in tumor cells, overall survival was also significantly longer at either dose of pembrolizumab when compared with docetaxel (pembrolizumab 2 mg/kg: 14.9 vs. 8.2 months; HR, 0.54; 95% CI, 0.38–0.77; $P = .0002$) (pembrolizumab 10 mg/kg: 17.3 vs. 8.2 months; HR, 0.50; 95% CI, 0.36–0.70; $P < .0001$). When compared with docetaxel, there were fewer grade 3 to 5 treatment-related adverse events at either dose of pembrolizumab (pembrolizumab 2 mg/kg: 13% [43/339] of patients, pembrolizumab 10 mg/kg: 16% [55/343] of patients; and docetaxel: 35% [109/309] of patients). A total of six treatment-related deaths occurred in patients receiving pembrolizumab (three at each dose) and five treatment-related deaths occurred in the docetaxel arm. Long-term data from KEYNOTE-010 show that 5-year survival for patients with metastatic NSCLC is approximately 15.6% for patients who received subsequent pembrolizumab monotherapy versus 6.5% for those receiving docetaxel. For patients with PD-L1 levels of 50% or more, 5-year overall survival was about 25% for patients receiving

pembrolizumab versus 8.2% for those receiving docetaxel, respectively.¹⁴

If patients have not previously received a PD-1/PD-L1 inhibitor, the NCCN NSCLC Panel recommends single-agent pembrolizumab (category 1; preferred) as a subsequent therapy option for patients with metastatic nonsquamous or squamous NSCLC and PD-L1 expression levels of 1% or more based on clinical trial data and FDA approval.^{946,981,982} Nivolumab and atezolizumab are also recommended subsequent therapy options in this setting (category 1; preferred). Testing for PD-L1 expression levels is recommended before prescribing pembrolizumab monotherapy (see *Principles of Molecular and Biomarker Analysis* in the NCCN Guidelines for NSCLC).

Tremelimumab-actl and Durvalumab

Tremelimumab-actl is a monoclonal IgG2 antibody that enhances immune function by inhibiting CTLA-4 and preventing interactions with CD80/CD86; CTLA-4 inhibits T-cell activation.⁹⁸³ Durvalumab is a human immunoglobulin G1 monoclonal antibody that inhibits PD-L1 binding to PD-1 and CD80, thus increasing immune function (see *Durvalumab* in this Discussion). Immune-mediated adverse events may occur with ICIs, including tremelimumab-actl and durvalumab.⁹⁸³ For patients with immune-mediated adverse events, intravenous high-dose corticosteroids should be administered based on the severity of the reaction. Tremelimumab-actl and durvalumab should be permanently discontinued for patients with grades 2, 3, or 4 myocarditis and should be discontinued for other severe or life-threatening immune-mediated adverse events when indicated (see prescribing information).

POSEIDON, a phase 3 randomized trial, assessed three different first-line treatment arms in 1013 patients with metastatic NSCLC who did not have *EGFR* or *ALK* mutations.⁹⁸⁴ The treatment arms included 1) tremelimumab plus durvalumab plus platinum-based chemotherapy; 2) durvalumab plus



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

platinum-based chemotherapy; and 3) platinum-based chemotherapy alone; there were 675 patients in arms 1 and 3. The chemotherapy regimens included 1) tremelimumab plus durvalumab plus albumin-bound paclitaxel plus carboplatin for either nonsquamous NSCLC or squamous cell histology; 2) tremelimumab plus durvalumab plus pemetrexed plus either cisplatin or carboplatin for nonsquamous NSCLC; or 3) tremelimumab plus durvalumab plus gemcitabine plus either cisplatin or carboplatin for squamous cell histology. Most patients received the pemetrexed regimen. Most patients either currently smoked cigarettes (25%) or had previously smoked (57%). The median overall survival was 14 months (95% CI, 11.7–16.1) in patients receiving tremelimumab plus durvalumab plus chemotherapy versus 11.7 months (95% CI, 10.5–13.1) in patients receiving chemotherapy alone (HR, 0.77; 95% CI, 0.65–0.92; $P=.003$). Subgroup analysis showed that overall survival was similar between the groups for squamous cell histology and other subgroups.

In patients receiving tremelimumab plus durvalumab plus chemotherapy, grade 3 or 4 adverse events included neutropenia, anemia, leukopenia, lymphocytopenia, increased lipase levels, hyponatremia, and thrombocytopenia.⁹⁸⁴ Common serious adverse events in patients receiving tremelimumab plus durvalumab plus chemotherapy included pneumonia (11%), anemia (5%), diarrhea (2.4%), thrombocytopenia (2.4%), pyrexia (2.4%), and febrile neutropenia (2.1%). Grade 3 to 4 immune-related adverse events occurred in 10% of patients receiving tremelimumab plus durvalumab plus chemotherapy versus 1.5% of those receiving chemotherapy alone. Treatment-related deaths occurred in 3.3% (11/330) of patients receiving tremelimumab plus durvalumab plus chemotherapy and 2.4% (8/333) receiving chemotherapy alone. An exploratory analysis suggests that tremelimumab plus durvalumab plus chemotherapy may be beneficial in patients with metastatic NSCLC who have *STK11*, *KEAP1*, or *KRAS* mutations; however, the sample size was too small to yield definitive conclusions.⁹⁸⁵

The NCCN NSCLC Panel recommends tremelimumab-actl plus durvalumab plus platinum-based chemotherapy regimens as first-line therapy options for eligible patients with metastatic NSCLC—regardless of histology or PD-L1 levels, and with negative test results for actionable driver mutations—based on clinical trial data and FDA approval; maintenance therapy with durvalumab with or without pemetrexed is also recommended.⁹⁸⁴ Combination therapy with tremelimumab-actl plus durvalumab plus platinum-based chemotherapy may be used as a first-line or subsequent therapy option, although it is not a preferred option, in eligible patients with actionable driver mutations as shown in the algorithm. Targeted therapy is a preferred first-line therapy option for patients with actionable driver mutations, except for *EGFR* exon 20 mutations or *KRAS* mutations. The category assignment (ie, category 1, 2A, or 2B) for the tremelimumab-actl plus durvalumab plus platinum-based chemotherapy regimens varies depending on the histology, chemotherapy, and PD-L1 levels. The NCCN Panel has preference stratified the tremelimumab-actl plus durvalumab plus chemotherapy regimens and decided that they are other recommended first-line therapy options for eligible patients with metastatic NSCLC based on clinical trial data.⁹⁸⁴

Maintenance Therapy

Maintenance therapy refers to therapy given for patients with advanced NSCLC after 4 to 6 cycles of first-line therapy.⁹⁸⁶ Patients are only candidates for maintenance therapy if their tumors have responded to their previous treatment or they have stable disease and their tumors have not progressed. *Continuation maintenance* therapy refers to the use of at least one of the agents that was given in the first-line regimen. *Switch maintenance* therapy refers to the initiation of a different agent that was not included as part of the first-line regimen. Selection of appropriate maintenance therapy depends on several factors, such as histologic type, presence of mutations or gene fusions, and PS. Maintenance therapy is



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

recommended in the NCCN Guidelines for select patients with tumor response or stable disease and is not recommended for all patients; it is not recommended for patients with PS 3 to 4 or those with progression (see the NCCN Guidelines for NSCLC).⁹⁸⁷ Monitoring is recommended during maintenance therapy with response assessment with CT, with or without contrast, of known or high-risk sites of disease every 6 to 12 weeks.

Continuation Maintenance Therapy

For continuation maintenance therapy, select agents (which were initially given with first-line therapy) may be continued until evidence of disease progression or unacceptable toxicity based on the design of the clinical trials that led to their approval. This section mainly discusses continuation maintenance with chemotherapy; continuation maintenance with ICIs is discussed in another section (see *Immune Checkpoint Inhibitors* in this Discussion). Use of continuation maintenance therapy depends on several factors, such as whether the patient had minimal toxicity during treatment. A drug vacation may be more appropriate for some patients.⁸¹⁰ Some clinicians feel that continuation maintenance therapy is only appropriate for select patients, because it has only been shown to improve overall survival or quality of life for a few agents and not all agents, although it has been shown to improve PFS.^{808,810} In addition, maintenance therapy has not been shown to be superior to subsequent therapy, which is initiated at disease progression. A phase 3 randomized trial suggests that conventional cytotoxic agents should not be continued beyond 4 to 6 cycles of therapy; however, many patients assigned to a longer duration of therapy did not receive the planned number of cycles (see *Maintenance Therapy* in this Discussion).^{807,808}

PARAMOUNT, a phase 3 randomized trial, reported that continuation maintenance therapy with pemetrexed slightly increased PFS when compared with placebo (4.1 vs. 2.8 months).⁹⁸⁸ Updated results from

PARAMOUNT reported that continuation maintenance therapy with pemetrexed also improves overall survival (13.9 vs. 11.0 months).⁹⁸⁹ The NCCN NSCLC Panel recommends single-agent pemetrexed as continuation maintenance therapy (category 1) in patients with nonsquamous NSCLC based on clinical trial data and FDA approval.⁹⁸⁸⁻⁹⁹⁰

POINTBREAK, a phase 3 randomized trial, assessed bevacizumab plus carboplatin plus pemetrexed compared with bevacizumab plus carboplatin plus paclitaxel in patients with metastatic NSCLC; patients received maintenance therapy with either bevacizumab plus pemetrexed or bevacizumab alone.⁸¹² PFS was 6 months with pemetrexed plus carboplatin plus bevacizumab versus 5.6 months with paclitaxel plus carboplatin plus bevacizumab.⁸¹² It is important to note that the pemetrexed-based arm was associated with less toxicity (eg, less neurotoxicity, less neutropenia, less hair loss) than the paclitaxel-based arm. AVAPERL, a phase 3 randomized trial, assessed maintenance therapy with bevacizumab plus pemetrexed versus bevacizumab alone in patients with advanced nonsquamous NSCLC; the initial regimen was bevacizumab plus cisplatin plus pemetrexed.^{991,992} An updated analysis reported that overall survival was 17.1 months with bevacizumab plus pemetrexed maintenance versus 13.2 months with bevacizumab alone (HR, 0.87; 95% CI, 0.63–1.21; $P = .29$).⁹⁹¹

The NCCN NSCLC Panel recommends continuation maintenance therapy with bevacizumab plus pemetrexed in patients with nonsquamous NSCLC who initially received bevacizumab plus pemetrexed plus platinum regimen based on clinical trial data.^{991,992} Single-agent bevacizumab (category 1) may be continued beyond 4 to 6 cycles of initial therapy (ie, platinum-doublet chemotherapy given with bevacizumab) in patients with nonsquamous NSCLC.^{806,990,993} The Panel recommends that bevacizumab biosimilars may be used in any of the systemic therapy regimens containing bevacizumab (eg, carboplatin plus paclitaxel plus



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

bevacizumab) that are used for eligible patients with metastatic NSCLC based on clinical data and FDA approvals.⁸⁹⁶⁻⁹⁰⁰ Therefore, if a bevacizumab biosimilar was initially used as part of first-line combination therapy, the biosimilar should be continued as maintenance therapy in eligible patients.

IFCT-GFPC 0502, a phase 3 randomized trial, compared using maintenance therapy with either gemcitabine or erlotinib after first-line therapy with cisplatin-gemcitabine in patients with advanced NSCLC. Continuation maintenance therapy with single-agent gemcitabine was reported to increase PFS to a greater extent (3.8 months) than switch maintenance therapy with erlotinib (2.9 months) when compared with observation (1.9 months).^{743,994} A phase 3 randomized trial from the ECOG assessed continuation maintenance therapy with gemcitabine versus best supportive care after an initial regimen of cisplatin plus gemcitabine.⁹⁹⁵ The data showed a slight difference in PFS but no difference in overall survival (13 vs. 11 months, respectively; $P = .195$). The NCCN NSCLC Panel recommends gemcitabine (category 2B) as continuation maintenance therapy regardless of histology in patients with metastatic NSCLC, negative test results for actionable driver mutations, and PD-L1 expression less than 1%.

Switch Maintenance Therapy

Issues have been raised about switch maintenance therapy, including the design of the trials, modest survival benefits, quality of life, and toxicity.^{810,996} Two phase 3 randomized trials reported a benefit in PFS and overall survival with the initiation of pemetrexed after first-line chemotherapy (4–6 cycles) in patients with nonsquamous NSCLC and no apparent disease progression.^{997,998} The NCCN NSCLC Panel recommends switch maintenance therapy with pemetrexed in eligible patients with nonsquamous cell carcinoma; negative test results for

actionable driver mutations, and PD-L1 expression less than 1% based on clinical trial data and FDA approval.^{998,999}

The NCCN NSCLC Panel does not recommend erlotinib as switch maintenance therapy (or as subsequent therapy) for patients with nonsquamous NSCLC, good PS, and negative test results for actionable driver mutations based on results from IUNO, a randomized trial, and a revised indication from the FDA.¹⁰⁰⁰ The NCCN NSCLC Panel also does not recommend switch maintenance therapy with erlotinib in patients with squamous cell NSCLC, because overall survival and quality of life were not improved.^{743,1001} A phase 3 trial assessed switch maintenance therapy with docetaxel given either immediately after gemcitabine plus carboplatin or delayed until progression in patients with advanced NSCLC; however, many patients in the delayed chemotherapy arm did not receive docetaxel.¹⁰⁰² Previously, the panel deleted the recommendation for switch maintenance therapy with docetaxel for patients with squamous cell NSCLC because there are better options.^{1002,1003}

Clinical Evaluation

The workup and evaluation of incidental lung nodules—that are detected on imaging for other conditions—are described in the NSCLC algorithm (see *Incidental Lung Nodules* in this Discussion and the NCCN Guidelines for NSCLC). The cutoff thresholds are 6 mm for a positive scan result for incidental solid and subsolid lung nodules detected on chest CT based on the Fleischner criteria (see the NCCN Guidelines for NSCLC).⁹⁴⁻⁹⁸ As previously described, low-dose CT screening is recommended for asymptomatic select patients who are at high risk for lung cancer and management of any nodules detected in these patients is described elsewhere (see the NCCN Guidelines for Lung Cancer Screening, available at www.NCCN.org).



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

After patients are confirmed to have NSCLC based on a pathologic diagnosis, a clinical evaluation needs to be done (see the NCCN Guidelines for NSCLC). In patients with symptoms, the clinical stage is initially determined from disease history (ie, cough, dyspnea, chest pain, weight loss) and physical examination together with a limited battery of tests (see *Initial Evaluation* and *Clinical Stage* in the NCCN Guidelines for NSCLC). The NCCN NSCLC Panel also recommends that smoking cessation advice, counseling, and pharmacotherapy be provided to patients.^{52,1004-1006} After the clinical stage is determined, the patient is assigned to one of the pathways that are defined by the stage, specific subdivision of the particular stage, and location of the tumor. Note that for some patients, diagnosis, staging, and surgical resection are done during the same operative procedure. A multidisciplinary evaluation should be done before treatment.

Additional Pretreatment Evaluation

As previously noted, evaluation of the mediastinal nodes is a key step in further staging of the patient. FDG PET/CT scans can be used as an initial assessment of the hilar and mediastinal nodes (ie, to determine whether the N1, N2, or N3 nodes are positive for cancer, which is a key determinant of stage II and stage III disease); however, CT scans have known limitations for evaluating the extent of lymph node involvement in lung cancer.^{112,1007-1009} When compared with noninvasive staging methods (EBUS, EUS), surgical staging with mediastinoscopy is more appropriate for certain settings when evaluating mediastinal nodes; however, clinicians use both methods when staging patients.¹¹² Thus, mediastinoscopy is encouraged as part of the initial evaluation, particularly if the results of imaging are not conclusive and the probability of mediastinal involvement is high (based on tumor size and location). Therefore, mediastinoscopy is appropriate for patients with T2 to T3 lesions even if the FDG PET/CT scan does not suggest mediastinal node involvement.

Mediastinoscopy may also be appropriate to confirm mediastinal node involvement in patients with a positive FDG PET/CT scan. Although mediastinal biopsy is generally preferred, the risks in selected patients may outweigh the benefits, such as patients who are medically inoperable. In patients with peripheral tumors (outer one third of the lung) less than 3 cm, pathologic mediastinal lymph node evaluation is optional if the nodes are FDG PET/CT negative because there is a low likelihood of positive mediastinal nodes.¹⁰¹⁰ Invasive mediastinal staging is recommended for central tumors. Mediastinal evaluation can be considered in patients with clinical stage IA disease (T1abc,N0). In patients with peripheral T2a, central T1abc, or T2a lesions with negative FDG PET/CT scans, the risk for mediastinal lymph node involvement is higher and mediastinoscopy and/or EUS-FNA and EBUS-TBNA are recommended. Dillemans et al have reported a selective mediastinoscopy strategy, proceeding straight to thoracotomy without mediastinoscopy for T1 peripheral tumors without enlarged mediastinal lymph nodes on preoperative CT.¹⁰¹¹ This strategy resulted in a 16% incidence of positive N2 nodes discovered only at the time of thoracotomy.

For identifying N2 disease, chest CT scans had sensitivity and specificity rates of 69% and 71%, respectively. Using a chest CT scan plus mediastinoscopy was significantly more accurate (89% vs. 71%) than using a chest CT scan alone for identifying N2 disease. When using CT scans, node positivity is based on the size of the lymph nodes. Therefore, a CT scan will miss small metastases that do not result in node enlargement. To address this issue, Arita et al specifically examined lung cancer metastases in normal size mediastinal lymph nodes in 90 patients and found an incidence of 16% (14/90) false-negative chest CT scans with histologic identification of occult N2 or N3 disease.¹⁰¹² Bronchoscopy is used in diagnosis and local staging of both central and peripheral lung lesions and is recommended for pretreatment evaluation of stage I to IIIA tumors. In patients who present with a solitary pulmonary nodule where



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

the suspicion of malignancy is high, surgical resection without prior invasive testing may be reasonable.

As previously mentioned, CT scans have known limitations for evaluating the extent of lymph node involvement in lung cancer.¹⁰⁰⁷ PET scans have been used to help evaluate the extent of disease and to provide more accurate staging. The NCCN NSCLC Panel reviewed the diagnostic performance of CT and PET scans. The panel assessed studies that examined the sensitivity and specificity of chest CT scans for mediastinal lymph node staging.¹⁰¹³ Depending on the clinical scenario, a sensitivity of 40% to 65% and a specificity of 45% to 90% were reported.¹⁰¹⁴ Because they detect tumor physiology, as opposed to anatomy, PET scans may be more sensitive than CT scans. Moreover, if postobstructive pneumonitis is present, there is little correlation between the size of the mediastinal lymph nodes and tumor involvement.¹⁰¹⁵ Chin et al found that PET, when used to stage the mediastinal nodes, was 78% sensitive and 81% specific with a negative predictive value of 89%.¹⁰¹⁶ Kernstine et al compared PET scan to CT scan for identifying N2 and N3 disease in NSCLC.¹⁰¹⁷ The PET scan was found to be more sensitive than the CT scan in identifying mediastinal node disease (70% vs. 65%). FDG PET/CT has been shown to be useful in restaging patients after adjuvant therapy.^{1018,1019} When patients with early-stage disease are accurately staged using FDG PET/CT, inappropriate surgery is avoided.¹⁰²⁰

The NCCN NSCLC Panel believes that PET scans can play a role in the evaluation and more accurate staging of NSCLC, for example, in identifying stage I (peripheral and central T1–2,N0), stage II, stage III, and stage IV diseases.^{112,1021,1022} However, FDG PET/CT is even more sensitive and is recommended by the panel.^{1020,1023,1024} PET/CT is typically done from the skull base to the knees; whole body PET/CT may also be done.

Positive FDG PET/CT scan findings for distant disease need pathologic or other radiologic confirmation (eg, MRI of bone). If the FDG PET/CT scan is positive in the mediastinum, the lymph node status needs pathologic confirmation.^{112,1025} Transesophageal EUS-FNA and EBUS-TBNA have proven useful to stage patients or to diagnose mediastinal lesions; these techniques can be used instead of invasive staging procedures in select patients.¹⁰²⁶⁻¹⁰²⁹ When compared with CT and PET, EBUS-TBNA has a high sensitivity and specificity for staging mediastinal and hilar lymph nodes in patients with lung cancer.¹⁰³⁰ In patients with positive nodes on CT or PET, EBUS-TBNA can be used to clarify the results.^{1031,1032} In patients with negative findings on EBUS-TBNA, conventional mediastinoscopy can be done to confirm the results.^{1027,1032-1034} Note that EBUS is also known as endosonography.

The routine use of bone scans (to exclude bone metastases) is not recommended. Brain MRI with contrast is recommended to rule out asymptomatic brain metastases in patients with stage II, III, and IV disease if aggressive combined-modality therapy is being considered.¹⁰³⁵ Patients with stage IB NSCLC are less likely to have brain metastases; therefore, brain MRI is optional in this setting and can be considered for select patients at high risk (eg, tumors >5 cm, central location). If brain MRI cannot be done, then CT of the head with contrast is an option. Note that PET scans are not recommended for assessing whether brain metastases are present (see the NCCN Guidelines for Central Nervous System Cancers, available at www.NCCN.org).

Initial Therapy

As previously mentioned, accurate pathologic assessment and staging are essential before treatment for NSCLC, because management varies depending on the stage, histology, presence of genetic variants, and PS. Before treatment, it is strongly recommended that determination of tumor resectability be made by thoracic surgeons who perform lung cancer



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

surgery as a prominent part of their practice (see *Principles of Surgical Therapy* in the NCCN Guidelines for NSCLC). RT doses are provided in the algorithm (see *Principles of Radiation Therapy* in the NCCN Guidelines for NSCLC). In addition, the NCCN Guidelines also recommend regimens for targeted therapy, immunotherapy, chemotherapy, and chemoradiation (see *Perioperative Systemic Therapy, Concurrent Chemoradiation Regimens, Molecular and Biomarker-Directed Therapy for Advanced or Metastatic Disease*, and *Systemic Therapy for Advanced or Metastatic Disease* in the NCCN Guidelines for NSCLC). First-line targeted therapy options are recommended for eligible patients with metastatic NSCLC and positive test results for actionable driver mutations such as *ALK*, *BRAF* p.V600E, *EGFR*, *MET*ex14 skipping, *NTRK1/2/3*, *RET*, and *ROS1*. Second-line targeted therapy options are recommended for eligible patients with metastatic NSCLC and positive test results for *EGFR* exon 20 insertions or *KRAS* p.G12C mutations. First-line immunotherapy options are recommended for eligible patients with metastatic NSCLC and negative test results for actionable driver mutations. Immunotherapy options may be used as first-line or subsequent therapy options in eligible patients with actionable driver mutations as shown in the algorithm, although they are generally not preferred options.

Stage I, Stage II, and Stage IIIA Disease

Depending on the extent and type of comorbidity present, patients with stage I or a subset of stage II (T1–2,N1) tumors are generally candidates for surgical resection and mediastinal lymph node dissection. Definitive RT, preferably SABR, is recommended for patients with early-stage NSCLC who are medically inoperable or refuse surgery; RT can be considered as an alternative to surgery in patients at high risk of complications (see *Stereotactic Ablative Radiotherapy* in this Discussion and see *Initial Treatment* for Stage I and II in the NCCN Guidelines for NSCLC).^{386,407,410,485,549,1036} Image-guided thermal ablation (eg,

cryotherapy, microwave, RFA) is an option for selected patients who are medically inoperable and not receiving SABR or definitive RT (see *Principles of Image-Guided Thermal Ablation Therapy* in the NCCN Guidelines for NSCLC).^{386,553-559} In some instances, positive mediastinal nodes (N2) are discovered at surgery; in this setting, an additional assessment of staging and tumor resectability must be made, and the treatment (ie, inclusion of systematic mediastinal lymph node dissection) must be modified accordingly. Therefore, the NCCN Guidelines include two different tracks for T1–2,N2 disease (ie, stage IIIA disease): 1) T1–2,N2 disease discovered unexpectedly at surgical exploration; and 2) T1–2,N2 disease confirmed before thoracotomy. In the second case, an initial brain MRI with contrast and FDG PET/CT scan (if not previously done) are recommended to rule out metastatic disease.

For patients with clinical stage IIB (T3,N0) and stage IIIA tumors who have different treatment options (surgery, RT, or chemotherapy), a multidisciplinary evaluation is recommended before treatment. For the subsets of stage IIB (T3,N0) and stage IIIA (T4,N0–1) tumors, treatment options are organized according to the location of the tumor, such as the superior sulcus, chest wall, proximal airway, or mediastinum.³⁹⁶ For each location, a thoracic surgeon needs to determine whether the tumor is resectable (see *Principles of Surgical Therapy* in the NCCN Guidelines for NSCLC). Preoperative concurrent chemoradiation followed by surgical resection of a superior sulcus tumor has shown 2-year survival in the 50% to 70% range.^{396,505,507,1037-1039} The overall 5-year survival rate is approximately 40%.⁵⁰⁵ For patients with resectable tumors (T3 invasion,N0–1) in the superior sulcus, the NCCN NSCLC Panel recommends preoperative concurrent chemoradiation followed by surgical resection and adjuvant chemotherapy plus either atezolizumab, pembrolizumab, or osimertinib, if patients have certain *EGFR* mutations or PD-L1 levels of 1% or more (see *Initial Treatment* for Superior Sulcus Tumors in the NCCN Guidelines for NSCLC). Patients with possibly



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

resectable superior sulcus tumors should undergo preoperative concurrent chemoradiation before surgical re-evaluation (including CT with or without contrast ± PET/CT). For patients with unresectable tumors (T4 extension, N0–1) in the superior sulcus, definitive concurrent chemoradiation is recommended followed by durvalumab (category 1).^{781,1040}

Definitive concurrent chemoradiation is recommended for patients with medically inoperable stage II or III NSCLC. The NCCN NSCLC Panel recommends durvalumab (category 1) as a consolidation immunotherapy option for eligible patients with unresectable stage III NSCLC and without disease progression after treatment with definitive concurrent chemoradiation based on clinical trial data and FDA approval (see *Chemoradiation: Trial Data and NCCN Recommendations* in this Discussion and the NCCN Guidelines for NSCLC).^{19,365,786} The panel also recommends durvalumab as a consolidation immunotherapy option (regardless of PD-L1 status) for eligible patients (PS 0–1) with unresectable stage II NSCLC but without disease progression after definitive concurrent platinum-based chemoradiation. The recommendation for consolidation immunotherapy with durvalumab occurs in multiple places in the NCCN Guidelines.

Surgical resection is the preferred treatment option for patients with tumors of the chest wall, proximal airway, or mediastinum (T3–4, N0–1). Other treatment options include preoperative systemic therapy or concurrent chemoradiation before surgical resection. For unresectable tumors (T4, N0–1) without pleural effusion, definitive concurrent chemoradiation (category 1) is recommended followed by consolidation immunotherapy with durvalumab (category 1).^{19,449,703,786} Additional chemotherapy (ie, consolidation chemotherapy) is an option if patients will not be receiving durvalumab.^{534,778} However, consolidation chemotherapy is not recommended if patients will be receiving

durvalumab, based on concerns that consolidation chemotherapy will increase the risk of pneumonitis if patients are also receiving durvalumab.

Multimodality therapy is recommended for most patients with stage III NSCLC.⁷⁷⁴ For patients with stage IIIA disease and positive mediastinal nodes (T1–2, N2), treatment is based on the findings of pathologic mediastinal lymph node evaluation (see the NCCN Guidelines for NSCLC). Patients with negative mediastinal biopsy findings are candidates for surgery. For those patients with resectable lesions, mediastinal lymph node dissection or lymph node sampling should be performed during the operation. Those individuals who are medically inoperable should be treated according to their clinical stage. For patients with (T1–2) N2 node-positive disease, a brain MRI with contrast and FDG PET/CT scan (if not done previously) are recommended to search for distant metastases. When distant metastases are not present, the NCCN NSCLC Panel recommends that the patient be treated with definitive concurrent chemoradiation therapy.^{484,704} Recommended therapy for metastatic disease depends on whether disease is in a solitary site or is widespread.

When a lung metastasis is present, it usually occurs in a patient with other systemic metastases; the prognosis is poor. Therefore, many of these patients are not candidates for surgery; however, systemic therapy is recommended. Although uncommon, patients with lung metastases but without systemic metastases have a better prognosis and are candidates for surgery (see *Multiple Lung Cancers* in this Discussion).¹⁰⁴¹ Patients with separate pulmonary nodule(s) in the same lobe (T3, N0–1) or ipsilateral non-primary lobe (T4, N0–1), without other systemic metastases, are potentially curable by surgery; 5-year survival rates are about 30%.¹⁰⁴² For those with N2 nodes after surgery, concurrent chemoradiation is recommended for those with positive margins and an R2 resection; either



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

sequential or concurrent chemoradiation is recommended after an R1 resection. Most NCCN Member Institutions favor concurrent chemoradiation for positive margins, but sequential chemoradiation is reasonable in frailer patients.⁷⁶⁸ For those with N2 nodes and negative margins, 1 chemotherapy (category 1) or sequential chemotherapy with RT is recommended. Adjuvant chemotherapy followed by adjuvant atezolizumab, pembrolizumab, or osimertinib is recommended for those with N0–1, depending on *EGFR* mutation levels and PD-L1 levels (see the NCCN Guidelines for NSCLC). In patients with synchronous solitary nodules (contralateral lung), the NCCN NSCLC Panel recommends treating them as two primary lung tumors if both are curable, even if the histology of the two tumors is similar.¹⁰⁴³

Multiple Lung Cancers

Patients with a history of lung cancer or those with biopsy-proven synchronous lesions may be suspected of or confirmed with having multiple lung cancers.^{1044,1045} It is important to determine whether the multiple lung cancers are metastases or separate lung primaries (synchronous or metachronous); most multiple lung tumors are metastases.^{89,396,1046,1047} Lesions with different cell types, such as squamous cell or adenocarcinoma, are usually different primary tumors. However, lesions of the same cell type may not be metastases. Therefore, it is essential to determine the histology of the lung tumor (see *Principles of Pathologic Review* in the NCCN Guidelines for NSCLC). Infection and other benign diseases also need to be ruled out (eg, inflammatory granulomas).^{1048,1049} Although criteria have been established for diagnosing multiple lung cancers, no definitive method has been established before treatment.¹⁰⁴⁹⁻¹⁰⁵² The Martini and Melamed criteria are often used to diagnose multiple lung cancers as follows: 1) the histologies are different; or 2) the histologies are the same, but there is no lymph node involvement and no extrathoracic metastases.¹⁰⁵² Data suggest that

NGS testing may help determine whether separate lung nodules are clonally related.¹⁰⁵³⁻¹⁰⁵⁵

Broad molecular profiling can be used to assess multiple lung lesions (see *Summary of the Guidelines Updates* in the NCCN Guidelines for NSCLC). For example, tumors with non-overlapping, unique mutations are considered to be clonally unrelated, separate primary lung cancers even if they are histologically similar. Therefore, these tumors can be treated with local therapy (see *Multiple Lung Cancers* in the NCCN Guidelines for NSCLC). Treatment of multiple lung cancers depends on the status of the lymph nodes (eg, N0–1) and on whether patients are asymptomatic, symptomatic, or at high or low risk of becoming symptomatic.^{1046,1056-1058} Patients should be evaluated in a multidisciplinary setting by thoracic surgeons, pulmonary medicine, radiation oncologists, and medical oncologists. In patients eligible for definitive local therapy, parenchymal-sparing resection is preferred (see the *Principles of Surgical Therapy* in the NCCN Guidelines for NSCLC).^{1045,1046} VATS or SABR are reasonable options depending on the number and distribution of the tumors requiring local treatment.¹⁰⁵⁹ Multiple lung nodules (eg, solid, subsolid nodules) may also be detected on CT scans; some of these nodules can be followed with imaging, whereas others need to be biopsied or excised (see *Incidental Lung Nodules* in this Discussion and the NCCN Guidelines for Lung Cancer Screening, available at www.NCCN.org).¹⁰⁶⁰

For the 2023 update (Version 1), the NCCN NSCLC Panel revised the algorithm for multiple lung cancers. For example, the NCCN Panel added an initial recommendation for multidisciplinary evaluation for patients suspected of or confirmed with having multiple lung cancers. One of the goals during multidisciplinary evaluation for multiple lung cancers is to assess whether patients have lung nodules that can be observed instead of erroneously assuming that patients have stage IV NSCLC. The panel added the following caveats: lesions at low risk of becoming symptomatic



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

can be observed, such as small subsolid nodules with slow growth. However, treatment should be considered if the lesion(s) show accelerating growth, increasing solid component, or increasing FDG uptake, even if the lesions are small.

Stage IIIB and IIIC NSCLC

Stage IIIB NSCLC comprises two unresectable groups, including: 1) T1–2,N3 tumors; and 2) T3–4,N2 tumors; stage IIIC NSCLC includes T3,N3 and contralateral mediastinal nodes (T4,N3), which are also unresectable. Surgical resection is not recommended in patients with T1–2,N3 disease. However, in patients with suspected N3 disease, the NCCN Guidelines recommend pathologic confirmation of nodal status (see *Pretreatment Evaluation* in the NCCN Guidelines for NSCLC).^{1061,1062} In addition, FDG PET/CT scans (if not previously done) and brain MRI with contrast should also be included in the pretreatment evaluation. If these imaging tests are negative, then treatment options for the appropriate nodal status should be followed (see the NCCN Guidelines for NSCLC). If N3 disease is confirmed, definitive concurrent chemoradiation (category 1) is recommended followed by durvalumab (category 1).^{449,703,781,786,1063-1065}

Durvalumab is recommended (category 1) as a consolidation immunotherapy option for eligible patients with unresectable stage III NSCLC and without disease progression after treatment with definitive concurrent chemoradiation (see *Chemoradiation: Trial Data and NCCN Recommendations* in this Discussion and the NCCN Guidelines for NSCLC).^{19,365,786} If patients will be receiving durvalumab, additional chemotherapy (ie, consolidation chemotherapy) is not recommended based on concerns that adding consolidation chemotherapy will increase the risk of pneumonitis if patients are also receiving durvalumab. If patients will not be receiving durvalumab because of medical contraindications or other reasons, consolidation chemotherapy is an option after concurrent chemoradiation.^{534,778} For metastatic disease that

is confirmed by FDG PET/CT scan and brain MRI with contrast, treatment is described in the NCCN Guidelines for limited or metastatic disease.

For patients with T4,N2–3 disease (stages IIIB and IIIC), surgical resection is not recommended. The initial workup includes biopsies of the N3 and N2 nodes. If these biopsies are negative, the same treatment options may be used as for stage IIIA (T4,N0–1) disease (see the NCCN Guidelines for NSCLC). If either the contralateral or ipsilateral mediastinal node is positive, definitive concurrent chemoradiation therapy is recommended (category 1) followed by durvalumab (see the NCCN Guidelines for NSCLC).^{449,703,781,786,1063-1066} Again, durvalumab is recommended (category 1) as a consolidation immunotherapy option for eligible patients with unresectable stage III NSCLC and without disease progression after treatment with definitive concurrent chemoradiation.^{19,365,786} Consolidation chemotherapy is an option for eligible patients.^{534,778} However, consolidation chemotherapy is not recommended if patients will be receiving durvalumab.

Limited Metastatic Disease

In general, systemic therapy is recommended for patients with metastatic disease (see *Systemic Therapy for Advanced or Metastatic Disease* in the NCCN Guidelines for NSCLC).⁸⁸⁶ In addition, palliative treatment, including RT, may be needed during the disease course to treat localized symptoms, diffuse brain metastases, or bone metastases (see *Therapy for Recurrence and Metastasis* in the NCCN Guidelines for NSCLC). This section focuses on patients with limited metastatic disease; management of widespread distant metastases is described in another section (see *Systemic Therapy for Advanced or Metastatic Disease* in the NCCN Guidelines for NSCLC). Biomarker testing is recommended for patients with stage IVA disease (see *Predictive and Prognostic Biomarkers* in this Discussion).



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

Pleural or pericardial effusion is a criterion for stage IV, M1a disease. T4 with pleural effusion is classified as stage IV, M1a (see Table 3 in *Staging* in the NCCN Guidelines for NSCLC).¹⁶¹ Pleural or pericardial effusions are malignant in 90% to 95% of patients; however, they may be related to obstructive pneumonitis, atelectasis, lymphatic or venous obstruction, or a pulmonary embolus. Therefore, pathologic confirmation of a malignant effusion by using thoracentesis or pericardiocentesis is recommended. In certain cases where thoracentesis is inconclusive, thoracoscopy may be performed. In the absence of nonmalignant causes (eg, obstructive pneumonia), an exudate or sanguineous effusion is considered malignant regardless of the results of cytologic examination. If the pleural or pericardial effusion is considered negative for malignancy (M0), recommended treatment is based on the confirmed T and N stage (see the NCCN Guidelines for NSCLC). Data suggest that most pleural or pericardial effusions are associated with unresectable disease in 95% of cases.¹⁰⁶⁷ In patients with effusions that are positive for malignancy, the tumor is defined as M1a and is treated with local therapy (ie, ambulatory small catheter drainage, pleurodesis, and pericardial window) in addition to treatment as for stage IV disease.^{1068,1069}

Care of patients with distant metastases in limited sites (ie, stage IVA, M1b) and good PS depends on the location and number of the metastases; the diagnosis is aided by mediastinoscopy, bronchoscopy, FDG PET/CT scan, and brain MRI with contrast. The increased sensitivity of FDG PET/CT scans, compared with other imaging methods, may identify additional metastases and, thus, spare some patients from unnecessary futile surgery. Positive FDG PET/CT scan findings for distant disease need pathologic or other radiologic confirmation. If the FDG PET/CT scan is positive in the mediastinum, the lymph node status needs pathologic confirmation. Patients with oligometastatic disease (eg, brain metastases) and otherwise limited disease in the chest may benefit from aggressive local therapy to both the primary chest and metastatic

sites.^{586,1070} Aggressive local therapy may comprise surgery and/or definitive RT, including SRS and SABR, and may be preceded or followed by chemotherapy. After progression on TKIs, patients with *EGFR* mutation–positive metastatic NSCLC may be able to continue with their current TKIs; local therapy can be considered to treat their limited progression (eg, SRS to brain metastases or other sites, SABR for limited thoracic or other metastatic disease; surgery).^{1071,1072}

Neoadjuvant or Adjuvant Therapy

Chemotherapy, Chemoradiation, Immunotherapy, and Targeted Therapy

On the basis of clinical studies,⁶⁹¹⁻⁶⁹³ the NCCN NSCLC Panel recommends cisplatin combined with docetaxel, etoposide, gemcitabine, or vinorelbine as neoadjuvant or adjuvant therapy options (also known as perioperative therapy) for all histologies in eligible patients with locally advanced disease. The NCCN NSCLC Panel has preference stratified all the systemic therapy regimens and decided that cisplatin plus pemetrexed is a preferred neoadjuvant or adjuvant therapy option (also known as preoperative and postoperative therapy) for nonsquamous NSCLC, whereas cisplatin plus either gemcitabine or docetaxel is preferred for squamous cell NSCLC (see *Perioperative Systemic Therapy* in the NCCN Guidelines for NSCLC).^{739,744,790} Cisplatin combined with either vinorelbine or etoposide are “other recommended” options. For patients with comorbidities or those who cannot tolerate cisplatin, carboplatin may be combined with pemetrexed (nonsquamous only), paclitaxel, or gemcitabine; thus, these regimens are useful in certain circumstances.^{739,1073} These neoadjuvant or adjuvant regimens may also be used for sequential chemoradiation.⁷⁴⁵⁻⁷⁴⁸

Because patients with stage III disease have both local and distant failures, theoretically, the use of chemotherapy may eradicate micrometastatic disease obviously present but undetectable at diagnosis.



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

The timing of this chemotherapy varies (see the NCCN Guidelines for NSCLC). Such chemotherapy may be given alone, sequentially, or concurrently with RT. In addition, chemotherapy could be given preoperatively or postoperatively in appropriate patients. Three phase 3 trials have assessed preoperative chemotherapy followed by surgery compared with surgery alone in the treatment of stage III NSCLC.^{698,1074-1076} All three studies showed a survival advantage for patients who received preoperative chemotherapy. SWOG S9900—one of the largest randomized trials examining preoperative chemotherapy in early-stage NSCLC—assessed surgery alone compared with surgery plus preoperative paclitaxel plus carboplatin in patients with stage IB/IIA and stage IIB/IIIA NSCLC (excluding superior sulcus tumors). PFS and overall survival were improved with preoperative chemotherapy.^{1075,1076} The two earlier phase 3 studies had a small number of patients, while the SWOG study was stopped early because of the positive results of the IALT study. A number of phase 2 studies have evaluated preoperative chemotherapy for stage III NSCLC, with (or without) RT, followed by surgery.¹⁰⁷⁷⁻¹⁰⁷⁹

Post-surgical treatment options for patients with stage IA tumors (T1abc,N0) and with positive surgical margins (R1, R2) include re-resection (preferred) or RT (category 2B); observation is recommended for patients with negative surgical margins (R0). Postoperative chemotherapy is a recommended option for patients with T2ab,N0 tumors and negative surgical margins who have high-risk features, including poorly differentiated tumors, vascular invasion, wedge resection, visceral pleural involvement, and unknown lymph node status (Nx) (see the NCCN Guidelines for NSCLC).^{738,1080} If the surgical margins are positive in patients with T2ab,N0 tumors, options include: 1) re-resection (preferred) with (or without) chemotherapy; or 2) RT with (or without) chemotherapy (chemotherapy is an option for T2b,N0).^{469,738}

The NCCN NSCLC Panel recommends atezolizumab as an adjuvant therapy option for eligible patients with completely resected (R0) stage IIB to IIIA, stage IIB (only T3,N2), or high-risk stage IIA NSCLC and PD-L1 of 1% or more, and negative for *EGFR* exon 19 deletions, *EGFR* L858R mutations, or *ALK* rearrangements, who have previously received adjuvant chemotherapy based on clinical trial data and FDA approval (see *Surgery Followed by Adjuvant Therapy: Trial Data and NCCN Recommendations* in this Discussion).⁷⁵¹ For the 2023 update (Version 2), the NCCN Panel also recommends pembrolizumab as an adjuvant therapy option following adjuvant chemotherapy for eligible patients with completely resected early-stage NSCLC based on clinical trial data and FDA approval.³² The panel recommends osimertinib as an adjuvant therapy option for eligible patients with completely resected (R0) stage IB to IIIA or stage IIB (only T3,N2) NSCLC and *EGFR* exon 19 deletions or *EGFR* L858R mutations who have previously received adjuvant chemotherapy or are ineligible to receive platinum-based chemotherapy based on clinical trial data and FDA approval.⁷⁴⁹ For patients who have these *EGFR* mutations and PD-L1 levels of 1% or more, the panel only recommends osimertinib; the panel does not recommend using 1) single-agent therapy with either atezolizumab or pembrolizumab; or 2) combination therapy with osimertinib plus atezolizumab or pembrolizumab.

The NCCN NSCLC Panel recommends adjuvant chemotherapy (category 1) followed by atezolizumab, pembrolizumab, or osimertinib for eligible patients with the appropriate biomarkers, negative surgical margins, and stage IIB disease, including 1) T1abc–T2a,N1; 2) T2b,N1; or 3) T3,N0 disease.^{734,1081} If surgical margins are positive in these patients, options after an R1 resection include: 1) re-resection and chemotherapy; or 2) chemoradiation (either sequential or concurrent). After an R2 resection, options include: 1) re-resection and chemotherapy; or 2) concurrent chemoradiation. Most NCCN Member Institutions favor concurrent



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

chemoradiation for positive margins, but sequential chemoradiation is reasonable in frailer patients.⁷⁶⁸ Postoperative chemotherapy or chemoradiation can also be used in patients with stage III NSCLC who have had surgery (see the NCCN Guidelines for NSCLC). Patients with T1–3,N2 or T3,N1 disease (discovered only at surgical exploration and mediastinal lymph node dissection) and positive margins may be treated with chemoradiation; either sequential or concurrent chemoradiation is recommended for an R1 resection, whereas concurrent chemoradiation is recommended for an R2 resection. For patients with negative margins (R0), treatment options include either 1) chemotherapy (category 1) followed by atezolizumab, pembrolizumab, or osimertinib for eligible patients with the appropriate biomarkers; or 2) sequential chemotherapy and consider RT.⁷³⁴

For stage IIIA superior sulcus tumors (T4 extension,N0–1) that become resectable after preoperative concurrent chemoradiation, resection followed by chemotherapy is recommended and then atezolizumab, pembrolizumab, or osimertinib, depending on biomarker status (see the NCCN Guidelines for NSCLC). Surgical reevaluation (including chest CT with or without contrast and with or without PET/CT) is done to determine whether the tumor is resectable after treatment. If the lesion remains unresectable after preoperative concurrent chemoradiation, then completion of definitive dose chemoradiation without interruption, followed by consolidation immunotherapy with durvalumab (category 1) is recommended for eligible patients. Among patients with chest wall lesions with T3 invasion–T4 extension, N0–1 disease, those who are initially treated with surgery (preferred) may receive chemotherapy and then either atezolizumab, pembrolizumab, or osimertinib depending on biomarker status, if the surgical margins are negative. For patients with positive margins, options include either 1) sequential or concurrent chemoradiation; or 2) re-resection and chemotherapy. As previously mentioned, most NCCN Member Institutions favor concurrent

chemoradiation for positive margins, but sequential is reasonable in frailer patients.⁷⁶⁸ Similar treatment plans are recommended for resectable tumors of the proximal airway or mediastinum (T3–4,N0–1).

For patients with stage III disease and positive mediastinal nodes (T1–3,N2) and no apparent disease progression after initial treatment with induction systemic therapy with or without RT, surgery is recommended (see the NCCN Guidelines for NSCLC).⁴⁵² Alternatively, if the disease progresses, treatment options include either 1) local therapy using RT (if feasible) with (or without) chemotherapy; or 2) systemic therapy. In patients with separate pulmonary nodules in the same lobe (T3,N0–1) or ipsilateral non-primary lobe (T4,N0–1), surgery is recommended; neoadjuvant systemic therapy with nivolumab plus chemotherapy is an option before surgery. In patients with N2 disease and negative margins, options include 1) chemotherapy (category 1); or 2) sequential chemotherapy with radiation. If the resection margins are positive in patients with N2 disease, concurrent chemoradiation is recommended for an R2 resection, whereas either concurrent or sequential chemoradiation is recommended for an R1 resection. Concurrent chemoradiation is often used for positive margins, but sequential is reasonable in frailer patients.

Radiation Therapy

After complete resection of clinical early-stage NSCLC, postoperative RT has been found to be detrimental for pathologic N0 or N1 stage disease in a meta-analysis (population-based analysis of data from SEER) of small randomized trials using older techniques and dosing regimens.¹⁰⁸² Data from two randomized phase 3 trials, the LungART and PORT-C trials, show that postoperative RT (also known as PORT) did not improve survival compared with no postoperative RT, although locoregional control was significantly improved in patients with completely resected stage IIIA (N2) NSCLC who received neoadjuvant or adjuvant chemotherapy.^{452,455} Postoperative RT may be considered for select



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

patients with negative margins and high-risk N2 disease (see *Chemoradiation: Trial Data and NCCN Recommendations in this Discussion and Principles of Radiation Therapy* in the algorithm). Previous trials and meta-analyses had suggested that there was an apparent survival benefit of postoperative RT in patients with N2 nodal stage diagnosed surgically.^{469,576,1081,1083,1084}

PROCLAIM, a phase 3 randomized trial, assessed concurrent thoracic RT with cisplatin plus pemetrexed versus cisplatin plus etoposide followed by consolidation chemotherapy in patients with unresectable stage III nonsquamous NSCLC.⁷⁷⁸ Both regimens were equivalent in terms of survival, but the cisplatin plus pemetrexed regimen was associated with less neutropenia (24.4% vs. 44.5%; $P < .001$) and fewer grade 3 to 4 adverse events (64.0% vs. 76.8%; $P = .001$). The NCCN Panel has preference stratified the concurrent chemoradiation regimens and decided that pemetrexed with either carboplatin or cisplatin are preferred concurrent chemoradiation regimens for eligible patients with nonsquamous NSCLC.^{783,1085,1086} Other preferred concurrent chemoradiation regimens include 1) paclitaxel plus carboplatin; and 2) cisplatin plus etoposide; these regimens may be used regardless of histology.^{534,780} The NCCN NSCLC Panel deleted the cisplatin plus etoposide consolidation regimen based on the PROCLAIM trial.⁷⁷⁸ Other consolidation chemotherapy regimens are an option for eligible patients receiving definitive chemoradiation; however, consolidation chemotherapy is not recommended if the patient will be receiving durvalumab.

Postoperative chemotherapy (category 1) followed by adjuvant atezolizumab, pembrolizumab, or osimertinib, depending on the biomarker status, is recommended for patients with T1–3, N2 disease and negative margins (see the NCCN Guidelines for NSCLC). Postoperative sequential chemotherapy with consideration of postoperative RT is recommended for select patients with negative margins and high-risk N2 disease such as

extracapsular extension, multistation involvement, inadequate lymph node dissection or sampling, and/or refusal or intolerance of adjuvant systemic therapy.^{452,455,767} Either concurrent or sequential chemoradiation may be used for postoperative therapy and positive margins, depending on the type of resection and the setting (eg, N2 disease). Concurrent chemoradiation is recommended for R2 resections, whereas either sequential or concurrent chemoradiation is recommended for R1 resections. Concurrent chemoradiation is often used for positive margins, but sequential is reasonable in frailer patients.⁷⁶⁸ Cisplatin plus etoposide and carboplatin plus paclitaxel are chemoradiation regimens recommended by the NCCN NSCLC Panel for all histologies (see *Concurrent Chemoradiation Regimens* in the NCCN Guidelines for NSCLC).⁷⁸⁰ When chemoradiation is recommended in the NCCN Guidelines, these regimens may be used for stage II to III disease.^{470,471,703,704,781,784,785}

Surveillance

Because recurrence is common after treatment for NSCLC, surveillance is recommended in the NCCN Guidelines for eligible patients with no evidence of clinical or radiographic disease after definitive therapy. Data from randomized phase 3 trials are not available to clarify surveillance recommendations; therefore, the most appropriate schedules are controversial.¹⁰⁸⁷⁻¹⁰⁹² The surveillance recommendations were compiled by polling the NCCN NSCLC Panel regarding their practice patterns. Details regarding the specific surveillance schedules are outlined in the algorithm based on stage and the definitive treatment (see *Surveillance After Completion of Definitive Therapy* in the algorithm). Surveillance schedules for most patients with metastatic disease are individualized for each patient, although the NCCN Guidelines provide a surveillance schedule for certain patients with stage IV oligometastatic disease.



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

NLST, a large randomized trial, assessed lung screening with low-dose CT screening versus chest radiography in individuals at high risk for lung cancer.⁷⁶ Low-dose CT screening decreased mortality from lung cancer (mainly adenocarcinoma) compared with chest radiography (247 vs. 309 deaths, respectively; 20% relative reduction in mortality; 95% CI, 6.8–26.7; $P = .004$).⁷⁶ Low-dose CT is recommended for screening individuals at high risk for lung cancer (see the NCCN Guidelines for Lung Cancer Screening, available at www.NCCN.org). The NCCN NSCLC Panel feels that low-dose CT is beneficial for identifying recurrences in patients previously treated for NSCLC. It is important to note that the surveillance recommendations for patients who have been treated for NSCLC are different from the screening recommendations for individuals at high risk for lung cancer (see the NCCN Guidelines for Lung Cancer Screening).

The NCCN Guidelines recommend a chest CT scan with (or without) contrast and an H&P for the initial surveillance schedules (2–5 years after definitive treatment) followed by annual low-dose non-contrast-enhanced CT and an H&P (see *Surveillance After Completion of Definitive Therapy* in the algorithm).^{1090,1091,1093-1096} Patients treated with chemotherapy with (or without) RT who have residual abnormalities may require more frequent imaging. FDG PET/CT or brain MRI is not routinely recommended for routine surveillance in patients without symptoms.¹⁰⁸⁷ But, PET may be useful for assessing CT scans that appear to show malignant neoplasms but may be radiation fibrosis, atelectasis, or other benign conditions. Areas previously treated with RT may remain FDG avid for up to 2 years; therefore, histologic confirmation of suspicious areas with apparent “recurrent” disease is needed.¹⁰⁹⁷ The NCCN NSCLC Panel recommends assessing patients with recurrences using PET/CT and brain MRI with contrast; if brain MRI is not possible, then CT with contrast of the head is recommended. Information about smoking cessation (eg, advice, counseling, therapy) should be provided for patients undergoing surveillance to improve their quality of life.

The NCCN Guidelines include information about the long-term follow-up care of NSCLC survivors (see *Cancer Survivorship Care* in the NCCN Guidelines for NSCLC). These recommendations include guidelines for routine cancer surveillance, immunizations, health monitoring, counseling for wellness and health promotion, and cancer screening (see the NCCN Guidelines for Colorectal Cancer Screening, NCCN Guidelines for Breast Cancer Screening and Diagnosis, and NCCN Guidelines for Prostate Cancer Early Detection, available at www.NCCN.org). An analysis suggests that patients who survive lung cancer have a high symptom burden 1 year after diagnosis and therefore need management after treatment.¹⁰⁹⁸

Treatment of Recurrences and Distant Metastases

Recurrences are subdivided into locoregional recurrences and distant metastases. Management of locoregional recurrences or symptomatic local disease—endobronchial obstruction, mediastinal lymph node recurrence, superior vena cava (SVC) obstruction, and severe hemoptysis—is described in the NCCN Guidelines (see *Therapy for Recurrence and Metastasis* in the NCCN Guidelines for NSCLC).⁶ An SVC stent may be used with either concurrent chemoradiation or RT to treat SVC obstruction. For patients with endobronchial obstruction, relieving airway obstruction may increase survival, especially in patients who are severely compromised, and may improve their quality of life.¹⁰⁹⁹ After treatment for locoregional recurrence, observation or systemic therapy (category 2B for systemic therapy) is recommended if disseminated disease is not evident. Systemic therapy is recommended for disseminated disease. The type of systemic therapy depends on the histologic type, whether somatic genomic alterations are present that can be treated with targeted therapy, and PS (see *Systemic Therapy for Advanced or Metastatic Disease* in the NCCN Guidelines for NSCLC).



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

Management of distant metastases—localized symptoms; bone; and limited, diffuse brain, or disseminated metastases—is described in the NCCN Guidelines (see *Therapy for Recurrence and Metastasis* in the NCCN Guidelines for NSCLC). Palliation of symptoms throughout the disease course can be achieved with external-beam RT for distant metastases with localized symptoms, diffuse brain metastases, or bone metastases (bisphosphonate or denosumab therapy can be considered).^{482,581,1100} For patients at risk of fracture in weight-bearing bone, orthopedic stabilization and palliative RT are recommended.

Of note, recurrent and metastatic disease have historically been regarded as incurable. However, select limited locoregional recurrences may be treated with curative intent therapy (re-resection preferred or RT or SABR) (see *Therapy for Recurrence and Metastasis* in the NCCN Guidelines for NSCLC). Similarly, patients with limited-site oligometastatic disease and good PS may benefit from aggressive local therapies to the metastatic and primary sites, with clinical data suggesting the possibility of long-term survival (see *Initial Treatment for Stage IVA, M1b* in the NCCN Guidelines for NSCLC).^{609,610,613,656,1101-1104} In addition, emerging clinical data suggest the feasibility of definitive reirradiation of local recurrences within prior RT fields using highly conformal techniques, although this should be limited to highly selected cases in specialty centers with appropriate expertise because of the potential for severe toxicity with high cumulative radiation doses to critical structures.^{479,633-635,1105-1108}

In patients with NSCLC who have bone metastases, data suggest that denosumab increases median overall survival when compared with zoledronic acid (9.5 vs. 8 months).¹¹⁰⁹ The FDA has approved the use of zoledronic acid and denosumab in patients with bone metastases from solid tumors.^{1110,1111} Denosumab and bisphosphonate therapy can be associated with severe hypocalcemia; patients with hypoparathyroidism and vitamin D deficiency are at increased risk for hypocalcemia.

Denosumab or intravenous bisphosphonate therapy can be considered in patients with bone metastases to decrease bone complications (eg, decrease pain, delay skeletal-related events) based on clinical trial data.^{184,1109,1112-1115}

For patients with recurrent and metastatic disease, the NCCN Guidelines recommend that histologic subtype should be determined before therapy so that the best treatment can be selected (see *Systemic Therapy for Advanced or Metastatic Disease* in the NCCN Guidelines for NSCLC).⁷⁹⁰ Biomarker testing for somatic, disease-associated variants/mutations (ie, oncogenic driver events) is recommended before starting therapy, if feasible, in eligible patients with metastatic NSCLC, because targeted therapy has been shown to decrease tumor burden, decrease symptoms, and dramatically improve the quality of life for patients with specific variants. If molecular testing results are pending and patients require an urgent start to therapy, clinicians can consider holding immunotherapy for one cycle (ie, just use platinum-based chemotherapy regimens). In the NCCN Guidelines, many targeted agents are recommended for first-line therapy in patients with specific actionable mutations such as afatinib, alectinib, brigatinib, capmatinib, ceritinib, crizotinib, dacomitinib (± trametinib), entrectinib, erlotinib, gefitinib, lorlatinib, osimertinib, pralsetinib, seliprecatinib, and tepotinib (see *Molecular and Biomarker-Driven Therapy for Advanced or Metastatic Disease* in the algorithm).⁸⁸⁶ The number of available targeted agents is increasing. For example, newer agents are now recommended as second-line and beyond (subsequent) therapy options—such as adagrasib, amivantamab, fam-trastuzumab deruxtecan-nxki, mobocertinib, and sotorasib—in patients with specific actionable driver mutations.

Additional targeted therapies for patients with other somatic genomic alterations are also recommended, although there is less evidence for these agents and they have not been FDA approved for lung cancer (see



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

Emerging Biomarkers to Identify Novel Therapies for Patients with Metastatic NSCLC in the NCCN Guidelines for NSCLC). The following targeted agents are recommended as monotherapy options for patients with metastatic NSCLC and emerging genetic variants: capmatinib, crizotinib, or tepotinib for high-level *MET* amplification; (see *Agents that Inhibit High-Level MET Amplifications* in this Discussion).^{169-172,921,922}

Certain targeted therapies—such as alectinib, brigatinib, ceritinib, lorlatinib, and osimertinib—are recommended as subsequent therapies (if not previously given) for patients with the indicated somatic genomic alterations whose disease becomes resistant to first-line targeted therapies; other targeted therapies are being investigated for resistance.

Biomarker testing for actionable oncogenic driver mutations is recommended in the NCCN Guidelines based on the improved outcomes associated with use of targeted therapy in eligible patients with metastatic NSCLC (see *Principles of Molecular and Biomarker Analysis* in the NCCN Guidelines for NSCLC and *Predictive and Prognostic Biomarkers* in this Discussion). It is important to note that 1) several different tests may be used to identify the same biomarker, including FDA-approved biomarker tests and validated laboratory tests done in CLIA-approved laboratories; and 2) biomarker testing is rapidly changing and improving. *EGFR* mutation testing (category 1) is recommended in patients with nonsquamous NSCLC (ie, adenocarcinoma, large cell carcinoma) or NSCLC NOS, because *EGFR* TKIs are recommended for patients with certain *EGFR* mutations.^{121,246,254,258,1116} Testing for *ALK* rearrangements (category 1) is also recommended in patients with nonsquamous NSCLC, because *ALK* inhibitors are effective for patients with metastatic NSCLC who are positive for *ALK* rearrangements.^{185,1117} The NCCN NSCLC Panel also recommends testing for other biomarkers such as *BRAF*, *ERBB2* (*HER2*), *KRAS*, *MET*ex14, *NTRK1/2/3*, *RET*, *ROS1* rearrangements (see *Predictive and Prognostic Biomarkers* in this Discussion). The NCCN NSCLC Panel also recommends upfront PD-L1 expression testing

(category 1) before first-line therapy in patients with metastatic NSCLC to assess whether patients are candidates for ICIs. In 2020, the NCCN Panel deleted TMB as an emerging immune biomarker for patients with metastatic NSCLC based on clinical trial data and other issues (see *TMB* in this Discussion).

Molecular testing is recommended in all patients with metastatic nonsquamous NSCLC and NSCLC NOS. Compared with patients with nonsquamous NSCLC, those with squamous cell carcinoma have fewer actionable mutations (eg, *ALK* rearrangements, the common *EGFR* mutations) when considered as individual mutations. However, the cumulative incidence of targetable molecular alterations in squamous cell carcinoma across all alterations ranges from 2% to 10%; therefore, molecular testing can be considered in these patients, particularly if a diagnosis is based on a small sampling.^{139,140,186,187} The NCCN Panel recommends that molecular testing be considered in all patients with metastatic NSCLC squamous cell carcinoma and not just those with certain characteristics, such as never smoking status and mixed histology. Treatment recommendations and eligibility criteria for patients with nonsquamous NSCLC, NSCLC NOS, and squamous cell carcinoma are described in the NCCN Guidelines. These recommendations are briefly summarized in the following paragraphs. Data supporting the combination platinum-based chemotherapy options are described in the following sections (see *Trial Data* in this Discussion).

Single-agent targeted therapy is recommended for patients with actionable driver mutations [ie, *ALK* rearrangements, *EGFR* activating mutations, *ERBB2* (*HER2*) mutations, *KRAS* p.G12C mutations, *MET*ex14 skipping, *NTRK1/2/3* fusions, *RET* rearrangements, *ROS1* rearrangements], or those with emerging driver mutations (see *Molecular and Biomarker-Driven Therapy for Advanced or Metastatic Disease* and *Emerging Biomarkers to Identify Novel Therapies for Patients with*



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

Metastatic NSCLC in the NCCN Guidelines for NSCLC). Chemotherapy plus immunotherapy regimens are recommended for patients without targetable somatic genomic alterations (see *Systemic Therapy for Advanced or Metastatic Disease* in the NCCN Guidelines for NSCLC). Chemotherapy plus immunotherapy regimens—such as pembrolizumab plus carboplatin (or cisplatin) plus pemetrexed—are recommended for patients with metastatic nonsquamous NSCLC and negative test results for actionable driver mutations (also known as wild-type), regardless of PD-L1 expression (see *Systemic Therapy for Advanced or Metastatic Disease* in the NCCN Guidelines for NSCLC, the NCCN Drugs & Biologics Compendium [NCCN Compendium®] for NSCLC, and the NCCN Guidelines with Evidence Blocks™ for NSCLC).⁷⁹⁰ These immunotherapy plus chemotherapy regimens may be used as first-line or subsequent therapy options in eligible patients with actionable driver mutations as shown in the algorithm, although they are generally not preferred options.

For patients with metastatic NSCLC and contraindications to pembrolizumab or other ICIs, chemotherapy options are recommended (such as carboplatin plus paclitaxel), although some regimens may be more appropriate for certain patients, depending on histology, PS, and other factors (see *Trial Data* in this Discussion, and *Systemic Therapy for Advanced or Metastatic Disease* in the NCCN Guidelines for NSCLC, the NCCN Compendium® for NSCLC, and the NCCN Guidelines with Evidence Blocks™ for NSCLC).^{886,1118} Chemotherapy with or without bevacizumab is an option if eligibility criteria are met for patients with nonsquamous NSCLC and negative test results for actionable driver mutations, and with PD-L1 expression less than 1%.¹¹¹⁹ Chemotherapy with or without bevacizumab regimens may be used as first-line or subsequent therapy options in eligible patients with actionable driver mutations as shown in the algorithm, although they are generally not preferred options. Targeted therapy is a preferred first-line therapy option for patients with actionable driver mutations, except for *EGFR* exon 20

mutations or *KRAS* mutations. Previously, patients with brain metastases were excluded from receiving bevacizumab because of concerns about CNS hemorrhage; however, data suggest that bevacizumab can be used in patients with treated CNS metastases.¹¹²⁰ A phase 3 randomized trial in older patients (70–89 years) with advanced NSCLC reported that combined therapy with weekly paclitaxel and monthly carboplatin improved survival when compared with single-agent therapy using either gemcitabine or vinorelbine (10.3 vs. 6.2 months).¹¹²¹ Systemic therapy for older patients with advanced NSCLC needs to be carefully selected to avoid adverse reactions.¹¹²² The NCCN NSCLC Panel does not recommend carboplatin plus vinorelbine, cisplatin plus vinorelbine, etoposide, irinotecan, and vinorelbine for patients with metastatic nonsquamous NSCLC or NSCLC NOS and negative test results for actionable driver mutations, because these regimens are rarely used in the United States.

For patients with metastatic squamous cell NSCLC and negative test results for actionable driver mutations, chemotherapy plus immunotherapy regimens—such as pembrolizumab plus carboplatin with either paclitaxel or albumin-bound paclitaxel—are recommended options (category 1; preferred). For patients with metastatic squamous cell NSCLC who have contraindications to ICIs, recommended options include cisplatin plus gemcitabine (category 1).⁷⁹⁰ Carboplatin plus paclitaxel, carboplatin plus gemcitabine (category 1 for both), and other regimens listed in the NSCLC algorithm are also recommended options (see *Systemic Therapy for Advanced or Metastatic Disease* in the NCCN Guidelines for NSCLC, the NCCN Compendium® for NSCLC, and the NCCN Guidelines with Evidence Blocks™ for NSCLC). Data supporting the combination platinum-based chemotherapy options are described in the following sections (see *Clinical Trial Data* in this Discussion). These chemotherapy with or without immunotherapy regimens may be used as first-line or subsequent therapy options in eligible patients with actionable driver



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

mutations as shown in the algorithm, although they are generally not preferred options. Targeted therapy is a preferred first-line therapy option for patients with actionable driver mutations, except for *EGFR* exon 20 mutations or *KRAS* mutations. The NCCN NSCLC Panel does not recommend carboplatin plus etoposide, carboplatin plus vinorelbine, cisplatin plus vinorelbine, cisplatin plus gemcitabine plus necitumumab, etoposide, irinotecan, and vinorelbine for patients with metastatic squamous cell NSCLC and negative test results for actionable driver mutations, because these regimens are rarely used in the United States although they may be used in other countries. Regimens containing pemetrexed or bevacizumab are not recommended for squamous cell carcinoma. Currently, fewer treatment options are available for patients with squamous cell carcinoma compared with nonsquamous NSCLC.

Clinical Trial Data

Data show that platinum-based combination therapy is superior to best supportive care for patients with advanced, incurable disease who are not eligible for targeted therapy or immunotherapy. Cisplatin or carboplatin have been proven effective in combination with many of the following agents: docetaxel, etoposide, gemcitabine, paclitaxel (and albumin-bound paclitaxel), and pemetrexed (see *Systemic Therapy for Advanced or Metastatic Disease* in the NCCN Guidelines for NSCLC).^{739,744,788-790,797,798,818} Carboplatin-based regimens are often used for patients with comorbidities or those who cannot tolerate cisplatin.¹¹²³ Non-platinum regimens (eg, gemcitabine plus docetaxel, gemcitabine plus vinorelbine) are reasonable alternatives, because data show they are active and less toxic than platinum-based regimens.^{800-803,1124}

ECOG 4599, a phase 2/3 trial, randomly assigned 878 patients to either 1) bevacizumab in combination with paclitaxel plus carboplatin; or 2) paclitaxel plus carboplatin alone.^{806,1125} Both regimens were well tolerated with selected toxicities. Patients receiving bevacizumab plus paclitaxel

plus carboplatin showed an improved median survival (12.3 vs. 10.3 months, $P = .003$) when compared to patients receiving paclitaxel plus carboplatin alone.⁸⁰⁶ The overall 1-year survival was 51% versus 44%; 2-year survival was 23% versus 15%, respectively, in favor of the bevacizumab plus paclitaxel plus carboplatin arm.⁸⁰⁶ More significant toxicities were observed with bevacizumab plus paclitaxel plus carboplatin compared to paclitaxel plus carboplatin (grade 4 neutropenia: 25.5% vs. 16.8%; grade 5 hemoptysis: 1.2% vs. 0%; and grade 3 hypertension: 6.8% vs. 0.5%). Treatment-related deaths were more common with bevacizumab plus paclitaxel plus carboplatin (15 patients) than with paclitaxel plus carboplatin (2 patients) ($P = .001$). An analysis of ECOG 4599 found that patients with adenocarcinoma histology receiving bevacizumab plus paclitaxel plus carboplatin had improved survival compared with chemotherapy alone (14.2 vs. 10.3 months).¹¹¹⁹ AVAIL, a phase 3 randomized trial, compared cisplatin plus gemcitabine with (or without) bevacizumab; survival was not increased with the addition of bevacizumab.^{1126,1127} The NCCN NSCLC Panel recommends that bevacizumab biosimilars may be used in any of the systemic therapy regimens containing bevacizumab (eg, carboplatin plus paclitaxel plus bevacizumab) that are used for eligible patients with metastatic NSCLC based on clinical data and FDA approvals.⁸⁹⁶⁻⁹⁰⁰

A noninferiority trial in 1725 patients with advanced NSCLC (either stage IIIB or IV; most were stage IV) assessed cisplatin plus gemcitabine compared with cisplatin plus pemetrexed.⁷⁹⁰ Patients with either adenocarcinoma or large cell carcinoma (ie, nonsquamous NSCLC) had improved survival with cisplatin plus pemetrexed (adenocarcinoma: 12.6 vs. 10.9 months). Patients with squamous cell carcinoma had improved survival with the cisplatin plus gemcitabine regimen (10.8 vs. 9.4 months). When compared with the cisplatin plus gemcitabine regimen, the cisplatin plus pemetrexed regimen had significantly lower rates of grade 3 or 4 neutropenia, anemia, and thrombocytopenia ($P \leq .001$); febrile



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

neutropenia ($P = .002$); and alopecia ($P < .001$). Treatment-related deaths were similar for both regimens (cisplatin plus pemetrexed, 9 patients [1.0%]; cisplatin plus gemcitabine, 6 patients [0.7%]). An analysis of three phase 3 trials confirmed that pemetrexed improves survival for patients with nonsquamous NSCLC in first-line, subsequent, and maintenance therapy.¹¹²⁸

Number of Cycles of First-Line Systemic Therapy

Data from the PARAMOUNT trial suggest that 4 cycles of platinum-based therapy is not optimal;⁹⁸⁸ tumors can shrink between 4 to 6 cycles of chemotherapy. However, patients may not be able to tolerate more than 4 cycles of chemotherapy, and most of the maintenance trials used only 4 cycles of chemotherapy.⁸¹⁰ A meta-analysis suggests that continuing the initial regimen beyond 4 to 6 cycles is associated with increased PFS; however, patients have more adverse events.¹¹²⁹ A phase 3 randomized trial suggested that continuing chemotherapy beyond 4 to 6 cycles is not beneficial; however, many patients assigned to a longer duration of therapy did not receive the planned number of cycles.^{807,808} In this phase 3 trial, taxane-based regimens were used and patients had increasing neurotoxicity as more cycles were used.⁸⁰⁷

Many patients with adenocarcinoma receive pemetrexed-based regimens and not taxane-based regimens. Pemetrexed-based regimens are less toxic than taxane-based regimens. Thus, data suggesting that more than 6 cycles of first-line chemotherapy are not appropriate may only apply to taxane-based regimens.⁸¹⁰ Studies report that 60% of patients were able to receive 6 cycles of pemetrexed-based chemotherapy (and had a low incidence of toxicity), whereas only 42% were able to receive more than 5 cycles of taxane-based chemotherapy and often stopped therapy because of neurotoxicity.^{807,990}

The NCCN Guidelines recommend that patients receiving first-line systemic therapy for advanced disease should be evaluated for tumor response with a CT scan. Response assessment should occur after 2 cycles of initial therapy and then every 2 to 4 cycles using CT of known or high-risk sites of disease (with or without contrast) or when clinically indicated.^{265,1130-1132} Patients with responsive or stable disease can continue to receive a total of 4 to 6 cycles of systemic therapy.^{715,807,1133} The NCCN Guidelines do not recommend continuing chemotherapy beyond 4 to 6 cycles. Generally, patients with metastatic NSCLC receive 4 cycles of initial systemic chemotherapy (eg, carboplatin plus pemetrexed plus pembrolizumab for nonsquamous NSCLC) before starting maintenance therapy. However, if patients are tolerating the therapy, then 6 cycles of systemic therapy can be considered. Approximately 25% of patients show disease progression after the initial cycle of chemotherapy; subsequent therapy is recommended for these patients (see the NCCN Guidelines for NSCLC).

Maintenance Therapy

Maintenance therapy is an option for patients with metastatic nonsquamous NSCLC, with responsive or stable disease after first-line systemic chemotherapy or immunotherapy with or without chemotherapy (see the NCCN Guidelines for NSCLC).

A phase 3 randomized trial in 663 patients with advanced NSCLC assessed the effect of best supportive care with (or without) switch maintenance pemetrexed in patients who had received platinum-based chemotherapy but whose tumors had not progressed.⁹⁹⁸ Overall survival was 13.4 months (95% CI, 11.9–15.9) with pemetrexed compared with 10.6 months (95% CI, 8.7–12.0) with placebo (HR, 0.50; 95% CI, 0.42–0.61; $P < .0001$).



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

IUNO, a phase 3 randomized trial, assessed erlotinib as switch maintenance therapy (and as subsequent therapy) for patients with nonsquamous NSCLC and PS 0 to 2 but without *EGFR* mutations.¹⁰⁰⁰ Overall survival and PFS were not improved in patients receiving erlotinib when compared with placebo. The NCCN NSCLC Panel does not recommend erlotinib as switch maintenance therapy (and as subsequent therapy) for patients with nonsquamous NSCLC and PS 0 to 2 but without *EGFR* mutations based on clinical trial data and a revised indication by the FDA.¹⁰⁰⁰

For patients with squamous cell NSCLC, pembrolizumab was used as continuation maintenance therapy if patients received either pembrolizumab plus carboplatin plus either paclitaxel or albumin-bound paclitaxel or they received pembrolizumab alone (see *Immune Checkpoint Inhibitors* in this Discussion). IFCT-GFPC 0502, a phase 3 randomized trial, compared maintenance therapy with either gemcitabine or erlotinib after initial cytotoxic therapy with cisplatin plus gemcitabine in patients with advanced NSCLC.^{743,994} Continuation maintenance therapy with single-agent gemcitabine increased PFS to a greater extent (3.8 months) than switch maintenance therapy with erlotinib (2.9 months) compared with observation (1.9 months).^{743,994} A phase 3 trial assessed switch maintenance therapy with docetaxel given either immediately after chemotherapy or delayed until progression in patients with advanced NSCLC.¹⁰⁰² Many patients in the delayed chemotherapy arm did not receive docetaxel.¹⁰⁰³

Depending on first-line therapy, the NCCN NSCLC Panel recommends continuation maintenance therapy options for patients with nonsquamous NSCLC with atezolizumab, atezolizumab plus bevacizumab (category 1), bevacizumab (category 1), bevacizumab plus pemetrexed, cemiplimab-rwlc with or without pemetrexed (category 1), durvalumab with or without pemetrexed (category 1), gemcitabine (category 2B), nivolumab

plus ipilimumab, pembrolizumab (category 1), pembrolizumab plus pemetrexed (category 1), or pemetrexed (category 1) (see the NCCN Guidelines for NSCLC).^{743,806,812,918,988,992,994} Switch maintenance therapy with pemetrexed is a recommended option for patients with nonsquamous NSCLC.^{743,994,997,998} The benefits of continuation maintenance therapy with gemcitabine were very slight; therefore, the recommendation is only category 2B.⁷⁴³ For patients with squamous cell NSCLC and depending on first-line therapy, the panel recommends continuation maintenance therapy with atezolizumab, cemiplimab-rwlc (category 1), durvalumab, gemcitabine (category 2B), nivolumab plus ipilimumab, or pembrolizumab. Previously, the NCCN Panel deleted the recommendation for docetaxel (category 2B) as switch maintenance therapy for patients with squamous cell NSCLC because there are better options.¹⁰⁰³

Continuation of Targeted Therapy After Progression on Initial Therapy

Patients may continue to derive benefit from EGFR TKIs, ALK inhibitors, or ROS1 inhibitors after disease progression on first-line targeted therapy; discontinuation of these TKIs leads to more rapid progression of disease (symptoms, tumor size, FDG avidity on PET scan) that is termed the *flare phenomenon*.⁸³⁶⁻⁸³⁹ This strategy mirrors the experience in other oncogene-addicted cancers.⁸³⁷ Afatinib, dacomitinib, erlotinib, gefitinib, or osimertinib may be continued after development of acquired resistance in patients with lung adenocarcinoma and *EGFR* exon 19 deletions or *EGFR* exon 21 L858R mutations; local therapy should be considered (eg, SRS to brain metastases or spine, SABR for limited thoracic or other metastatic disease).^{661,1071,1072,1134,1135}

Accumulating data suggest how cancers become resistant to EGFR inhibitors.^{1136,1137} The most common known mechanism is the acquisition of T790M (which is a secondary mutation in *EGFR*), which renders the kinase resistant to afatinib, dacomitinib, erlotinib, or gefitinib.^{1138,1139}



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

Therefore, if patients are T790M positive, osimertinib is recommended (category 1) and afatinib, dacomitinib, erlotinib, or gefitinib are discontinued. Amplification of the *MET* oncogene is another validated resistance mechanism in patients who become resistant to osimertinib.^{1136,1140,1141} Furthermore, data show that when cancers start to progress, which were once sensitive to EGFR inhibitors, discontinuation of the EGFR TKI can lead to a much more accelerated progression of the cancer.^{836,838} Thus, continuing EGFR TKIs is beneficial in many patients even after their cancers develop resistance to EGFR TKIs.¹¹³⁵

The NCCN NSCLC Panel recommends continuing erlotinib (± bevacizumab or ramucirumab), afatinib, dacomitinib, gefitinib, or osimertinib and considering local therapy in patients with asymptomatic progression; however, treatment varies for patients with symptomatic progression (see the NCCN Guidelines for NSCLC).^{1134,1142-1145} Osimertinib is recommended (category 1) for patients with symptomatic brain metastases, T790M positive, and disease progression on erlotinib, gefitinib, dacomitinib, or afatinib.²⁶⁶ Erlotinib (± bevacizumab or ramucirumab), gefitinib, dacomitinib, or afatinib can be continued for these patients with symptomatic brain metastases who are negative for T790M; additional therapy may be added or substituted (eg, local therapy, systemic therapy). First-line systemic therapy options are recommended for patients with multiple symptomatic lesions who are negative for T790M. Osimertinib is recommended (category 1) as subsequent therapy for patients positive for T790M and disease progression on erlotinib, gefitinib, dacomitinib, or afatinib. After progression on osimertinib, patients with *EGFR* exon 19 deletions or *EGFR* exon 21 L858R mutations may continue to derive benefit from osimertinib; other options are also recommended [see *Second-Line and Beyond (Subsequent) Systemic Therapy* in this Discussion]. After progression on alectinib, brigatinib, crizotinib, or ceritinib, patients with *ALK* rearrangements may continue to derive benefit from these agents; other options are also recommended.

Second-Line and Beyond (Subsequent) Systemic Therapy

The phrase *subsequent* therapy is substituted for the terms *second-line*, *third-line*, and *beyond* systemic therapy in the NCCN Guidelines, because the line of therapy may vary depending on previous treatment with targeted agents. Subsequent systemic therapy options for patients with disease progression during or after first-line therapy are described in the NSCLC algorithm and depend on the specific genetic variant, the histologic subtype, whether the patient has symptoms, and PS (see the NCCN Guidelines for NSCLC).¹¹⁴⁶⁻¹¹⁵⁵ The NCCN NSCLC Panel recommends response assessment of known or high-risk sites of disease with CT with or without contrast every 6 to 12 weeks in patients receiving subsequent therapy or maintenance therapy. Note that traditional RECIST response criteria (1.1) are used to assess response for most types of systemic therapy, but different response criteria may be useful for assessing response in patients receiving PD-1 or PD-L1 inhibitors.^{265,1130,1132,1156-1158} If patients have received and completed first-line systemic therapy (such as carboplatin plus paclitaxel), including maintenance therapy, before receiving targeted therapy for an actionable mutation, but have disease progression on targeted therapy, then certain subsequent therapy options are recommended, such as docetaxel.

Response rates to single-agent ICIs vary among patients with different actionable mutations.¹⁸⁶ Data suggest that patients with *EGFR* mutations or *ALK* rearrangements have a low response rate to single-agent ICIs when compared with patients without these mutations (response rate, 3.6% vs. 23%, respectively).^{186,840,1159} Therefore, subsequent therapy with atezolizumab, nivolumab, or pembrolizumab is less effective in patients with *EGFR* mutations or *ALK* rearrangements. Patients with *ALK*-positive NSCLC and very high PD-L1 expression do not respond to pembrolizumab.⁸⁴⁰ In patients with *MET*ex14 mutation-positive metastatic NSCLC, the response rate is about 16% for single-agent ICIs, even those with high PD-L1 expression.^{186,324} Data in the second-line setting suggest



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

that PD-1/PD-L1 inhibitor monotherapy is less effective in patients with metastatic NSCLC who have *EGFR* exon 19 deletions, *EGFR* exon 21 L858R mutations, or *ALK* rearrangements, regardless of PD-L1 expression.^{186,360,945,946,965}

Among patients with *EGFR* exon 19 deletions or *EGFR* exon 21 L858R mutations, no improvement in overall survival has been noted in the phase 3 trials assessing subsequent therapy with atezolizumab, nivolumab, or pembrolizumab compared to docetaxel, but there were not enough patients with these mutations to determine whether there were statistically significant differences.^{360,840,946,958} The PD-1 or PD-L1 inhibitors were not worse than chemotherapy and were better tolerated. In the phase 3 trials for atezolizumab, nivolumab, or pembrolizumab versus docetaxel as subsequent therapy for patients with metastatic NSCLC, subset analyses were done in patients with *EGFR* mutations to determine the best subsequent therapy.^{360,946,958} The HRs for overall survival do not favor docetaxel over atezolizumab (HR, 1.24; CI, 0.7–2.2), nivolumab (HR, 1.18; CI, 0.69–2.0), or pembrolizumab (HR, 0.88; CI, 0.45–1.7); the CIs for the HRs are wide probably because there were so few patients with *EGFR* mutations. The HRs for PFS do favor docetaxel for patients with *EGFR* mutations when compared with either nivolumab (HR, 1.46; CI, 0.90–2.37) or pembrolizumab (HR, 1.79; CI, 0.94–3.42). But again, the CIs are wide. The evidence is weak for recommending docetaxel, atezolizumab, nivolumab, or pembrolizumab as subsequent therapy options for patients with *EGFR* mutations. A meta-analysis suggests that docetaxel improves overall survival when compared with atezolizumab, nivolumab, or pembrolizumab.¹¹⁵⁹

If patients have not previously received an ICI, the NCCN NSCLC Panel recommends (category 1) atezolizumab, nivolumab, or pembrolizumab as preferred subsequent therapy options in eligible patients with metastatic NSCLC based on clinical trial data and FDA approvals (see *Immune*

Checkpoint Inhibitors in this Discussion).^{360,364,945,946,958,965} Human ICI antibodies inhibit the PD-1 receptor or PD-L1, which improves antitumor immunity; PD-1 receptors are expressed on activated cytotoxic T cells.³⁶⁰⁻³⁶²

For patients with *EGFR* exon 19 deletions or *EGFR* exon 21 L858R mutations and disease progression during or after first-line EGFR TKIs (ie, erlotinib [\pm bevacizumab or ramucirumab], afatinib, gefitinib, dacomitinib, or osimertinib therapy), recommended subsequent therapy options depend on whether the progression is asymptomatic or symptomatic and include: 1) considering local therapy; 2) continuing erlotinib (\pm bevacizumab or ramucirumab), afatinib, gefitinib, dacomitinib, or osimertinib; 3) switching to osimertinib if not previously given and T790M positive; or 4) taking a first-line systemic therapy regimen for metastatic NSCLC, such as carboplatin plus paclitaxel.^{246,1143-1145} The NCCN NSCLC Panel recommends osimertinib (category 1) as a subsequent therapy option for patients with T790M-positive metastatic NSCLC, brain metastases, and disease progression on afatinib, dacomitinib, erlotinib (\pm bevacizumab or ramucirumab), or gefitinib.^{266,904,914,915} Data suggest that an afatinib plus cetuximab regimen may be useful for patients who have progressed after receiving erlotinib, afatinib, or gefitinib and after chemotherapy.⁹¹⁷ Patients with T790M-positive and T790M-negative tumors had a similar response rate to an afatinib plus cetuximab regimen (32% vs. 25%; $P = .341$). The NCCN NSCLC Panel recommends considering afatinib plus cetuximab as an option for patients with *EGFR* mutations and disease progression after receiving EGFR TKIs—afatinib, dacomitinib, erlotinib (\pm bevacizumab or ramucirumab), gefitinib, or osimertinib—and after chemotherapy based on these data.

Subsequent therapy is recommended for patients with advanced NSCLC and *EGFR* exon 19 deletions or *EGFR* exon 21 L858R mutations and disease progression during or after first-line therapy with osimertinib.



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

Recommended subsequent therapy options depend on whether the progression is asymptomatic or symptomatic and include: 1) considering definitive local therapy; 2) continuing osimertinib; or 3) switching to a first-line systemic therapy regimen for metastatic NSCLC (eg, carboplatin plus paclitaxel). There are no data to support using erlotinib (\pm bevacizumab or ramucirumab), gefitinib, dacomitinib, or afatinib after progression on osimertinib. Doublet chemotherapy options, such as carboplatin plus paclitaxel, are recommended for eligible patients with metastatic NSCLC, genetic variants, and disease progression after first-line targeted therapy.⁸⁰⁶ Flare phenomenon may occur in some patients who discontinue ALK, EGFR, METex14, RET, or ROS1 inhibitors. If disease flare occurs, then the appropriate TKIs should be restarted.⁸³⁶⁻⁸³⁹ The IMPRESS trial indicated that chemotherapy should be used alone and not be combined with EGFR inhibitors, such as gefitinib, in patients with disease progression on gefitinib.¹¹⁶⁰

For patients with ALK-positive metastatic NSCLC and disease progression during or after first-line targeted therapy with alectinib, brigatinib, ceritinib, or lorlatinib, recommended subsequent therapy options depend on whether the progression is asymptomatic or symptomatic and include: 1) considering definitive local therapy (eg, SABR, SRS, surgery); 2) continuing alectinib, brigatinib, ceritinib, or lorlatinib; 3) switching to lorlatinib for ALK G1202R; or 4) switching to a first-line systemic therapy regimen for metastatic NSCLC, such as carboplatin/paclitaxel. The NCCN NSCLC Panel recommends lorlatinib as a subsequent therapy option for select patients with ALK G1202R metastatic NSCLC and disease progression after treatment with alectinib, brigatinib, or ceritinib based on clinical trial data.⁸³³ After further progression on subsequent targeted therapy, options include: 1) lorlatinib (if not previously given); or 2) first-line combination chemotherapy options for NSCLC (eg, carboplatin plus paclitaxel) (see *Systemic Therapy for Advanced or Metastatic Disease* in the NCCN Guidelines for NSCLC).^{184,1161} The recommended subsequent

therapy options are slightly different if patients have disease progression after initial targeted therapy with crizotinib.

For patients with ROS1-positive metastatic NSCLC and disease progression during or after first-line targeted therapy with ceritinib, crizotinib, or entrectinib, recommended subsequent therapy options depend on whether the progression is asymptomatic or symptomatic and include: 1) considering definitive local therapy (eg, SABR, SRS, surgery); 2) continuing ceritinib, crizotinib, or entrectinib; 3) switching to entrectinib (if previously treated with crizotinib or ceritinib) or lorlatinib (if not previously given); or 4) switching to a first-line systemic therapy regimen for metastatic NSCLC, such as carboplatin plus paclitaxel.

The NCCN NSCLC Panel recommends capmatinib, crizotinib, or tepotinib as subsequent therapy options for select patients with metastatic NSCLC and METex14 skipping mutations who have not previously received any of these agents (see *Oral TKIs that Inhibit METex14 Skipping Mutations and High-Level MET Amplification* in this Discussion). The panel preference stratified the subsequent therapy options for patients with METex14 skipping mutations and voted that capmatinib or tepotinib are preferred options, whereas crizotinib is useful in certain circumstances.

The NCCN NSCLC Panel recommends pralsetinib, selpercatinib, or cabozantinib as subsequent therapy options for select patients with RET rearrangement-positive metastatic NSCLC who have not previously received any of these agents (see *Oral TKIs that Inhibit RET Rearrangements* in this Discussion). The panel preference stratified the subsequent therapy options for patients with RET-positive metastatic NSCLC and voted that pralsetinib or selpercatinib are preferred options, whereas cabozantinib is useful in certain circumstances.

The NCCN NSCLC Panel recommends dabrafenib plus trametinib as a subsequent therapy option for select patients with BRAF p.V600E-positive



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

metastatic NSCLC who have not previously received this regimen (see *Oral Agents that Inhibit BRAF Mutations* in this Discussion). Single-agent dabrafenib or vemurafenib are treatment options if the combination of dabrafenib plus trametinib is not tolerated. The panel recommends entrectinib or larotrectinib as subsequent therapy options for select patients with *NTRK1/2/3* fusion-positive metastatic NSCLC who have not previously received any of these agents (see *Oral TKIs that Inhibit NTRK1/2/3 Gene Fusions* in this Discussion). If patients with *BRAF* p.V600E-, *MET*ex14 skipping-, *NTRK1/2/3*-, or *RET*-positive metastatic NSCLC have disease progression on first-line targeted therapy, then platinum-based doublet therapy options (eg, carboplatin plus paclitaxel) are recommended as subsequent therapy options; note that these platinum-based doublet therapy options are the identical regimens that may be used as first-line combination chemotherapy options for NSCLC, depending on the setting.⁸⁰⁶

Most patients with NSCLC do not have actionable driver mutations [eg, *ALK*, *BRAF* p.V600E, *EGFR*, *ERBB2* (*HER2*), *KRAS*, *MET*ex14 skipping, *NTRK1/2/3*, *RET*, *ROS1*]. Platinum-based doublet therapy is recommended (eg, carboplatin plus paclitaxel) as subsequent systemic therapy options for patients with metastatic NSCLC (but without these genetic variants), positive PD-L1 levels (≥50%), and disease progression after first-line therapy with single-agent ICIs. For patients with metastatic NSCLC and disease progression after first-line therapy with ICIs plus chemotherapy, subsequent systemic therapy options include docetaxel (± ramucirumab), gemcitabine, albumin-bound paclitaxel, or pemetrexed (for nonsquamous only), depending on which agent was not previously given. For patients with all histologic subtypes and PS of 0 to 2, but without these genetic variants, who have disease progression during or after first-line platinum-based combination therapy, recommended subsequent systemic therapy options include single-agent ICIs or chemotherapy. If ICIs have not previously been given, the panel

recommends (category 1) atezolizumab, nivolumab, or pembrolizumab as preferred subsequent therapy options for all histologic subtypes based on improved survival rates, longer duration of response, and fewer adverse events when compared with cytotoxic chemotherapy (see *Immune Checkpoint Inhibitors* in this Discussion).^{360,364,945,946,958,965}

Data show that overall survival is improved with subsequent therapy with atezolizumab, nivolumab, or pembrolizumab compared with docetaxel.^{945,946,965} However, some patients cannot tolerate immunotherapy or have had disease progression on immunotherapy. A phase 3 randomized trial assessed ramucirumab plus docetaxel versus docetaxel as subsequent therapy options for all histologic subtypes.⁹³⁸ Docetaxel has been proven superior to best supportive care, vinorelbine, or ifosfamide with improved survival and quality of life.^{1152,1153} When compared with docetaxel, pemetrexed has similar median survival but less toxicity.^{1154,1162} Pemetrexed is a recommended subsequent therapy option for eligible patients with metastatic nonsquamous NSCLC.⁹⁹⁸ Docetaxel is recommended for patients with wild-type *EGFR* tumors based on two randomized trials comparing erlotinib versus docetaxel.^{1163,1164} Patients often have a limited response to subsequent therapy other than ICIs, although chemotherapy may serve a useful palliative role.¹¹⁶⁵

ABOUND, a phase 2 trial, assessed albumin-bound paclitaxel with or without oral 5-azacitidine as subsequent therapy in 161 patients with advanced nonsquamous NSCLC.¹¹⁶⁶ Median overall survival was 8.1 months with combination therapy and 17 months with single-agent albumin-bound paclitaxel (HR, 1.7; 95% CI, 1.08–2.57). Grade 3 or greater adverse events were reported in 41% of patients receiving combination therapy and 32% of single-agent albumin-bound paclitaxel.

The NSCLC Panel recommends the following subsequent therapy options for patients with metastatic NSCLC, depending on which agents have not previously been given: 1) atezolizumab, nivolumab, or pembrolizumab if



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

none has been previously given (all are category 1, preferred); 2) docetaxel (\pm ramucirumab); 3) gemcitabine; 4) albumin-bound paclitaxel; or 5) pemetrexed (nonsquamous only). In patients with PS of 3 to 4, best supportive care is recommended (see the NCCN Guidelines for NSCLC).^{6,722,723} After second disease progression, subsequent therapy is recommended for certain patients if the following agents have not already been given: 1) atezolizumab, nivolumab, or pembrolizumab if none has been previously given; 2) docetaxel (\pm ramucirumab; category 2B for both); 3) gemcitabine (category 2B); 4) albumin-bound paclitaxel (category 2B); or 5) pemetrexed (nonsquamous only) (category 2B).^{1147,1164,1167,1168} These patients include those with advanced NSCLC and a PS of 0 to 2.

UNO, a randomized trial, showed that overall survival and PFS were not improved in patients receiving erlotinib as a subsequent therapy option when compared with placebo.¹⁰⁰⁰ The NCCN NSCLC Panel does not recommend erlotinib as a subsequent therapy option (or as switch maintenance therapy) for patients with nonsquamous NSCLC and PS of 0 to 2 but without *EGFR* mutations based on clinical trial data and a revised indication by the FDA.¹⁰⁰⁰ A phase 3 randomized trial compared afatinib with erlotinib as subsequent therapy options for patients with squamous cell NSCLC.⁸⁷⁸ Overall survival was 7.9 months (95% CI, 7.2–8.7) for afatinib versus 6.8 months (95% CI, 5.9–7.8) for erlotinib (HR, 0.81; 95% CI, 0.69–0.95; $P = .0077$). Almost 60% of patients in each arm had grade 3 or higher adverse events. In contrast, the median overall survival was 9.2 months with nivolumab compared with 6.0 months for docetaxel for patients with squamous cell NSCLC.³⁶⁴ In addition, only 7% of patients receiving nivolumab had grade 3 or higher adverse events. The NCCN NSCLC Panel does not recommend erlotinib as a subsequent therapy option for patients with squamous cell NSCLC based on this study comparing afatinib with erlotinib; this study was statistically significant but not clinically significant.⁸⁷⁸ Erlotinib or afatinib are not recommended as

subsequent therapy options for patients with squamous cell NSCLC; they are less efficacious and safe compared to other available options.⁸⁷⁸

Summary

Management of NSCLC is described in the NCCN Guidelines for NSCLC, which include the algorithm and this supporting Discussion text. Revisions for the 2023 update of the NCCN Guidelines for NSCLC are described in this Discussion and outlined in the algorithm (see *Summary of the Guidelines Updates* in the algorithm). Most lung cancer (approximately 85%) is due to NSCLC; SCLC accounts for about 15% of lung cancer. Recommendations for SCLC are provided in a different guideline (see the NCCN Guidelines for Small Cell Lung Cancer, available at www.NCCN.org). The NCCN Guidelines for NSCLC are updated at least once a year by the NCCN NSCLC Panel; there were 6 updates to the 2022 guidelines.

For the 2023 update (Version 1), the NCCN NSCLC Panel revised the *Principles of Molecular and Biomarker Analysis*. For example, the NCCN Panel added testing recommendations for *ERBB2* (*HER2*) mutations. NGS-based approaches are best able to detect the broad spectrum of genomic *ERBB2* (*HER2*) alterations that may occur, although Sanger sequencing and targeted PCR approaches may also be used. The NCCN Panel also revised the *Principles of Diagnostic Evaluation* in the NCCN Guidelines for NSCLC. For example, the NCCN Panel clarified that a preoperative biopsy may be useful for patients with early-stage NSCLC (ie, clinical stage IB or more) who may be candidates for systemic therapy before surgery (also known as neoadjuvant, induction, or preoperative therapy).¹¹⁰ The NCCN Panel added a recommendation that patients should be evaluated for perioperative (neoadjuvant or adjuvant) therapy before surgery. Neoadjuvant therapy should not be used to attempt to induce resectability in patients who do not already meet criteria for resectability on initial evaluation.



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

For the 2023 update (Version 2), the NCCN Panel added a recommendation for adjuvant pembrolizumab for eligible patients with completely resected (R0) stage IIB to IIIA, stage IIIB (only T3,N2), or high-risk stage IIA NSCLC who are negative for *EGFR* exon 19 deletions, *EGFR* exon 21 L858R mutations, or *ALK* fusions and who have previously received adjuvant chemotherapy.³² The panel added a caveat that the benefit of adjuvant pembrolizumab is unclear for patients with PD-L1 levels less than 1%. The NCCN Panel also expanded the molecular testing criteria to include *ALK* rearrangements because patients with completely resected (R0) NSCLC who have *ALK* rearrangements, *EGFR* exon 19 deletions, or *EGFR* exon 21 L858R mutations have less benefit from ICIs.¹⁸⁶ Single-agent therapy with osimertinib is recommended for eligible patients with completely resected (R0) stage IB–IIIA or stage IIIB (T3, N2) NSCLC if patients have *EGFR* exon 19 deletions or *EGFR* exon 21 L858R mutations, regardless of PD-L1 levels. The NCCN Panel clarified that adjuvant chemotherapy followed by single-agent therapy with osimertinib is recommended for eligible patients with completely resected (R0) early-stage NSCLC if patients have both *EGFR* mutations and PD-L1 levels of 1% or more; the panel does not recommend using atezolizumab or pembrolizumab, or combined therapy with osimertinib and either atezolizumab or pembrolizumab in this setting.

For the 2023 update (Version 1), the NCCN NSCLC Panel expanded the criteria for using adjuvant osimertinib, atezolizumab, or pembrolizumab for eligible patients with completely resected (R0) early-stage NSCLC to include certain patients with stage IIIB (only T3,N2) disease based on differences between the AJCC staging systems for the 7th and 8th editions.¹⁶² Although the 8th edition of the AJCC staging manual is currently being used, the clinical trials used the 7th edition.

For the 2023 update (Version 1), the NCCN NSCLC Panel recommends adagrasib as a subsequent therapy option for select patients with

metastatic NSCLC and *KRAS* p.G12C mutations who have disease progression after treatment with platinum-based chemotherapy (± immunotherapy) based on clinical trial data and FDA approval.³¹⁷

Responsiveness to adagrasib has not been assessed for mutations other than *KRAS* p.G12C. The NCCN Panel clarified that adagrasib is recommended as third-line or beyond therapy if the patient has not previously received targeted therapy for *KRAS* p.G12C mutations. The NCCN Panel also recommends lorlatinib as a subsequent therapy option, if not previously given, for patients with *ROS1*-positive metastatic NSCLC and symptomatic brain metastases.^{935,936}

For the 2023 update (Version 1), the NCCN Panel clarified that postoperative RT may be considered for select patients with high-risk N2 disease such as extracapsular extension, multistation involvement, inadequate lymph node dissection or sampling, and/or refusal or intolerance of adjuvant systemic therapy.^{452,455,767} The Panel revised the algorithm for multiple lung cancers. For example, the NCCN Panel added an initial recommendation for multidisciplinary evaluation for patients suspected of or confirmed with having multiple lung cancers. One of the goals during multidisciplinary evaluation for multiple lung cancers is to assess whether patients have lung nodules that can be observed instead of erroneously assuming that patients have stage IV NSCLC. The panel also added the following caveats: lesions at low risk of becoming symptomatic can be observed, such as small subsolid nodules with slow growth. However, treatment should be considered if the lesion(s) show accelerating growth, increasing solid component, or increasing FDG uptake, even if the lesions are small.



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

References

1. Siegel RL, Miller KD, Wagle NS, Jemal A. Cancer statistics, 2023. CA Cancer J Clin 2023;73:17-48. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/36633525>.
2. Torre LA, Siegel RL, Jemal A. Lung Cancer Statistics. Adv Exp Med Biol 2016;893:1-19. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26667336>.
3. Siegel RL, Miller KD, Fuchs HE, Jemal A. Cancer statistics, 2022. CA Cancer J Clin 2022;72:7-33. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/35020204>.
4. SEER*Explorer: An interactive website for SEER cancer statistics [Internet]. Surveillance Research Program [Cited 2022 October 4] National Cancer Institute; 2022. Available at: <https://seer.cancer.gov/statistics-network/explorer/>.
5. Howlader N, Noone AM, Krapcho M, et al. SEER Cancer Statistics Review, 1975-2017, based on November 2019 SEER data submission, posted to the SEER web site, April 2020. Bethesda, MD: National Cancer Institute; 2020. Available at: https://seer.cancer.gov/csr/1975_2017/.
6. Simoff MJ, Lally B, Slade MG, et al. Symptom management in patients with lung cancer: Diagnosis and management of lung cancer, 3rd ed: American College of Chest Physicians evidence-based clinical practice guidelines. Chest 2013;143:e455S-e497S. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23649452>.
7. Brahmer JR, Govindan R, Anders RA, et al. The Society for Immunotherapy of Cancer consensus statement on immunotherapy for the treatment of non-small cell lung cancer (NSCLC). J Immunother Cancer 2018;6:75. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30012210>.
8. Johnson DH, Schiller JH, Bunn PA, Jr. Recent clinical advances in lung cancer management. J Clin Oncol 2014;32:973-982. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24567433>.
9. Reck M, Heigener DF, Mok T, et al. Management of non-small-cell lung cancer: recent developments. Lancet 2013;382:709-719. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23972814>.
10. Forde PM, Ettinger DS. Targeted therapy for non-small-cell lung cancer: past, present and future. Expert Rev Anticancer Ther 2013;13:745-758. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23773106>.
11. Ettinger DS. Ten years of progress in non-small cell lung cancer. J Natl Compr Canc Netw 2012;10:292-295. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22393190>.
12. Reck M, Rodriguez-Abreu D, Robinson AG, et al. Updated analysis of KEYNOTE-024: pembrolizumab versus platinum-based chemotherapy for advanced non-small-cell lung cancer with PD-L1 tumor proportion score of 50% or greater. J Clin Oncol 2019;37:537-546. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30620668>.
13. Siegel RL, Miller KD, Fuchs HE, Jemal A. Cancer statistics, 2021. CA Cancer J Clin 2021;71:7-33. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/33433946>.
14. Herbst RS, Garon EB, Kim DW, et al. Five year survival update from KEYNOTE-010: pembrolizumab versus docetaxel for previously treated, programmed death-ligand 1-positive advanced NSCLC. J Thorac Oncol 2021;16:1718-1732. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/34048946>.
15. Reck M, Rodriguez-Abreu D, Robinson AG, et al. Five-year outcomes with pembrolizumab versus chemotherapy for metastatic non-small-cell lung cancer with PD-L1 tumor proportion score \geq 50. J Clin Oncol 2021;39:2339-2349. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/33872070>.
16. Spigel DR, Faivre-Finn C, Gray JE, et al. Five-year survival outcomes from the PACIFIC trial: durvalumab after chemoradiotherapy in stage III non-small-cell lung cancer. J Clin Oncol 2022;40:1301-1311. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/35108059>.



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

17. Mok T, Camidge DR, Gadgeel SM, et al. Updated overall survival and final progression-free survival data for patients with treatment-naïve advanced ALK-positive non-small-cell lung cancer in the ALEX study. *Ann Oncol* 2020;31:1056-1064. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32418886>.
18. Planchard D, Besse B, Groen HJM, et al. Phase 2 Study of Dabrafenib Plus Trametinib in Patients With BRAF V600E-Mutant Metastatic NSCLC: Updated 5-Year Survival Rates and Genomic Analysis. *J Thorac Oncol* 2022;17:103-115. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/34455067>.
19. Faivre-Finn C, Vicente D, Kurata T, et al. Four-Year Survival With Durvalumab After Chemoradiotherapy in Stage III NSCLC—an Update From the PACIFIC Trial. *J Thorac Oncol* 2021;16:860-867. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/33476803>.
20. Borghaei H, Gettinger S, Vokes EE, et al. Five-Year Outcomes From the Randomized, Phase III Trials CheckMate 017 and 057: Nivolumab Versus Docetaxel in Previously Treated Non-Small-Cell Lung Cancer. *J Clin Oncol* 2021;39:723-733. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/33449799>.
21. Ramalingam SS, Vansteenkiste J, Planchard D, et al. Overall Survival with Osimertinib in Untreated, EGFR-Mutated Advanced NSCLC. *N Engl J Med* 2020;382:41-50. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31751012>.
22. Garon EB, Hellmann MD, Rizvi NA, et al. Five-Year Overall Survival for Patients With Advanced NonSmall-Cell Lung Cancer Treated With Pembrolizumab: Results From the Phase I KEYNOTE-001 Study. *J Clin Oncol* 2019;37:2518-2527. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31154919>.
23. Leighl NB, Hellmann MD, Hui R, et al. Pembrolizumab in patients with advanced non-small-cell lung cancer (KEYNOTE-001): 3-year results from an open-label, phase 1 study. *Lancet Respir Med* 2019;7:347-357. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30876831>.
24. Pacheco JM, Gao D, Smith D, et al. Natural History and Factors Associated with Overall Survival in Stage IV ALK-Rearranged Non-Small Cell Lung Cancer. *J Thorac Oncol* 2019;14:691-700. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30599201>.
25. Shaw AT, Riely GJ, Bang YJ, et al. Crizotinib in ROS1-rearranged advanced non-small-cell lung cancer (NSCLC): updated results, including overall survival, from PROFILE 1001. *Ann Oncol* 2019;30:1121-1126. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30980071>.
26. Lin JJ, Cardarella S, Lydon CA, et al. Five-Year Survival in EGFR-Mutant Metastatic Lung Adenocarcinoma Treated with EGFR-TKIs. *J Thorac Oncol* 2016;11:556-565. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26724471>.
27. Singhi EK, Horn L, Sequist LV, et al. Advanced Non-Small Cell Lung Cancer: Sequencing Agents in the EGFR-Mutated/ALK-Rearranged Populations. *Am Soc Clin Oncol Educ Book* 2019;39:e187-e197. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31099642>.
28. Antonia SJ, Borghaei H, Ramalingam SS, et al. Four-year survival with nivolumab in patients with previously treated advanced non-small-cell lung cancer: a pooled analysis. *Lancet Oncol* 2019;20:1395-1408. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31422028>.
29. Zhao D, Chen X, Qin N, et al. The prognostic role of EGFR-TKIs for patients with advanced non-small cell lung cancer. *Sci Rep* 2017;7:40374. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28079142>.
30. Johung KL, Yeh N, Desai NB, et al. Extended Survival and Prognostic Factors for Patients With ALK-Rearranged Non-Small-Cell Lung Cancer and Brain Metastasis. *J Clin Oncol* 2016;34:123-129. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26438117>.
31. Ettinger DS, Cox JD, Ginsberg RJ, et al. NCCN Non-Small-Cell Lung Cancer Practice Guidelines. The National Comprehensive Cancer Network. *Oncology (Williston Park)* 1996;10:81-111. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/8953597>.



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

32. O'Brien M, Paz-Ares L, Marreaud S, et al. Pembrolizumab versus placebo as adjuvant therapy for completely resected stage IB-IIIA non-small-cell lung cancer (PEARLS/KEYNOTE-091): an interim analysis of a randomised, triple-blind, phase 3 trial. *Lancet Oncol* 2022;23:1274-1286. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/36108662>.

33. Alberg AJ, Brock MV, Ford JG, et al. Epidemiology of lung cancer: Diagnosis and management of lung cancer, 3rd ed: American College of Chest Physicians evidence-based clinical practice guidelines. *Chest* 2013;143:e1S-e29S. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23649439>.

34. Subramanian J, Govindan R. Lung cancer in never smokers: a review. *J Clin Oncol* 2007;25:561-570. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17290066>.

35. The health consequences of smoking: A report of the Surgeon General. (ed 2010/07/30). Atlanta (GA): U.S. Department of Health and Human Services. Centers for Disease Control and Prevention (US); 2004.

36. Secretan B, Straif K, Baan R, et al. A review of human carcinogens--Part E: tobacco, areca nut, alcohol, coal smoke, and salted fish. *Lancet Oncol* 2009;10:1033-1034. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19891056>.

37. Doll R, Peto R. Mortality in relation to smoking: 20 years' observations on male British doctors. *Br Med J* 1976;2:1525-1536. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/1009386>.

38. Taylor R, Najafi F, Dobson A. Meta-analysis of studies of passive smoking and lung cancer: effects of study type and continent. *Int J Epidemiol* 2007;36:1048-1059. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17690135>.

39. The health consequences of involuntary exposure to tobacco smoke: A report of the Surgeon General. (ed 2010/07/30). Atlanta (GA): U.S. Department of Health and Human Services, Centers for Disease Control and Prevention; 2006.

40. Hackshaw AK, Law MR, Wald NJ. The accumulated evidence on lung cancer and environmental tobacco smoke. *BMJ* 1997;315:980-988. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/9365295>.

41. Wald NJ, Nanchahal K, Thompson SG, Cuckle HS. Does breathing other people's tobacco smoke cause lung cancer? *Br Med J (Clin Res Ed)* 1986;293:1217-1222. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/3096439>.

42. Fraumeni JF, Jr. Respiratory carcinogenesis: an epidemiologic appraisal. *J Natl Cancer Inst* 1975;55:1039-1046. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/1107567>.

43. Janerich DT, Thompson WD, Varela LR, et al. Lung cancer and exposure to tobacco smoke in the household. *N Engl J Med* 1990;323:632-636. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/2385268>.

44. Straif K, Benbrahim-Tallaa L, Baan R, et al. A review of human carcinogens--Part C: metals, arsenic, dusts, and fibres. *Lancet Oncol* 2009;10:453-454. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19418618>.

45. Driscoll T, Nelson DI, Steenland K, et al. The global burden of disease due to occupational carcinogens. *Am J Ind Med* 2005;48:419-431. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/16299703>.

46. Humans IWGoTEoCRt. Arsenic, metals, fibres, and dusts. *IARC Monogr Eval Carcinog Risks Hum* 2012;100:11-465. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23189751>.

47. Omenn GS, Merchant J, Boatman E, et al. Contribution of environmental fibers to respiratory cancer. *Environ Health Perspect* 1986;70:51-56. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/3830113>.

48. Chlebowski RT, Schwartz AG, Wakelee H, et al. Oestrogen plus progestin and lung cancer in postmenopausal women (Women's Health Initiative trial): a post-hoc analysis of a randomised controlled trial. *Lancet*



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

2009;374:1243-1251. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/19767090>.

49. Chlebowski RT, Anderson GL, Manson JE, et al. Lung cancer among postmenopausal women treated with estrogen alone in the women's health initiative randomized trial. *J Natl Cancer Inst* 2010;102:1413-1421.

Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20709992>.

50. Thun MJ, Carter BD, Feskanich D, et al. 50-year trends in smoking-related mortality in the United States. *N Engl J Med* 2013;368:351-364. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/23343064>.

51. Shiels MS, Gibson T, Sampson J, et al. Cigarette smoking prior to first cancer and risk of second smoking-associated cancers among survivors of bladder, kidney, head and neck, and stage I lung cancers. *J Clin Oncol* 2014;32:3989-3995. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/25385740>.

52. Leone FT, Evers-Casey S, Toll BA, Vachani A. Treatment of tobacco use in lung cancer: Diagnosis and management of lung cancer, 3rd ed: American College of Chest Physicians evidence-based clinical practice guidelines. *Chest* 2013;143:e61S-e77S. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/23649454>.

53. Jha P, Ramasundarahettige C, Landsman V, et al. 21st-century hazards of smoking and benefits of cessation in the United States. *N Engl J Med* 2013;368:341-350. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/23343063>.

54. Rigotti NA. Strategies to help a smoker who is struggling to quit. *JAMA* 2012;308:1573-1580. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/23073954>.

55. Five major steps to intervention (the "5 A's"). Vol. December. Rockville, MD: Agency for Healthcare Research and Quality; 2012.

Available at: <http://bit.ly/1jXzEvC>.

56. Tao L, Wang R, Gao YT, Yuan JM. Impact of postdiagnosis smoking on long-term survival of cancer patients: the Shanghai cohort study. *Cancer Epidemiol Biomarkers Prev* 2013;22:2404-2411. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/24319070>.

57. Marino KA, Little MA, Bursac Z, et al. Operating on Patients Who Smoke: A Survey of Thoracic Surgeons in the United States. *Ann Thorac Surg* 2016;102:911-916. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/27474514>.

58. Treating tobacco use and dependence. Vol. April. Rockville, MD: Agency for Healthcare Research and Quality; 2013. Available at:

<http://bit.ly/28KMo4K>.

59. Cahill K, Stevens S, Perera R, Lancaster T. Pharmacological interventions for smoking cessation: an overview and network meta-analysis. *Cochrane Database Syst Rev* 2013;5:CD009329. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/23728690>.

60. Koegelenberg CF, Noor F, Bateman ED, et al. Efficacy of varenicline combined with nicotine replacement therapy vs varenicline alone for smoking cessation: a randomized clinical trial. *JAMA* 2014;312:155-161. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/25005652>.

61. Walker N, Howe C, Glover M, et al. Cytisine versus nicotine for smoking cessation. *N Engl J Med* 2014;371:2353-2362. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/25517706>.

62. Aubin HJ, Bobak A, Britton JR, et al. Varenicline versus transdermal nicotine patch for smoking cessation: results from a randomised open-label trial. *Thorax* 2008;63:717-724. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/18263663>.

63. Jorenby DE, Hays JT, Rigotti NA, et al. Efficacy of varenicline, an alpha4beta2 nicotinic acetylcholine receptor partial agonist, vs placebo or sustained-release bupropion for smoking cessation: a randomized controlled trial. *JAMA* 2006;296:56-63. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/16820547>.



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

64. Gonzales D, Rennard SI, Nides M, et al. Varenicline, an alpha4beta2 nicotinic acetylcholine receptor partial agonist, vs sustained-release bupropion and placebo for smoking cessation: a randomized controlled trial. JAMA 2006;296:47-55. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/16820546>.

65. Cahill K, Stead LF, Lancaster T. Nicotine receptor partial agonists for smoking cessation. Cochrane Database Syst Rev 2011:CD006103. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21328282>.

66. Ware JH, Vetrovec GW, Miller AB, et al. Cardiovascular safety of varenicline: patient-level meta-analysis of randomized, blinded, placebo-controlled trials. Am J Ther 2013;20:235-246. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23615317>.

67. Haber SL, Boomers V, Raney E. Safety of varenicline in patients with cardiovascular disease. J Pharm Pract 2014;27:65-70. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24080536>.

68. Mills EJ, Thorlund K, Eapen S, et al. Cardiovascular events associated with smoking cessation pharmacotherapies: a network meta-analysis. Circulation 2014;129:28-41. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24323793>.

69. Xi ZX. Preclinical Pharmacology, Efficacy and Safety of Varenicline in Smoking Cessation and Clinical Utility in High Risk Patients. Drug Healthc Patient Saf 2010;2010:39-48. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21278851>.

70. Gonzales D, Hajek P, Pliamm L, et al. Retreatment with varenicline for smoking cessation in smokers who have previously taken varenicline: a randomized, placebo-controlled trial. Clin Pharmacol Ther 2014;96:390-396. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24911368>.

71. Garrison GD, Dugan SE. Varenicline: a first-line treatment option for smoking cessation. Clin Ther 2009;31:463-491. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19393839>.

72. Hays JT, Ebbert JO. Adverse effects and tolerability of medications for the treatment of tobacco use and dependence. Drugs 2010;70:2357-2372. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21142259>.

73. Carney DN. Lung cancer--time to move on from chemotherapy. N Engl J Med 2002;346:126-128. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/11784881>.

74. Chute JP, Chen T, Feigal E, et al. Twenty years of phase III trials for patients with extensive-stage small-cell lung cancer: perceptible progress. J Clin Oncol 1999;17:1794-1801. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/10561217>.

75. National Lung Screening Trial Research T, Aberle DR, Berg CD, et al. The National Lung Screening Trial: overview and study design. Radiology 2011;258:243-253. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21045183>.

76. National Lung Screening Trial Research T, Aberle DR, Adams AM, et al. Reduced lung-cancer mortality with low-dose computed tomographic screening. N Engl J Med 2011;365:395-409. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21714641>.

77. National Lung Screening Trial Research T, Aberle DR, Adams AM, et al. Baseline characteristics of participants in the randomized national lung screening trial. J Natl Cancer Inst 2010;102:1771-1779. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21119104>.

78. de Koning HJ, van der Aalst CM, de Jong PA, et al. Reduced lung-cancer mortality with volume CT screening in a randomized trial. N Engl J Med 2020;382:503-513. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31995683>.

79. Krist AH, Davidson KW, Mangione CM, et al. Screening for lung cancer: US Preventive Services Task Force Recommendation Statement. JAMA 2021;325:962-970. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/33687470>.



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

80. Detterbeck FC, Mazzone PJ, Naidich DP, Bach PB. Screening for lung cancer: Diagnosis and management of lung cancer, 3rd ed: American College of Chest Physicians evidence-based clinical practice guidelines. Chest 2013;143:e78S-e92S. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23649455>.

81. Vansteenkiste J, Crino L, Doms C, et al. 2nd ESMO Consensus Conference on Lung Cancer: early-stage non-small-cell lung cancer consensus on diagnosis, treatment and follow-up. Ann Oncol 2014;25:1462-1474. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24562446>.

82. Smith RA, Brooks D, Cokkinides V, et al. Cancer screening in the United States, 2013: a review of current American Cancer Society guidelines, current issues in cancer screening, and new guidance on cervical cancer screening and lung cancer screening. CA Cancer J Clin 2013;63:88-105. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23378235>.

83. Moyer VA, Force USPST. Screening for lung cancer: U.S. Preventive Services Task Force recommendation statement. Ann Intern Med 2014;160:330-338. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24378917>.

84. Tsao MS, Nicholson AG, Maleszewski JJ, et al. Introduction to 2021 WHO Classification of Thoracic Tumors. J Thorac Oncol 2022;17:e1-e4. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/34930611>.

85. WHO Classification of Tumours Editorial Board. WHO Classification of Tumours Thoracic Tumours. WHO classification of tumours series, 5th ed. Lyon, France: International Agency for Research on Cancer; 2021.

86. Travis WD, Brambilla E, Nicholson AG, et al. The 2015 World Health Organization Classification of Lung Tumors: Impact of Genetic, Clinical and Radiologic Advances Since the 2004 Classification. J Thorac Oncol 2015;10:1243-1260. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26291008>.

87. Travis WD, Brambilla E, Burke AP, et al. WHO Classification of Tumours of the Lung, Pleura, Thymus and Heart, Volume 7. Lyon: International Agency for Research on Cancer; 2015:412.

88. Howlader N, Noone AM, Krapcho M, et al. SEER Cancer Statistics Review, 1975-2018, based on November 2020 SEER data submission, posted to the SEER web site, April 2021. Bethesda, MD: National Cancer Institute; 2021. Available at: https://seer.cancer.gov/csr/1975_2018/.

89. Travis WD, Brambilla E, Noguchi M, et al. International association for the study of lung cancer/american thoracic society/european respiratory society international multidisciplinary classification of lung adenocarcinoma. J Thorac Oncol 2011;6:244-285. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21252716>.

90. Finkelstein DM, Ettinger DS, Ruckdeschel JC. Long-term survivors in metastatic non-small-cell lung cancer: an Eastern Cooperative Oncology Group Study. J Clin Oncol 1986;4:702-709. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/3701389>.

91. Tammemagi MC, Church TR, Hocking WG, et al. Evaluation of the lung cancer risks at which to screen ever- and never-smokers: screening rules applied to the PLCO and NLST cohorts. PLoS Med 2014;11:e1001764. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25460915>.

92. Gould MK, Donington J, Lynch WR, et al. Evaluation of individuals with pulmonary nodules: when is it lung cancer? Diagnosis and management of lung cancer, 3rd ed: American College of Chest Physicians evidence-based clinical practice guidelines. Chest 2013;143:e93S-e120S. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23649456>.

93. American College of Radiology. lung CT screening reporting and data system (Lung-RADS); 2016. Available at: <https://www.acr.org/Quality-Safety/Resources/LungRADS>.

94. MacMahon H, Naidich DP, Goo JM, et al. Guidelines for Management of Incidental Pulmonary Nodules Detected on CT Images: From the



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

Fleischner Society 2017. Radiology 2017;284:228-243. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28240562>.

95. MacMahon H, Austin JH, Gamsu G, et al. Guidelines for management of small pulmonary nodules detected on CT scans: a statement from the Fleischner Society. Radiology 2005;237:395-400. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/16244247>.

96. Naidich DP, Bankier AA, MacMahon H, et al. Recommendations for the management of subsolid pulmonary nodules detected at CT: a statement from the Fleischner Society. Radiology 2013;266:304-317. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23070270>.

97. Blagev DP, Lloyd JF, Conner K, et al. Follow-up of incidental pulmonary nodules and the radiology report. J Am Coll Radiol 2014;11:378-383. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24316231>.

98. Gould MK, Tang T, Liu IL, et al. Recent Trends in the Identification of Incidental Pulmonary Nodules. Am J Respir Crit Care Med 2015;192:1208-1214. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26214244>.

99. Gardiner N, Jogai S, Wallis A. The revised lung adenocarcinoma classification-an imaging guide. J Thorac Dis 2014;6:S537-546. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25349704>.

100. Seidelman JL, Myers JL, Quint LE. Incidental, subsolid pulmonary nodules at CT: etiology and management. Cancer Imaging 2013;13:365-373. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24061063>.

101. Hansell DM, Bankier AA, MacMahon H, et al. Fleischner Society: glossary of terms for thoracic imaging. Radiology 2008;246:697-722. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/18195376>.

102. Yankelevitz DF, Yip R, Smith JP, et al. CT Screening for Lung Cancer: Nonsolid Nodules in Baseline and Annual Repeat Rounds.

Radiology 2015;277:555-564. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26101879>.

103. Travis WD, Brambilla E, Noguchi M, et al. Diagnosis of lung cancer in small biopsies and cytology: implications of the 2011 International Association for the Study of Lung Cancer/American Thoracic Society/European Respiratory Society classification. Arch Pathol Lab Med 2013;137:668-684. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22970842>.

104. Kim HY, Shim YM, Lee KS, et al. Persistent pulmonary nodular ground-glass opacity at thin-section CT: histopathologic comparisons. Radiology 2007;245:267-275. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17885195>.

105. Marshall HM, Bowman RV, Yang IA, et al. Screening for lung cancer with low-dose computed tomography: a review of current status. J Thorac Dis 2013;5 Suppl 5:S524-539. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24163745>.

106. Brawley OW, Flanagan EL. Low-dose spiral CT screening and evaluation of the solitary pulmonary nodule. Oncology (Williston Park) 2014;28:441-446. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25004661>.

107. Pinsky PF, Gierada DS, Black W, et al. Performance of Lung-RADS in the National Lung Screening Trial: a retrospective assessment. Ann Intern Med 2015;162:485-491. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25664444>.

108. McKee BJ, Regis SM, McKee AB, et al. Performance of ACR Lung-RADS in a clinical CT lung screening program. J Am Coll Radiol 2015;12:273-276. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25176499>.

109. Kazerooni EA, Austin JH, Black WC, et al. ACR-STR practice parameter for the performance and reporting of lung cancer screening thoracic computed tomography (CT): 2014 (Resolution 4). J Thorac



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

Imaging 2014;29:310-316. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/24992501>.

110. Forde PM, Spicer J, Lu S, et al. Neoadjuvant nivolumab plus chemotherapy in resectable lung cancer. *N Engl J Med* 2022;386:1973-1985. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/35403841>.

111. Rivera MP, Mehta AC, Wahidi MM. Establishing the diagnosis of lung cancer: Diagnosis and management of lung cancer, 3rd ed: American College of Chest Physicians evidence-based clinical practice guidelines. *Chest* 2013;143:e142S-e165S. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/23649436>.

112. Silvestri GA, Gonzalez AV, Jantz MA, et al. Methods for staging non-small cell lung cancer: Diagnosis and management of lung cancer, 3rd ed: American College of Chest Physicians evidence-based clinical practice guidelines. *Chest* 2013;143:e211S-e250S. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/23649440>.

113. Schwartz AM, Rezaei MK. Diagnostic surgical pathology in lung cancer: Diagnosis and management of lung cancer, 3rd ed: American College of Chest Physicians evidence-based clinical practice guidelines. *Chest* 2013;143:e251S-e262S. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/23649441>.

114. Li MM, Datto M, Duncavage EJ, et al. Standards and Guidelines for the Interpretation and Reporting of Sequence Variants in Cancer: A Joint Consensus Recommendation of the Association for Molecular Pathology, American Society of Clinical Oncology, and College of American Pathologists. *J Mol Diagn* 2017;19:4-23. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/27993330>.

115. Leichsenring J, Horak P, Kreutzfeldt S, et al. Variant classification in precision oncology. *Int J Cancer* 2019;145:2996-3010. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/31008532>.

116. Dy GK, Nesline MK, Papanicolaou-Sengos A, et al. Treatment recommendations to cancer patients in the context of FDA guidance for

next generation sequencing. *BMC Med Inform Decis Mak* 2019;19:14.

Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30658646>.

117. Jordan EJ, Kim HR, Arcila ME, et al. Prospective Comprehensive Molecular Characterization of Lung Adenocarcinomas for Efficient Patient Matching to Approved and Emerging Therapies. *Cancer Discov* 2017;7:596-609. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/28336552>.

118. Oxnard GR, Binder A, Janne PA. New targetable oncogenes in non-small-cell lung cancer. *J Clin Oncol* 2013;31:1097-1104. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/23401445>.

119. Cooper WA, O'Toole S, Boyer M, et al. What's new in non-small cell lung cancer for pathologists: the importance of accurate subtyping, EGFR mutations and ALK rearrangements. *Pathology* 2011;43:103-115.

Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21233671>.

120. Fossella FV, Putnam JB, Komaki R, eds. *Lung Cancer*. M.D. Anderson Cancer Care Series. New York: Springer; 2003:316.

121. Eberhard DA, Johnson BE, Amler LC, et al. Mutations in the epidermal growth factor receptor and in KRAS are predictive and prognostic indicators in patients with non-small-cell lung cancer treated with chemotherapy alone and in combination with erlotinib. *J Clin Oncol* 2005;23:5900-5909. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/16043828>.

122. Cappuzzo F, Ligorio C, Toschi L, et al. EGFR and HER2 gene copy number and response to first-line chemotherapy in patients with advanced non-small cell lung cancer (NSCLC). *J Thorac Oncol* 2007;2:423-429.

Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17473658>.

123. Kwak EL, Bang YJ, Camidge DR, et al. Anaplastic lymphoma kinase inhibition in non-small-cell lung cancer. *N Engl J Med* 2010;363:1693-1703. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/20979469>.



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

124. Zakowski MF, Rekhtman N, Auger M, et al. Morphologic Accuracy in Differentiating Primary Lung Adenocarcinoma From Squamous Cell Carcinoma in Cytology Specimens. *Arch Pathol Lab Med* 2016;140:1116-1120. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27552093>.

125. Yatabe Y, Dacic S, Borczuk AC, et al. Best Practices Recommendations for Diagnostic Immunohistochemistry in Lung Cancer. *J Thorac Oncol* 2019;14:377-407. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30572031>.

126. Travis WD, Brambilla E, Noguchi M, et al. Diagnosis of lung adenocarcinoma in resected specimens: implications of the 2011 International Association for the Study of Lung Cancer/American Thoracic Society/European Respiratory Society classification. *Arch Pathol Lab Med* 2013;137:685-705. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22913371>.

127. Cameron SE, Andrade RS, Pambuccian SE. Endobronchial ultrasound-guided transbronchial needle aspiration cytology: a state of the art review. *Cytopathology* 2010;21:6-26. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20015257>.

128. Moreira AL, Thornton RH. Personalized medicine for non-small-cell lung cancer: implications of recent advances in tissue acquisition for molecular and histologic testing. *Clin Lung Cancer* 2012;13:334-339. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22424871>.

129. Wang H, Wei N, Tang Y, et al. The utility of rapid on-site evaluation during bronchoscopic biopsy: A 2-year respiratory endoscopy central experience. *Biomed Res Int* 2019;2019:5049248. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31886221>.

130. Jain D, Allen TC, Aisner DL, et al. Rapid on-site evaluation of endobronchial ultrasound-guided transbronchial needle aspirations for the diagnosis of lung cancer: A perspective from members of the Pulmonary Pathology Society. *Arch Pathol Lab Med* 2018;142:253-262. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28639854>.

131. Yuan M, Wang Y, Yin W, et al. Efficacy of rapid on-site cytological evaluation (ROSE) by a pulmonologist in determining specimen adequacy and diagnostic accuracy in interventional diagnosis of lung lesions. *J Int Med Res* 2021;49:300060520982687. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/33527860>.

132. Diacon AH, Schuurmans MM, Theron J, et al. Utility of rapid on-site evaluation of transbronchial needle aspirates. *Respiration* 2005;72:182-188. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/15824529>.

133. Wahidi MM, Herth F, Yasufuku K, et al. Technical Aspects of Endobronchial Ultrasound-Guided Transbronchial Needle Aspiration: CHEST Guideline and Expert Panel Report. *Chest* 2016;149:816-835. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26402427>.

134. Thompson GR, 3rd. Pulmonary coccidioidomycosis. *Semin Respir Crit Care Med* 2011;32:754-763. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22167403>.

135. Iannuzzi MC, Fontana JR. Sarcoidosis: clinical presentation, immunopathogenesis, and therapeutics. *JAMA* 2011;305:391-399. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21266686>.

136. Centers for Disease C, Prevention. CDC Grand Rounds: the TB/HIV syndemic. *MMWR Morb Mortal Wkly Rep* 2012;61:484-489. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22763886>.

137. Travis WD, Brambilla E, Muller-Hermelink HK, Harris CC. Pathology and genetics of tumours of the lung, pleura, thymus and heart, World Health Organization Classification of Tumours. Lyon, France: IARC Press; 2004.

138. Travis WD, Brambilla E, Riely GJ. New pathologic classification of lung cancer: relevance for clinical practice and clinical trials. *J Clin Oncol* 2013;31:992-1001. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23401443>.



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

139. Lam VK, Tran HT, Banks KC, et al. Targeted Tissue and Cell-Free Tumor DNA Sequencing of Advanced Lung Squamous-Cell Carcinoma Reveals Clinically Significant Prevalence of Actionable Alterations. Clin Lung Cancer 2019;20:30-36 e33. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30279110>.

140. Sands JM, Nguyen T, Shivdasani P, et al. Next-generation sequencing informs diagnosis and identifies unexpected therapeutic targets in lung squamous cell carcinomas. Lung Cancer 2020;140:35-41. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31855703>.

141. Reck M, Rodriguez-Abreu D, Robinson AG, et al. Pembrolizumab versus Chemotherapy for PD-L1-Positive Non-Small-Cell Lung Cancer. N Engl J Med 2016;375:1823-1833. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27718847>.

142. Travis WD, Rekhtman N. Pathological diagnosis and classification of lung cancer in small biopsies and cytology: strategic management of tissue for molecular testing. Semin Respir Crit Care Med 2011;32:22-31. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21500121>.

143. Rekhtman N, Ang DC, Sima CS, et al. Immunohistochemical algorithm for differentiation of lung adenocarcinoma and squamous cell carcinoma based on large series of whole-tissue sections with validation in small specimens. Mod Pathol 2011;24:1348-1359. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21623384>.

144. Mukhopadhyay S, Katzenstein AL. Subclassification of non-small cell lung carcinomas lacking morphologic differentiation on biopsy specimens: Utility of an immunohistochemical panel containing TTF-1, napsin A, p63, and CK5/6. Am J Surg Pathol 2011;35:15-25. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21164283>.

145. Terry J, Leung S, Laskin J, et al. Optimal immunohistochemical markers for distinguishing lung adenocarcinomas from squamous cell carcinomas in small tumor samples. Am J Surg Pathol 2010;34:1805-1811. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21107086>.

146. Ordóñez NG. Thyroid transcription factor-1 is a marker of lung and thyroid carcinomas. Adv Anat Pathol 2000;7:123-127. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/10721419>.

147. Rivera MP, Mehta AC, American College of Chest P. Initial diagnosis of lung cancer: ACCP evidence-based clinical practice guidelines (2nd edition). Chest 2007;132:131S-148S. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17873165>.

148. Tan D, Zander DS. Immunohistochemistry for assessment of pulmonary and pleural neoplasms: a review and update. Int J Clin Exp Pathol 2008;1:19-31. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/18784820>.

149. Zhang H, Liu J, Cagle PT, et al. Distinction of pulmonary small cell carcinoma from poorly differentiated squamous cell carcinoma: an immunohistochemical approach. Mod Pathol 2005;18:111-118. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/15309021>.

150. Staaf J, Tran L, Soderlund L, et al. Diagnostic Value of Insulinoma-Associated Protein 1 (INSM1) and Comparison With Established Neuroendocrine Markers in Pulmonary Cancers. Arch Pathol Lab Med 2020;144:1075-1085. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31913660>.

151. Husain AN, Colby TV, Ordóñez NG, et al. Guidelines for pathologic diagnosis of malignant mesothelioma 2017 update of the Consensus Statement From the International Mesothelioma Interest Group. Arch Pathol Lab Med 2018;142:89-108. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28686500>.

152. Husain AN, Colby T, Ordóñez N, et al. Guidelines for pathologic diagnosis of malignant mesothelioma: 2012 update of the consensus statement from the International Mesothelioma Interest Group. Arch Pathol Lab Med 2013;137:647-667. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22929121>.

153. Husain AN, Colby TV, Ordóñez NG, et al. Guidelines for pathologic diagnosis of malignant mesothelioma: a consensus statement from the



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

International Mesothelioma Interest Group. Arch Pathol Lab Med 2009;133:1317-1331. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/19653732>.

154. Ordonez NG. D2-40 and podoplanin are highly specific and sensitive immunohistochemical markers of epithelioid malignant mesothelioma. Hum Pathol 2005;36:372-380. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/15891998>.

155. Amin MB, Greene FL, Edge SB, et al. AJCC staging manual, 8th ed: Springer International Publishing; 2017:1-1024.

156. Detterbeck FC, Boffa DJ, Kim AW, Tanoue LT. The Eighth Edition Lung Cancer Stage Classification. Chest 2017;151:193-203. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/27780786>.

157. Goldstraw P, Chansky K, Crowley J, et al. The IASLC Lung Cancer Staging Project: Proposals for Revision of the TNM Stage Groupings in the Forthcoming (Eighth) Edition of the TNM Classification for Lung Cancer. J Thorac Oncol 2016;11:39-51. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/26762738>.

158. Goldstraw P, Crowley J, Chansky K, et al. The IASLC Lung Cancer Staging Project: proposals for the revision of the TNM stage groupings in the forthcoming (seventh) edition of the TNM Classification of malignant tumours. J Thorac Oncol 2007;2:706-714. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/17762336>.

159. Detterbeck FC, Boffa DJ, Tanoue LT. The new lung cancer staging system. Chest 2009;136:260-271. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/19584208>.

160. Rami-Porta R, Bolejack V, Goldstraw P. The new tumor, node, and metastasis staging system. Semin Respir Crit Care Med 2011;32:44-51. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/21500123>.

161. Rami-Porta R, Crowley JJ, Goldstraw P. The revised TNM staging system for lung cancer. Ann Thorac Cardiovasc Surg 2009;15:4-9. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/19262443>.

162. Rami-Porta R, Asamura H, Travis WD, Rusch VW. Lung cancer - major changes in the American Joint Committee on Cancer eighth edition cancer staging manual. CA Cancer J Clin 2017;67:138-155. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28140453>.

163. Postmus PE, Kerr KM, Oudkerk M, et al. Early and locally advanced non-small-cell lung cancer (NSCLC): ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up. Ann Oncol 2017;28:iv1-iv21. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/28881918>.

164. Edge SB, Byrd DR, Compton CC, et al. AJCC Cancer Staging Manual, 7th ed. New York: Springer; 2010.

165. Ou SH, Zell JA, Ziogas A, Anton-Culver H. Prognostic factors for survival of stage I nonsmall cell lung cancer patients : a population-based analysis of 19,702 stage I patients in the California Cancer Registry from 1989 to 2003. Cancer 2007;110:1532-1541. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/17702091>.

166. Raz DJ, Zell JA, Ou SH, et al. Natural history of stage I non-small cell lung cancer: implications for early detection. Chest 2007;132:193-199. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/17505036>.

167. Ou SH, Kwak EL, Siwak-Tapp C, et al. Activity of crizotinib (PF02341066), a dual mesenchymal-epithelial transition (MET) and anaplastic lymphoma kinase (ALK) inhibitor, in a non-small cell lung cancer patient with de novo MET amplification. J Thorac Oncol 2011;6:942-946. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/21623265>.

168. Camidge DR, Ou S-HI, Shapiro G, et al. Efficacy and safety of crizotinib in patients with advanced c-MET-amplified non-small cell lung cancer (NSCLC) [abstract]. J Clin Oncol 2014;32(Suppl 5):Abstract 8001. Available at: https://ascopubs.org/doi/10.1200/jco.2014.32.15_suppl.8001.

169. Le X, Paz-Ares LG, Van Meerbeeck J, et al. Tepotinib in patients (pts) with advanced non-small cell lung cancer (NSCLC) with MET amplification (METamp). Journal of Clinical Oncology 2021;39:9021-9021.



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

Available at:

https://ascopubs.org/doi/abs/10.1200/JCO.2021.39.15_suppl.9021.

170. Li BT, Shen R, Buonocore D, et al. Ado-trastuzumab emtansine for patients with HER2-mutant lung cancers: results from a phase II basket trial. *J Clin Oncol* 2018;36:2532-2537. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/29989854>.

171. Li BT, Smit EF, Goto Y, et al. Trastuzumab Deruxtecan in HER2-Mutant Non-Small-Cell Lung Cancer. *N Engl J Med* 2022;386:241-251. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/34534430>.

172. Wolf J, Seto T, Han JY, et al. Capmatinib in MET exon 14-mutated or MET-amplified non-small-cell lung cancer. *N Engl J Med* 2020;383:944-957. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/32877583>.

173. Ettinger DS, Wood DE, Aisner DL, et al. NCCN Guidelines Insights: Non-Small Cell Lung Cancer, Version 2.2021. *J Natl Compr Canc Netw* 2021;19:254-266. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/33668021>.

174. Ettinger DS, Wood DE, Aisner DL, et al. NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) for Non-Small Cell Lung Cancer (Version 1.2021). © 2020 National Comprehensive Cancer Network, Inc. To view the most recent and complete version of the NCCN Guidelines®, go online to NCCN.org. 2021. Available at: www.NCCN.org.

175. Kerr KM, Bubendorf L, Edelman MJ, et al. Second ESMO consensus conference on lung cancer: pathology and molecular biomarkers for non-small-cell lung cancer. *Ann Oncol* 2014;25:1681-1690. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/24718890>.

176. Chakravarty D, Gao J, Phillips SM, et al. OncoKB: A Precision Oncology Knowledge Base. *JCO Precis Oncol* 2017;2017. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/28890946>.

177. Miller VA, Riely GJ, Zakowski MF, et al. Molecular characteristics of bronchioloalveolar carcinoma and adenocarcinoma, bronchioloalveolar carcinoma subtype, predict response to erlotinib. *J Clin Oncol* 2008;26:1472-1478. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/18349398>.

178. Sequist LV, Martins RG, Spigel D, et al. First-line gefitinib in patients with advanced non-small-cell lung cancer harboring somatic EGFR mutations. *J Clin Oncol* 2008;26:2442-2449. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/18458038>.

179. Tsao MS, Sakurada A, Cutz JC, et al. Erlotinib in lung cancer - molecular and clinical predictors of outcome. *N Engl J Med* 2005;353:133-144. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/16014883>.

180. Park K, Haura EB, Leighl NB, et al. Amivantamab in EGFR Exon 20 Insertion-Mutated Non-Small-Cell Lung Cancer Progressing on Platinum Chemotherapy: Initial Results From the CHRYSALIS Phase I Study. *J Clin Oncol* 2021;39:3391-3402. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/34339292>.

181. Zhou C, Ramalingam SS, Kim TM, et al. Treatment Outcomes and Safety of Mobocertinib in Platinum-Pretreated Patients With EGFR Exon 20 Insertion-Positive Metastatic Non-Small Cell Lung Cancer: A Phase 1/2 Open-label Nonrandomized Clinical Trial. *JAMA Oncol* 2021;7:e214761. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/34647988>.

182. Ou SH, Ahn JS, De Petris L, et al. Alectinib in Crizotinib-Refractory ALK-Rearranged Non-Small-Cell Lung Cancer: A Phase II Global Study. *J Clin Oncol* 2016;34:661-668. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/26598747>.

183. Douillard JY, Ostoros G, Cobo M, et al. First-line gefitinib in Caucasian EGFR mutation-positive NSCLC patients: a phase-IV, open-label, single-arm study. *Br J Cancer* 2014;110:55-62. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/24263064>.



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

184. Besse B, Adjei A, Baas P, et al. 2nd ESMO Consensus Conference on Lung Cancer: non-small-cell lung cancer first-line/second and further lines of treatment in advanced disease. *Ann Oncol* 2014;25:1475-1484. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24669016>.

185. Lindeman NI, Cagle PT, Beasley MB, et al. Molecular testing guideline for selection of lung cancer patients for EGFR and ALK tyrosine kinase inhibitors: guideline from the College of American Pathologists, International Association for the Study of Lung Cancer, and Association for Molecular Pathology. *J Thorac Oncol* 2013;8:823-859. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23552377>.

186. Mazieres J, Drilon A, Lusque A, et al. Immune checkpoint inhibitors for patients with advanced lung cancer and oncogenic driver alterations: results from the IMMUNOTARGET registry. *Ann Oncol* 2019;30:1321-1328. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31125062>.

187. Sholl LM, Aisner DL, Varella-Garcia M, et al. Multi-institutional Oncogenic Driver Mutation Analysis in Lung Adenocarcinoma: The Lung Cancer Mutation Consortium Experience. *J Thorac Oncol* 2015;10:768-777. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25738220>.

188. Kim AS, Bartley AN, Bridge JA, et al. Comparison of Laboratory-Developed Tests and FDA-Approved Assays for BRAF, EGFR, and KRAS Testing. *JAMA Oncol* 2018;4:838-841. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29242895>.

189. Vansteenkiste JF, Van De Kerkhove C, Wauters E, Van Mol P. Capmatinib for the treatment of non-small cell lung cancer. *Expert Rev Anticancer Ther* 2019;19:659-671. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31368815>.

190. Planchard D, Besse B, Groen HJM, et al. Dabrafenib plus trametinib in patients with previously treated BRAF(V600E)-mutant metastatic non-small cell lung cancer: an open-label, multicentre phase 2 trial. *Lancet Oncol* 2016;17:984-993. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27283860>.

191. Gautschi O, Milia J, Cabarrou B, et al. Targeted Therapy for Patients with BRAF-Mutant Lung Cancer: Results from the European EURAF Cohort. *J Thorac Oncol* 2015;10:1451-1457. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26200454>.

192. Mazieres J, Zalcman G, Crino L, et al. Crizotinib therapy for advanced lung adenocarcinoma and a ROS1 rearrangement: results from the EUROS1 cohort. *J Clin Oncol* 2015;33:992-999. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25667280>.

193. Ali G, Proietti A, Pelliccioni S, et al. ALK rearrangement in a large series of consecutive non-small cell lung cancers: comparison between a new immunohistochemical approach and fluorescence in situ hybridization for the screening of patients eligible for crizotinib treatment. *Arch Pathol Lab Med* 2014;138:1449-1458. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24885803>.

194. Slebos RJ, Kibbelaar RE, Dalesio O, et al. K-ras oncogene activation as a prognostic marker in adenocarcinoma of the lung. *N Engl J Med* 1990;323:561-565. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/2199829>.

195. Lindeman NI, Cagle PT, Aisner DL, et al. Updated Molecular Testing Guideline for the Selection of Lung Cancer Patients for Treatment With Targeted Tyrosine Kinase Inhibitors: Guideline From the College of American Pathologists, the International Association for the Study of Lung Cancer, and the Association for Molecular Pathology. *Arch Pathol Lab Med* 2018;142:321-346. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29355391>.

196. Mitchell RL, Kosche C, Burgess K, et al. Misdiagnosis of Li-Fraumeni Syndrome in a Patient With Clonal Hematopoiesis and a Somatic TP53 Mutation. *J Natl Compr Canc Netw* 2018;16:461-466. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29752319>.

197. Jaiswal S, Fontanillas P, Flannick J, et al. Age-related clonal hematopoiesis associated with adverse outcomes. *N Engl J Med* 2014;371:2488-2498. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25426837>.



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

198. Coombs CC, Zehir A, Devlin SM, et al. Therapy-Related Clonal Hematopoiesis in Patients with Non-hematologic Cancers Is Common and Associated with Adverse Clinical Outcomes. *Cell Stem Cell* 2017;21:374-382 e374. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28803919>.

199. Rolfo C, Mack P, Scagliotti GV, et al. Liquid biopsy for advanced NSCLC: A Consensus Statement From the International Association for the Study of Lung Cancer. *J Thorac Oncol* 2021;16:1647-1662. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/34246791>.

200. Rolfo C, Mack PC, Scagliotti GV, et al. Liquid Biopsy for Advanced Non-Small Cell Lung Cancer (NSCLC): A Statement Paper from the IASLC. *J Thorac Oncol* 2018;13:1248-1268. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29885479>.

201. Palmero R, Taus A, Viteri S, et al. P2.03-02 cell-free DNA (cfDNA) testing in lung adenocarcinoma (LUAC) patients: Spanish Lung Liquid Versus Invasive Biopsy Program (SLLIP) [abstract]. *J Thorac Oncol* 2018;13:Abstract S716-717. Available at: [https://www.ito.org/article/S1556-0864\(18\)32147-6/pdf](https://www.ito.org/article/S1556-0864(18)32147-6/pdf).

202. Leighl NB, Page RD, Raymond VM, et al. Clinical Utility of Comprehensive Cell-free DNA Analysis to Identify Genomic Biomarkers in Patients with Newly Diagnosed Metastatic Non-small Cell Lung Cancer. *Clin Cancer Res* 2019;25:4691-4700. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30988079>.

203. McCoach CE, Blakely CM, Banks KC, et al. Clinical Utility of Cell-Free DNA for the Detection of ALK Fusions and Genomic Mechanisms of ALK Inhibitor Resistance in Non-Small Cell Lung Cancer. *Clin Cancer Res* 2018;24:2758-2770. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29599410>.

204. Aggarwal C, Thompson JC, Black TA, et al. Clinical implications of plasma-based genotyping with the delivery of personalized therapy in metastatic non-small cell lung cancer. *JAMA Oncol* 2019;5:173-180. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30325992>.

205. Hanna NH, Robinson AG, Temin S, et al. Therapy for Stage IV Non-Small-Cell Lung Cancer With Driver Alterations: ASCO and OH (CCO) Joint Guideline Update. *J Clin Oncol* 2021;39:1040-1091. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/33591844>.

206. Roberts PJ, Stinchcombe TE. KRAS mutation: should we test for it, and does it matter? *J Clin Oncol* 2013;31:1112-1121. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23401440>.

207. Jennings LJ, Arcila ME, Corless C, et al. Guidelines for Validation of Next-Generation Sequencing-Based Oncology Panels: A Joint Consensus Recommendation of the Association for Molecular Pathology and College of American Pathologists. *J Mol Diagn* 2017;19:341-365. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28341590>.

208. Aziz N, Zhao Q, Bry L, et al. College of American Pathologists' laboratory standards for next-generation sequencing clinical tests. *Arch Pathol Lab Med* 2015;139:481-493. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25152313>.

209. Luthra R, Chen H, Roy-Chowdhuri S, Singh RR. Next-Generation Sequencing in Clinical Molecular Diagnostics of Cancer: Advantages and Challenges. *Cancers (Basel)* 2015;7:2023-2036. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26473927>.

210. Drilon A, Wang L, Arcila ME, et al. Broad, Hybrid Capture-Based Next-Generation Sequencing Identifies Actionable Genomic Alterations in Lung Adenocarcinomas Otherwise Negative for Such Alterations by Other Genomic Testing Approaches. *Clin Cancer Res* 2015;21:3631-3639. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25567908>.

211. Robson ME, Bradbury AR, Arun B, et al. American Society of Clinical Oncology Policy Statement Update: Genetic and Genomic Testing for Cancer Susceptibility. *J Clin Oncol* 2015;33:3660-3667. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26324357>.

212. Yu PP, Vose JM, Hayes DF. Genetic Cancer Susceptibility Testing: Increased Technology, Increased Complexity. *J Clin Oncol*



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

2015;33:3533-3534. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/26324366>.

213. Cardarella S, Ortiz TM, Joshi VA, et al. The introduction of systematic genomic testing for patients with non-small-cell lung cancer. *J Thorac Oncol* 2012;7:1767-1774. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/23154547>.

214. Li T, Kung HJ, Mack PC, Gandara DR. Genotyping and genomic profiling of non-small-cell lung cancer: implications for current and future therapies. *J Clin Oncol* 2013;31:1039-1049. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/23401433>.

215. Planchard D. Identification of driver mutations in lung cancer: first step in personalized cancer. *Target Oncol* 2013;8:3-14. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/23371030>.

216. Plon SE, Eccles DM, Easton D, et al. Sequence variant classification and reporting: recommendations for improving the interpretation of cancer susceptibility genetic test results. *Hum Mutat* 2008;29:1282-1291. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/18951446>.

217. Richards S, Aziz N, Bale S, et al. Standards and guidelines for the interpretation of sequence variants: a joint consensus recommendation of the American College of Medical Genetics and Genomics and the Association for Molecular Pathology. *Genet Med* 2015;17:405-424. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25741868>.

218. Shaw AT, Yeap BY, Mino-Kenudson M, et al. Clinical features and outcome of patients with non-small-cell lung cancer who harbor EML4-ALK. *J Clin Oncol* 2009;27:4247-4253. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/19667264>.

219. Shaw AT, Bauer TM, de Marinis F, et al. First-Line Lorlatinib or Crizotinib in Advanced ALK-Positive Lung Cancer. *N Engl J Med* 2020;383:2018-2029. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/33207094>.

220. Peters S, Camidge DR, Shaw AT, et al. Alectinib versus Crizotinib in Untreated ALK-Positive Non-Small-Cell Lung Cancer. *N Engl J Med* 2017;377:829-838. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/28586279>.

221. Solomon BJ, Mok T, Kim DW, et al. First-line crizotinib versus chemotherapy in ALK-positive lung cancer. *N Engl J Med* 2014;371:2167-2177. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/25470694>.

222. Soria JC, Tan DSW, Chiari R, et al. First-line ceritinib versus platinum-based chemotherapy in advanced ALK-rearranged non-small-cell lung cancer (ASCEND-4): a randomised, open-label, phase 3 study. *Lancet* 2017;389:917-929. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/28126333>.

223. Camidge DR, Kim HR, Ahn MJ, et al. Brigatinib versus Crizotinib in ALK-Positive Non-Small-Cell Lung Cancer. *N Engl J Med* 2018;379:2027-2039. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/30280657>.

224. Rogers TM, Russell PA, Wright G, et al. Comparison of methods in the detection of ALK and ROS1 rearrangements in lung cancer. *J Thorac Oncol* 2015;10:611-618. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/25789833>.

225. von Laffert M, Warth A, Penzel R, et al. Multicenter immunohistochemical ALK-testing of non-small-cell lung cancer shows high concordance after harmonization of techniques and interpretation criteria. *J Thorac Oncol* 2014;9:1685-1692. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/25436802>.

226. Wynes MW, Sholl LM, Dietel M, et al. An international interpretation study using the ALK IHC antibody D5F3 and a sensitive detection kit demonstrates high concordance between ALK IHC and ALK FISH and between evaluators. *J Thorac Oncol* 2014;9:631-638. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/24722153>.



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

227. Zhou J, Zhao J, Sun K, et al. Accurate and economical detection of ALK positive lung adenocarcinoma with semiquantitative immunohistochemical screening. PLoS One 2014;9:e92828. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24667320>.

228. Thunnissen E, Bubendorf L, Dietel M, et al. EML4-ALK testing in non-small cell carcinomas of the lung: a review with recommendations. Virchows Arch 2012;461:245-257. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22825000>.

229. Kim H, Yoo SB, Choe JY, et al. Detection of ALK gene rearrangement in non-small cell lung cancer: a comparison of fluorescence in situ hybridization and chromogenic in situ hybridization with correlation of ALK protein expression. J Thorac Oncol 2011;6:1359-1366. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21587085>.

230. Rodig SJ, Mino-Kenudson M, Dacic S, et al. Unique clinicopathologic features characterize ALK-rearranged lung adenocarcinoma in the western population. Clin Cancer Res 2009;15:5216-5223. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19671850>.

231. Mino-Kenudson M, Chirieac LR, Law K, et al. A novel, highly sensitive antibody allows for the routine detection of ALK-rearranged lung adenocarcinomas by standard immunohistochemistry. Clin Cancer Res 2010;16:1561-1571. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20179225>.

232. Ali SM, Hensing T, Schrock AB, et al. Comprehensive genomic profiling identifies a subset of crizotinib-responsive ALK-rearranged non-small cell lung cancer not detected by fluorescence in situ hybridization. Oncologist 2016;21:762-770. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27245569>.

233. Wallander ML, Geiersbach KB, Tripp SR, Layfield LJ. Comparison of reverse transcription-polymerase chain reaction, immunohistochemistry, and fluorescence in situ hybridization methodologies for detection of echinoderm microtubule-associated proteinlike 4-anaplastic lymphoma kinase fusion-positive non-small cell lung carcinoma: implications for

optimal clinical testing. Arch Pathol Lab Med 2012;136:796-803. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22742552>.

234. Weickhardt AJ, Aisner DL, Franklin WA, et al. Diagnostic assays for identification of anaplastic lymphoma kinase-positive non-small cell lung cancer. Cancer 2013;119:1467-1477. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23280244>.

235. Gainor JF, Varghese AM, Ou SH, et al. ALK rearrangements are mutually exclusive with mutations in EGFR or KRAS: an analysis of 1,683 patients with non-small cell lung cancer. Clin Cancer Res 2013;19:4273-4281. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23729361>.

236. Takahashi T, Sonobe M, Kobayashi M, et al. Clinicopathologic features of non-small-cell lung cancer with EML4-ALK fusion gene. Ann Surg Oncol 2010;17:889-897. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20183914>.

237. Magnuson WJ, Lester-Coll NH, Wu AJ, et al. Management of Brain Metastases in Tyrosine Kinase Inhibitor-Naive Epidermal Growth Factor Receptor-Mutant Non-Small-Cell Lung Cancer: A Retrospective Multi-Institutional Analysis. J Clin Oncol 2017;35:1070-1077. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28113019>.

238. Brown PD, Jaeckle K, Ballman KV, et al. Effect of Radiosurgery Alone vs Radiosurgery With Whole Brain Radiation Therapy on Cognitive Function in Patients With 1 to 3 Brain Metastases: A Randomized Clinical Trial. JAMA 2016;316:401-409. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27458945>.

239. Yamamoto M, Serizawa T, Shuto T, et al. Stereotactic radiosurgery for patients with multiple brain metastases (JLKG0901): a multi-institutional prospective observational study. Lancet Oncol 2014;15:387-395. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24621620>.

240. Reck M, Popat S, Reinmuth N, et al. Metastatic non-small-cell lung cancer (NSCLC): ESMO Clinical Practice Guidelines for diagnosis,



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

treatment and follow-up. *Ann Oncol* 2014;25 Suppl 3:iii27-39. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25115305>.

241. Tsao MN, Lloyd N, Wong RK, et al. Whole brain radiotherapy for the treatment of newly diagnosed multiple brain metastases. *Cochrane Database Syst Rev* 2012;4:CD003869. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22513917>.

242. Chang WS, Kim HY, Chang JW, et al. Analysis of radiosurgical results in patients with brain metastases according to the number of brain lesions: is stereotactic radiosurgery effective for multiple brain metastases? *J Neurosurg* 2010;113 Suppl:73-78. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21121789>.

243. Davies H, Bignell GR, Cox C, et al. Mutations of the BRAF gene in human cancer. *Nature* 2002;417:949-954. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/12068308>.

244. Paik PK, Arcila ME, Fara M, et al. Clinical characteristics of patients with lung adenocarcinomas harboring BRAF mutations. *J Clin Oncol* 2011;29:2046-2051. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21483012>.

245. Mazieres J, Cropet C, Montane L, et al. Vemurafenib in non-small-cell lung cancer patients with BRAF(V600) and BRAF(nonV600) mutations. *Ann Oncol* 2020;31:289-294. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31959346>.

246. Langer CJ. Epidermal growth factor receptor inhibition in mutation-positive non-small-cell lung cancer: is afatinib better or simply newer? *J Clin Oncol* 2013;31:3303-3306. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23980079>.

247. Hirsch FR, Bunn PA, Jr. EGFR testing in lung cancer is ready for prime time. *Lancet Oncol* 2009;10:432-433. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19410185>.

248. Riely GJ, Politi KA, Miller VA, Pao W. Update on epidermal growth factor receptor mutations in non-small cell lung cancer. *Clin Cancer Res*

2006;12:7232-7241. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17189394>.

249. O'Kane GM, Bradbury PA, Feld R, et al. Uncommon EGFR mutations in advanced non-small cell lung cancer. *Lung Cancer* 2017;109:137-144. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28577943>.

250. Paik PK, Varghese AM, Sima CS, et al. Response to erlotinib in patients with EGFR mutant advanced non-small cell lung cancers with a squamous or squamous-like component. *Mol Cancer Ther* 2012;11:2535-2540. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22896669>.

251. Sequist LV, Yang JC, Yamamoto N, et al. Phase III study of afatinib or cisplatin plus pemetrexed in patients with metastatic lung adenocarcinoma with EGFR mutations. *J Clin Oncol* 2013;31:3327-3334. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23816960>.

252. Rosell R, Carcereny E, Gervais R, et al. Erlotinib versus standard chemotherapy as first-line treatment for European patients with advanced EGFR mutation-positive non-small-cell lung cancer (EORTAC): a multicentre, open-label, randomised phase 3 trial. *Lancet Oncol* 2012;13:239-246. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22285168>.

253. Mitsudomi T, Morita S, Yatabe Y, et al. Gefitinib versus cisplatin plus docetaxel in patients with non-small-cell lung cancer harbouring mutations of the epidermal growth factor receptor (WJTOG3405): an open label, randomised phase 3 trial. *Lancet Oncol* 2010;11:121-128. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20022809>.

254. Maemondo M, Inoue A, Kobayashi K, et al. Gefitinib or chemotherapy for non-small-cell lung cancer with mutated EGFR. *N Engl J Med* 2010;362:2380-2388. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20573926>.

255. Zhou C, Wu YL, Chen G, et al. Erlotinib versus chemotherapy as first-line treatment for patients with advanced EGFR mutation-positive non-small-cell lung cancer (OPTIMAL, CTONG-0802): a multicentre,



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

open-label, randomised, phase 3 study. *Lancet Oncol* 2011;12:735-742. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21783417>.

256. Zhou C, Wu YL, Chen G, et al. Updated efficacy and quality-of-life (QoL) analyses in OPTIMAL, a phase III, randomized, open-label study of first-line erlotinib versus gemcitabine/carboplatin in patients with EGFR-activating mutation-positive (EGFR Act Mut+) advanced non-small cell lung cancer (NSCLC) [abstract]. *J Clin Oncol* 2011;29(Suppl 15):Abstract 7520. Available at: http://meeting.ascopubs.org/cgi/content/abstract/29/15_suppl/7520.

257. Yang JC, Wu YL, Schuler M, et al. Afatinib versus cisplatin-based chemotherapy for EGFR mutation-positive lung adenocarcinoma (LUX-Lung 3 and LUX-Lung 6): analysis of overall survival data from two randomised, phase 3 trials. *Lancet Oncol* 2015;16:141-151. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25589191>.

258. Mok TS, Wu YL, Thongprasert S, et al. Gefitinib or carboplatin-paclitaxel in pulmonary adenocarcinoma. *N Engl J Med* 2009;361:947-957. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19692680>.

259. Vasconcelos P, Gergis C, Viray H, et al. EGFR-A763_Y764insFQEA Is a Unique Exon 20 Insertion Mutation That Displays Sensitivity to Approved and In-Development Lung Cancer EGFR Tyrosine Kinase Inhibitors. *JTO Clin Res Rep* 2020;1. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/34104899>.

260. Yasuda H, Park E, Yun CH, et al. Structural, biochemical, and clinical characterization of epidermal growth factor receptor (EGFR) exon 20 insertion mutations in lung cancer. *Sci Transl Med* 2013;5:216ra177. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24353160>.

261. Arcila ME, Nafa K, Chaff JE, et al. EGFR exon 20 insertion mutations in lung adenocarcinomas: prevalence, molecular heterogeneity, and clinicopathologic characteristics. *Mol Cancer Ther* 2013;12:220-229. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23371856>.

262. Oxnard GR, Lo PC, Nishino M, et al. Natural history and molecular characteristics of lung cancers harboring EGFR exon 20 insertions. *J Thorac Oncol* 2013;8:179-184. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23328547>.

263. Lund-Iversen M, Kleinberg L, Fjellbirkeland L, et al. Clinicopathological characteristics of 11 NSCLC patients with EGFR-exon 20 mutations. *J Thorac Oncol* 2012;7:1471-1473. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22895145>.

264. Yasuda H, Kobayashi S, Costa DB. EGFR exon 20 insertion mutations in non-small-cell lung cancer: preclinical data and clinical implications. *Lancet Oncol* 2012;13:e23-31. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21764376>.

265. Nishino M, Cardarella S, Jackman DM, et al. RECIST 1.1 in NSCLC patients with EGFR mutations treated with EGFR tyrosine kinase inhibitors: comparison with RECIST 1.0. *AJR Am J Roentgenol* 2013;201:W64-71. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23789698>.

266. Mok TS, Wu YL, Ahn MJ, et al. Osimertinib or Platinum-Pemetrexed in EGFR T790M-Positive Lung Cancer. *N Engl J Med* 2017;376:629-640. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27959700>.

267. Yu HA, Arcila ME, Rekhtman N, et al. Analysis of tumor specimens at the time of acquired resistance to EGFR-TKI therapy in 155 patients with EGFR-mutant lung cancers. *Clin Cancer Res* 2013;19:2240-2247. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23470965>.

268. Riely GJ, Yu HA. EGFR: The Paradigm of an Oncogene-Driven Lung Cancer. *Clin Cancer Res* 2015;21:2221-2226. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25979928>.

269. Finlay MR, Anderton M, Ashton S, et al. Discovery of a potent and selective EGFR inhibitor (AZD9291) of both sensitizing and T790M resistance mutations that spares the wild type form of the receptor. *J Med Chem* 2014;57:8249-8267. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25271963>.



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

270. Gainor JF, Shaw AT. Emerging paradigms in the development of resistance to tyrosine kinase inhibitors in lung cancer. *J Clin Oncol* 2013;31:3987-3996. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/24101047>.

271. Pao W, Miller VA, Politi KA, et al. Acquired resistance of lung adenocarcinomas to gefitinib or erlotinib is associated with a second mutation in the EGFR kinase domain. *PLoS Med* 2005;2:e73. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/15737014>.

272. Kosaka T, Yatabe Y, Endoh H, et al. Analysis of epidermal growth factor receptor gene mutation in patients with non-small cell lung cancer and acquired resistance to gefitinib. *Clin Cancer Res* 2006;12:5764-5769. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/17020982>.

273. Onitsuka T, Uramoto H, Nose N, et al. Acquired resistance to gefitinib: the contribution of mechanisms other than the T790M, MET, and HGF status. *Lung Cancer* 2010;68:198-203. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/19589612>.

274. Rosell R, Molina MA, Costa C, et al. Pretreatment EGFR T790M mutation and BRCA1 mRNA expression in erlotinib-treated advanced non-small-cell lung cancer patients with EGFR mutations. *Clin Cancer Res* 2011;17:1160-1168. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/21233402>.

275. Berry DK, Wang X, Michalski ST, et al. Clinical Cohort Analysis of Germline EGFR T790M Demonstrates Penetrance Across Ethnicities and Races, Sexes, and Ages. *JCO Precis Oncol* 2020;4. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/34164592>.

276. Oxnard GR, Miller VA, Robson ME, et al. Screening for germline EGFR T790M mutations through lung cancer genotyping. *J Thorac Oncol* 2012;7:1049-1052. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/22588155>.

277. Gazdar A, Robinson L, Oliver D, et al. Hereditary lung cancer syndrome targets never smokers with germline EGFR gene T790M

mutations. *J Thorac Oncol* 2014;9:456-463. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/24736066>.

278. Schoenfeld AJ, Chan JM, Kubota D, et al. Tumor Analyses Reveal Squamous Transformation and Off-Target Alterations As Early Resistance Mechanisms to First-line Osimertinib in EGFR-Mutant Lung Cancer. *Clin Cancer Res* 2020;26:2654-2663. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/31911548>.

279. Marcoux N, Gettinger SN, O'Kane G, et al. EGFR-Mutant Adenocarcinomas That Transform to Small-Cell Lung Cancer and Other Neuroendocrine Carcinomas: Clinical Outcomes. *J Clin Oncol* 2019;37:278-285. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/30550363>.

280. Sequist LV, Waltman BA, Dias-Santagata D, et al. Genotypic and histological evolution of lung cancers acquiring resistance to EGFR inhibitors. *Sci Transl Med* 2011;3:75ra26. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/21430269>.

281. Oxnard GR. Strategies for overcoming acquired resistance to epidermal growth factor receptor: targeted therapies in lung cancer. *Arch Pathol Lab Med* 2012;136:1205-1209. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/23020725>.

282. Suda K, Mizuuchi H, Maehara Y, Mitsudomi T. Acquired resistance mechanisms to tyrosine kinase inhibitors in lung cancer with activating epidermal growth factor receptor mutation--diversity, ductility, and destiny. *Cancer Metastasis Rev* 2012;31:807-814. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/22736441>.

283. Yu HA, Suzawa K, Jordan E, et al. Concurrent Alterations in EGFR-Mutant Lung Cancers Associated with Resistance to EGFR Kinase Inhibitors and Characterization of MTOR as a Mediator of Resistance. *Clin Cancer Res* 2018;24:3108-3118. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/29530932>.

284. Sholl LM, Cagle PT, Lindeman NI, et al. Template for reporting results of biomarker testing of specimens from patients with non-small cell



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

carcinoma of the lung; Version: LungBiomarkers 1.3.0.0: College of American Pathologists; 2016. Available at: <http://www.cap.org>.

285. Han SW, Kim TY, Jeon YK, et al. Optimization of patient selection for gefitinib in non-small cell lung cancer by combined analysis of epidermal growth factor receptor mutation, K-ras mutation, and Akt phosphorylation. Clin Cancer Res 2006;12:2538-2544. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/16638863>.

286. Dacic S. EGFR assays in lung cancer. Adv Anat Pathol 2008;15:241-247. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/18580100>.

287. Sholl LM, Xiao Y, Joshi V, et al. EGFR mutation is a better predictor of response to tyrosine kinase inhibitors in non-small cell lung carcinoma than FISH, CISH, and immunohistochemistry. Am J Clin Pathol 2010;133:922-934. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20472851>.

288. Eberhard DA, Giaccone G, Johnson BE, Non-Small-Cell Lung Cancer Working G. Biomarkers of response to epidermal growth factor receptor inhibitors in Non-Small-Cell Lung Cancer Working Group: standardization for use in the clinical trial setting. J Clin Oncol 2008;26:983-994. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/18281673>.

289. Pao W, Ladanyi M. Epidermal growth factor receptor mutation testing in lung cancer: searching for the ideal method. Clin Cancer Res 2007;13:4954-4955. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17785543>.

290. Shepherd FA, Tsao MS. Epidermal growth factor receptor biomarkers in non-small-cell lung cancer: a riddle, wrapped in a mystery, inside an enigma. J Clin Oncol 2010;28:903-905. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20100955>.

291. Dias-Santagata D, Akhavanfard S, David SS, et al. Rapid targeted mutational analysis of human tumours: a clinical platform to guide

personalized cancer medicine. EMBO Mol Med 2010;2:146-158. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20432502>.

292. Janne PA, Yang JC, Kim DW, et al. AZD9291 in EGFR inhibitor-resistant non-small-cell lung cancer. N Engl J Med 2015;372:1689-1699. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25923549>.

293. Cho JH, Lim SH, An HJ, et al. Osimertinib for Patients With Non-Small-Cell Lung Cancer Harboring Uncommon EGFR Mutations: A Multicenter, Open-Label, Phase II Trial (KCSG-LU15-09). J Clin Oncol 2020;38:488-495. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31825714>.

294. Tu HY, Ke EE, Yang JJ, et al. A comprehensive review of uncommon EGFR mutations in patients with non-small cell lung cancer. Lung Cancer 2017;114:96-102. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29173773>.

295. Otsuka T, Mori M, Yano Y, et al. Effectiveness of Tyrosine Kinase Inhibitors in Japanese Patients with Non-small Cell Lung Cancer Harboring Minor Epidermal Growth Factor Receptor Mutations: Results from a Multicenter Retrospective Study (HANSHIN Oncology Group 0212). Anticancer Res 2015;35:3885-3891. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26124334>.

296. Riess JW, Gandara DR, Frampton GM, et al. Diverse EGFR Exon 20 Insertions and Co-Occurring Molecular Alterations Identified by Comprehensive Genomic Profiling of NSCLC. J Thorac Oncol 2018;13:1560-1568. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29981927>.

297. Riely GJ, Neal JW, Camidge DR, et al. Activity and Safety of Mobocertinib (TAK-788) in Previously Treated Non-Small Cell Lung Cancer with EGFR Exon 20 Insertion Mutations from a Phase I/II Trial. Cancer Discov 2021;11:1688-1699. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/33632775>.



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

298. Piotrowska Z, Wang Y, Sequist LV, Ramalingam SS. ECOG-ACRIN 5162: A phase II study of osimertinib 160 mg in NSCLC with EGFR exon 20 insertions. *Journal of Clinical Oncology* 2020;38:9513-9513. Available at: https://ascopubs.org/doi/abs/10.1200/JCO.2020.38.15_suppl.9513.

299. Chelabi S, Mignard X, Leroy K, et al. EGFR Exon 20 Insertion in Metastatic Non-Small-Cell Lung Cancer: Survival and Clinical Efficacy of EGFR Tyrosine-Kinase Inhibitor and Chemotherapy. *Cancers (Basel)* 2021;13. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/34680280>.

300. Byeon S, Kim Y, Lim SW, et al. Clinical Outcomes of EGFR Exon 20 Insertion Mutations in Advanced Non-small Cell Lung Cancer in Korea. *Cancer Res Treat* 2019;51:623-631. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30049203>.

301. Naidoo J, Sima CS, Rodriguez K, et al. Epidermal growth factor receptor exon 20 insertions in advanced lung adenocarcinomas: Clinical outcomes and response to erlotinib. *Cancer* 2015;121:3212-3220. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26096453>.

302. Dersarkissian M, Bhak R, Lin H, et al. P2.01-103 Real-World Treatment Patterns and Survival in Non-Small Cell Lung Cancer Patients with EGFR Exon 20 Insertion Mutations. *Journal of Thoracic Oncology* 2019;14:S681. Available at: <https://doi.org/10.1016/j.jtho.2019.08.1446>.

303. Morita C, Yoshida T, Shirasawa M, et al. Clinical characteristics of advanced non-small cell lung cancer patients with EGFR exon 20 insertions. *Sci Rep* 2021;11:18762. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/34548567>.

304. Ou S-HI, Lin HM, Hong J-L, et al. Real-world response and outcomes in NSCLC patients with EGFR exon 20 insertion mutations. *Journal of Clinical Oncology* 2021;39:9098-9098. Available at: https://ascopubs.org/doi/abs/10.1200/JCO.2021.39.15_suppl.9098.

305. Udagawa H, Matsumoto S, Ohe Y, et al. OA07.03 Clinical Outcome of Non-Small Cell Lung Cancer with EGFR/HER2 Exon 20 Insertions Identified in the LC-SCRUM-Japan. *Journal of Thoracic Oncology* 2019;14:S224. Available at: <https://doi.org/10.1016/j.jtho.2019.08.443>.

306. Pillai RN, Behera M, Berry LD, et al. HER2 mutations in lung adenocarcinomas: A report from the Lung Cancer Mutation Consortium. *Cancer* 2017;123:4099-4105. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28743157>.

307. Offin M, Feldman D, Ni A, et al. Frequency and outcomes of brain metastases in patients with HER2-mutant lung cancers. *Cancer* 2019;125:4380-4387. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31469421>.

308. Saalfeld FC, Wenzel C, Christopoulos P, et al. Efficacy of immune checkpoint inhibitors alone or in combination with chemotherapy in NSCLC harboring ERBB2 mutations. *J Thorac Oncol* 2021;16:1952-1958. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/34245914>.

309. Sequist LV, Heist RS, Shaw AT, et al. Implementing multiplexed genotyping of non-small-cell lung cancers into routine clinical practice. *Ann Oncol* 2011;22:2616-2624. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22071650>.

310. Tsao MS, Aviel-Ronen S, Ding K, et al. Prognostic and predictive importance of p53 and RAS for adjuvant chemotherapy in non small-cell lung cancer. *J Clin Oncol* 2007;25:5240-5247. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/18024870>.

311. Slebos RJ, Hruban RH, Dalesio O, et al. Relationship between K-ras oncogene activation and smoking in adenocarcinoma of the human lung. *J Natl Cancer Inst* 1991;83:1024-1027. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/2072410>.

312. Mitsudomi T, Steinberg SM, Oie HK, et al. ras gene mutations in non-small cell lung cancers are associated with shortened survival irrespective of treatment intent. *Cancer Res* 1991;51:4999-5002. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/1654209>.

313. Febbo PG, Ladanyi M, Aldape KD, et al. NCCN Task Force report: Evaluating the clinical utility of tumor markers in oncology. *J Natl Compr Canc Netw* 2011;9 Suppl 5:S1-32; quiz S33. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22138009>.



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

314. Lee SE, Lee B, Hong M, et al. Comprehensive analysis of RET and ROS1 rearrangement in lung adenocarcinoma. *Mod Pathol* 2015;28:468-479. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25234288>.

315. Kim JO, Lee J, Shin JY, et al. KIF5B-RET Fusion gene may coincide oncogenic mutations of EGFR or KRAS gene in lung adenocarcinomas. *Diagn Pathol* 2015;10:143. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26268359>.

316. Skoulidis F, Goldberg ME, Greenawalt DM, et al. STK11/LKB1 mutations and PD-1 inhibitor resistance in KRAS-mutant lung adenocarcinoma. *Cancer Discov* 2018;8:822-835. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29773717>.

317. Janne PA, Riely GJ, Gadgeel SM, et al. Adagrasib in non-small-cell lung cancer harboring a KRAS(G12C) mutation. *N Engl J Med* 2022;387:120-131. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/35658005>.

318. de Langen AJ, Johnson ML, Mazieres J, et al. Sotorasib versus docetaxel for previously treated non-small-cell lung cancer with KRAS(G12C) mutation: a randomised, open-label, phase 3 trial. *Lancet* 2023. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/36764316>.

319. Skoulidis F, Li BT, Dy GK, et al. Sotorasib for lung cancers with KRAS p.G12C mutation. *N Engl J Med* 2021;384:2371-2381. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/34096690>.

320. Awad MM, Oxnard GR, Jackman DM, et al. MET exon 14 mutations in non-small-cell lung cancer are associated with advanced age and stage-dependent MET genomic amplification and c-Met overexpression. *J Clin Oncol* 2016;34:721-730. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26729443>.

321. Frampton GM, Ali SM, Rosenzweig M, et al. Activation of MET via diverse exon 14 splicing alterations occurs in multiple tumor types and confers clinical sensitivity to MET inhibitors. *Cancer Discov*

2015;5:850-859. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25971938>.

322. Paik PK, Drilon A, Fan PD, et al. Response to MET inhibitors in patients with stage IV lung adenocarcinomas harboring MET mutations causing exon 14 skipping. *Cancer Discov* 2015;5:842-849. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25971939>.

323. Vuong HG, Ho ATN, Altibi AMA, et al. Clinicopathological implications of MET exon 14 mutations in non-small cell lung cancer - A systematic review and meta-analysis. *Lung Cancer* 2018;123:76-82. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30089599>.

324. Sabari JK, Leonardi GC, Shu CA, et al. PD-L1 expression, tumor mutational burden, and response to immunotherapy in patients with MET exon 14 altered lung cancers. *Ann Oncol* 2018;29:2085-2091. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30165371>.

325. Kron A, Scheffler M, Heydt C, et al. Genetic Heterogeneity of MET-Abserrant NSCLC and Its Impact on the Outcome of Immunotherapy. *J Thorac Oncol* 2021;16:572-582. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/33309988>.

326. Wolf J, Setons T, Han J-Y, et al. Capmatinib (INC280) in METΔex14-mutated advanced non-small cell lung cancer (NSCLC): Efficacy data from the phase II GEOMETRY mono-1 study [abstract]. *J Clin Oncol* 2019;37:Abstract 9004. Available at: https://ascopubs.org/doi/abs/10.1200/JCO.2019.37.15_suppl.9004.

327. Drilon A, Clark JW, Weiss J, et al. Antitumor activity of crizotinib in lung cancers harboring a MET exon 14 alteration. *Nat Med* 2020;26:47-51. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31932802>.

328. Gatalica Z, Xiu J, Swensen J, Vranic S. Molecular characterization of cancers with NTRK gene fusions. *Mod Pathol* 2019;32:147-153. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30171197>.

329. Farago AF, Taylor MS, Doebele RC, et al. Clinicopathologic Features of Non-Small-Cell Lung Cancer Harboring an NTRK Gene Fusion. *JCO*



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

Precis Oncol 2018;2018. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/30215037>.

330. Drilon A, Laetsch TW, Kummar S, et al. Efficacy of Larotrectinib in TRK Fusion-Positive Cancers in Adults and Children. *N Engl J Med* 2018;378:731-739. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/29466156>.

331. Benayed R, Offin M, Mullaney K, et al. High Yield of RNA Sequencing for Targetable Kinase Fusions in Lung Adenocarcinomas with No Mitogenic Driver Alteration Detected by DNA Sequencing and Low Tumor Mutation Burden. *Clin Cancer Res* 2019;25:4712-4722. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31028088>.

332. Drilon A, Siena S, Ou SI, et al. Safety and Antitumor Activity of the Multitargeted Pan-TRK, ROS1, and ALK Inhibitor Entrectinib: Combined Results from Two Phase I Trials (ALKA-372-001 and STARTRK-1). *Cancer Discov* 2017;7:400-409. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28183697>.

333. Doebele RC, Drilon A, Paz-Ares L, et al. Entrectinib in patients with advanced or metastatic NTRK fusion-positive solid tumours: integrated analysis of three phase 1-2 trials. *Lancet Oncol* 2020;21:271-282. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31838007>.

334. Gautschi O, Milia J, Filleron T, et al. Targeting RET in Patients With RET-Rearranged Lung Cancers: Results From the Global, Multicenter RET Registry. *J Clin Oncol* 2017;35:1403-1410. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28447912>.

335. Ferrara R, Auger N, Auclin E, Besse B. Clinical and Translational Implications of RET Rearrangements in Non-Small Cell Lung Cancer. *J Thorac Oncol* 2018;13:27-45. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29128428>.

336. Michels S, Scheel AH, Scheffler M, et al. Clinicopathological Characteristics of RET Rearranged Lung Cancer in European Patients. *J Thorac Oncol* 2016;11:122-127. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26762747>.

337. Takeuchi K, Soda M, Togashi Y, et al. RET, ROS1 and ALK fusions in lung cancer. *Nat Med* 2012;18:378-381. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22327623>.

338. Tsuta K, Kohno T, Yoshida A, et al. RET-rearranged non-small-cell lung carcinoma: a clinicopathological and molecular analysis. *Br J Cancer* 2014;110:1571-1578. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24504365>.

339. Gainor JF, Curigliano G, Kim DW, et al. Pralsetinib for RET fusion-positive non-small-cell lung cancer (ARROW): a multi-cohort, open-label, phase 1/2 study. *Lancet Oncol* 2021;22:959-969. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/34118197>.

340. Drilon A, Oxnard GR, Tan DSW, et al. Efficacy of Selpercatinib in RET Fusion-Positive Non-Small-Cell Lung Cancer. *N Engl J Med* 2020;383:813-824. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32846060>.

341. Drilon A, Rekhtman N, Arcila M, et al. Cabozantinib in patients with advanced RET-rearranged non-small-cell lung cancer: an open-label, single-centre, phase 2, single-arm trial. *Lancet Oncol* 2016;17:1653-1660. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27825636>.

342. Drilon A, Wang L, Hasanovic A, et al. Response to Cabozantinib in patients with RET fusion-positive lung adenocarcinomas. *Cancer Discov* 2013;3:630-635. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23533264>.

343. Yoh K, Seto T, Satouchi M, et al. Final survival results for the LURET phase II study of vandetanib in previously treated patients with RET-rearranged advanced non-small cell lung cancer. *Lung Cancer* 2021;155:40-45. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/33725547>.

344. Lee SH, Lee JK, Ahn MJ, et al. Vandetanib in pretreated patients with advanced non-small cell lung cancer-harboring RET rearrangement: a phase II clinical trial. *Ann Oncol* 2017;28:292-297. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27803005>.



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

345. Robinson DR, Wu YM, Lin SF. The protein tyrosine kinase family of the human genome. *Oncogene* 2000;19:5548-5557. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/11114734>.

346. Shaw AT, Ou SH, Bang YJ, et al. Crizotinib in ROS1-rearranged non-small-cell lung cancer. *N Engl J Med* 2014;371:1963-1971. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25264305>.

347. Dugay F, Llamas-Gutierrez F, Gournay M, et al. Clinicopathological characteristics of ROS1- and RET-rearranged NSCLC in caucasian patients: Data from a cohort of 713 non-squamous NSCLC lacking KRAS/EGFR/HER2/BRAF/PIK3CA/ALK alterations. *Oncotarget* 2017;8:53336-53351. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28881815>.

348. Bergeth K, Shaw AT, Ou SH, et al. ROS1 rearrangements define a unique molecular class of lung cancers. *J Clin Oncol* 2012;30:863-870. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22215748>.

349. Kim HR, Lim SM, Kim HJ, et al. The frequency and impact of ROS1 rearrangement on clinical outcomes in never smokers with lung adenocarcinoma. *Ann Oncol* 2013;24:2364-2370. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23788756>.

350. Drilon A, Barlesi F, De Braud F, et al. Entrectinib in locally advanced or metastatic ROS1 fusion positive non-small cell lung cancer (NSCLC): Integrated analysis of STARTRK-2, STARTRK-1, and ALKA-372-001 [abstract]. *Cancer Research* 2019;79:Abstract CT192. Available at: https://cancerres.aacrjournals.org/content/79/13_Supplement/CT192.

351. Kazandjian D, Blumenthal GM, Luo L, et al. Benefit-Risk Summary of Crizotinib for the Treatment of Patients With ROS1 Alteration-Positive, Metastatic Non-Small Cell Lung Cancer. *Oncologist* 2016;21:974-980. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27328934>.

352. Cohen D, Hondelink LM, Solleveld-Westerink N, et al. Optimizing Mutation and Fusion Detection in NSCLC by Sequential DNA and RNA Sequencing. *J Thorac Oncol* 2020;15:1000-1014. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32014610>.

353. Bubendorf L, Buttner R, Al-Dayel F, et al. Testing for ROS1 in non-small cell lung cancer: a review with recommendations. *Virchows Arch* 2016;469:489-503. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27535289>.

354. Davies KD, Le AT, Theodoro MF, et al. Identifying and targeting ROS1 gene fusions in non-small cell lung cancer. *Clin Cancer Res* 2012;18:4570-4579. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22919003>.

355. Clave S, Gimeno J, Munoz-Marmol AM, et al. ROS1 copy number alterations are frequent in non-small cell lung cancer. *Oncotarget* 2016;7:8019-8028. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26783962>.

356. Zacharias M, Absenger G, Kashofer K, et al. Reflex testing in non-small cell lung carcinoma using DNA- and RNA-based next-generation sequencing-a single-center experience. *Transl Lung Cancer Res* 2021;10:4221-4234. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/35004252>.

357. Doebele RC, Paz-Ares L, Farago AF, et al. Entrectinib in NTRK fusion-positive non-small cell lung cancer (NSCLC): integrated analysis of patients enrolled in three trials (STARTRK-2, STARTRK-1 and ALKA-372-001) [abstract]. *AACR Annual Meeting*. Atlanta, GA:Abstract CT131. Available at: https://cancerres.aacrjournals.org/content/79/13_Supplement/CT131.

358. Shaw AT, Solomon BJ, Chiari R, et al. Lorlatinib in advanced ROS1-positive non-small-cell lung cancer: a multicentre, open-label, single-arm, phase 1-2 trial. *Lancet Oncol* 2019;20:1691-1701. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31669155>.

359. Drilon A, Siena S, Dziadziuszko R, et al. Entrectinib in ROS1 fusion-positive non-small-cell lung cancer: integrated analysis of three phase 1-2 trials. *Lancet Oncol* 2020;21:261-270. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31838015>.



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

360. Borghaei H, Paz-Ares L, Horn L, et al. Nivolumab versus docetaxel in advanced nonsquamous non-small-cell lung cancer. *N Engl J Med* 2015;373:1627-1639. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26412456>.

361. Ribas A. Releasing the Brakes on Cancer Immunotherapy. *N Engl J Med* 2015;373:1490-1492. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26348216>.

362. Brahmer JR, Hammers H, Lipson EJ. Nivolumab: targeting PD-1 to bolster antitumor immunity. *Future Oncol* 2015;11:1307-1326. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25798726>.

363. Sezer A, Kilickap S, Gumus M, et al. Cemiplimab monotherapy for first-line treatment of advanced non-small-cell lung cancer with PD-L1 of at least 50%: a multicentre, open-label, global, phase 3, randomised, controlled trial. *Lancet* 2021;397:592-604. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/33581821>.

364. Brahmer J, Reckamp KL, Baas P, et al. Nivolumab versus Docetaxel in Advanced Squamous-Cell Non-Small-Cell Lung Cancer. *N Engl J Med* 2015;373:123-135. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26028407>.

365. Antonia SJ, Villegas A, Daniel D, et al. Durvalumab after chemoradiotherapy in stage III non-small-cell lung cancer. *N Engl J Med* 2017;377:1919-1929. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28885881>.

366. Fehrenbacher L, Spira A, Ballinger M, et al. Atezolizumab versus docetaxel for patients with previously treated non-small-cell lung cancer (POPLAR): a multicentre, open-label, phase 2 randomised controlled trial. *Lancet* 2016;387:1837-1846. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26970723>.

367. Gadgeel SM, Stevenson J, Langer C, et al. Pembrolizumab (pembro) plus chemotherapy as front-line therapy for advanced NSCLC: KEYNOTE-021 cohorts A-C [abstract]. *J Clin Oncol* 2016;34:Abstract 9016. Available at: <http://meetinglibrary.asco.org/content/167088-176>.

368. Kerr KM, Nicolson MC. Non-Small Cell Lung Cancer, PD-L1, and the Pathologist. *Arch Pathol Lab Med* 2016;140:249-254. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26927720>.

369. Kerr KM, Hirsch FR. Programmed Death Ligand-1 Immunohistochemistry: Friend or Foe? *Arch Pathol Lab Med* 2016;140:326-331. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26756647>.

370. Rimm DL, Han G, Taube JM, et al. A Prospective, Multi-institutional, Pathologist-Based Assessment of 4 Immunohistochemistry Assays for PD-L1 Expression in Non-Small Cell Lung Cancer. *JAMA Oncol* 2017;3:1051-1058. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28278348>.

371. Buttner R, Gosney JR, Skov BG, et al. Programmed Death-Ligand 1 Immunohistochemistry Testing: A Review of Analytical Assays and Clinical Implementation in Non-Small-Cell Lung Cancer. *J Clin Oncol* 2017;35:3867-3876. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29053400>.

372. Kerr KM, Tsao MS, Nicholson AG, et al. Programmed Death-Ligand 1 Immunohistochemistry in Lung Cancer: In what state is this art? *J Thorac Oncol* 2015;10:985-989. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26134220>.

373. Soria JC, Ohe Y, Vansteenkiste J, et al. Osimertinib in Untreated EGFR-Mutated Advanced Non-Small-Cell Lung Cancer. *N Engl J Med* 2018;378:113-125. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29151359>.

374. Lisberg A, Cummings A, Goldman JW, et al. A Phase II Study of Pembrolizumab in EGFR-Mutant, PD-L1+, Tyrosine Kinase Inhibitor Naive Patients With Advanced NSCLC. *J Thorac Oncol* 2018;13:1138-1145. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29874546>.

375. Camidge DR, Dziadziuszko R, Peters S, et al. Updated Efficacy and Safety Data and Impact of the EML4-ALK Fusion Variant on the Efficacy of Alectinib in Untreated ALK-Positive Advanced Non-Small Cell Lung



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

Cancer in the Global Phase III ALEX Study. J Thorac Oncol 2019;14:1233-1243. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30902613>.

376. Gainor JF, Dardaei L, Yoda S, et al. Molecular Mechanisms of Resistance to First- and Second-Generation ALK Inhibitors in ALK-Rearranged Lung Cancer. Cancer Discov 2016;6:1118-1133. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27432227>.

377. Sholl LM, Hirsch FR, Hwang D, et al. The promises and challenges of tumor mutation burden as an immunotherapy biomarker: A perspective from the International Association for the Study of Lung Cancer Pathology Committee. J Thorac Oncol 2020;15:1409-1424. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32522712>.

378. Sesma A, Pardo J, Cruellas M, et al. From Tumor Mutational Burden to Blood T Cell Receptor: Looking for the Best Predictive Biomarker in Lung Cancer Treated with Immunotherapy. Cancers (Basel) 2020;12. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/33066479>.

379. Offin M, Rizvi H, Tenet M, et al. Tumor Mutation Burden and Efficacy of EGFR-Tyrosine Kinase Inhibitors in Patients with EGFR-Mutant Lung Cancers. Clin Cancer Res 2019;25:1063-1069. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30045933>.

380. Hellmann MD, Ciuleanu TE, Pluzanski A, et al. Nivolumab plus ipilimumab in lung cancer with a high tumor mutational burden. N Engl J Med 2018;378:2093-2104. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29658845>.

381. Hellmann MD, Paz-Ares L, Bernabe Caro R, et al. Nivolumab plus ipilimumab in advanced non-small-cell lung cancer. N Engl J Med 2019;381:2020-2031. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31562796>.

382. Rizvi H, Sanchez-Vega F, La K, et al. Molecular Determinants of Response to Anti-Programmed Cell Death (PD)-1 and Anti-Programmed Death-Ligand 1 (PD-L1) Blockade in Patients With Non-Small-Cell Lung Cancer Profiled With Targeted Next-Generation Sequencing. J Clin Oncol

2018;36:633-641. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29337640>.

383. Carbone DP, Reck M, Paz-Ares L, et al. First-Line Nivolumab in Stage IV or Recurrent Non-Small-Cell Lung Cancer. N Engl J Med 2017;376:2415-2426. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28636851>.

384. Marabelle A, Fakih M, Lopez J, et al. Association of tumour mutational burden with outcomes in patients with advanced solid tumours treated with pembrolizumab: prospective biomarker analysis of the multicohort, open-label, phase 2 KEYNOTE-158 study. Lancet Oncol 2020;21:1353-1365. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32919526>.

385. Chow WB, Rosenthal RA, Merkow RP, et al. Optimal preoperative assessment of the geriatric surgical patient: a best practices guideline from the American College of Surgeons National Surgical Quality Improvement Program and the American Geriatrics Society. J Am Coll Surg 2012;215:453-466. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22917646>.

386. Howington JA, Blum MG, Chang AC, et al. Treatment of stage I and II non-small cell lung cancer: Diagnosis and management of lung cancer, 3rd ed: American College of Chest Physicians evidence-based clinical practice guidelines. Chest 2013;143:e278S-e313S. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23649443>.

387. Altorki N, Wang X, Kozono D, et al. Lobar or sublobar resection for peripheral stage IA non-small-cell lung cancer. N Engl J Med 2023;388:489-498. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/36780674>.

388. Pennathur A, Brunelli A, Criner GJ, et al. Definition and assessment of high risk in patients considered for lobectomy for stage I non-small cell lung cancer: The American Association for Thoracic Surgery expert panel consensus document. J Thorac Cardiovasc Surg 2021;162:1605-1618 e1606. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/34716030>.



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

389. Handforth C, Clegg A, Young C, et al. The prevalence and outcomes of frailty in older cancer patients: a systematic review. *Ann Oncol* 2015;26:1091-1101. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/25403592>.

390. Caillet P, Laurent M, Bastuji-Garin S, et al. Optimal management of elderly cancer patients: usefulness of the Comprehensive Geriatric Assessment. *Clin Interv Aging* 2014;9:1645-1660. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/25302022>.

391. Pallis AG, Gridelli C, Wedding U, et al. Management of elderly patients with NSCLC; updated expert's opinion paper: EORTC Elderly Task Force, Lung Cancer Group and International Society for Geriatric Oncology. *Ann Oncol* 2014;25:1270-1283. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/24638905>.

392. Brunelli A, Kim AW, Berger KI, Addrizzo-Harris DJ. Physiologic evaluation of the patient with lung cancer being considered for resectional surgery: Diagnosis and management of lung cancer, 3rd ed: American College of Chest Physicians evidence-based clinical practice guidelines. *Chest* 2013;143:e166S-e190S. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/23649437>.

393. Turner G, Clegg A, British Geriatrics S, et al. Best practice guidelines for the management of frailty: a British Geriatrics Society, Age UK and Royal College of General Practitioners report. *Age Ageing* 2014;43:744-747. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/25336440>.

394. Vairaktarakis C, Tsiamis V, Soursoy G, et al. A computer-aided diagnosis system for geriatrics assessment and frailty evaluation. *Adv Exp Med Biol* 2015;820:69-77. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/25417017>.

395. Wildiers H, Heeren P, Puts M, et al. International Society of Geriatric Oncology consensus on geriatric assessment in older patients with cancer. *J Clin Oncol* 2014;32:2595-2603. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/25071125>.

396. Kozower BD, Lerner JM, Detterbeck FC, Jones DR. Special treatment issues in non-small cell lung cancer: Diagnosis and management of lung cancer, 3rd ed: American College of Chest Physicians evidence-based clinical practice guidelines. *Chest* 2013;143:e369S-e399S. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/23649447>.

397. Boffa DJ, Allen MS, Grab JD, et al. Data from The Society of Thoracic Surgeons General Thoracic Surgery database: the surgical management of primary lung tumors. *J Thorac Cardiovasc Surg* 2008;135:247-254. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/18242243>.

398. Scott WJ, Howington J, Feigenberg S, et al. Treatment of non-small cell lung cancer stage I and stage II: ACCP evidence-based clinical practice guidelines (2nd edition). *Chest* 2007;132:234S-242S. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/17873171>.

399. Saji H, Okada M, Tsuboi M, et al. Segmentectomy versus lobectomy in small-sized peripheral non-small-cell lung cancer (JCOG0802/WJOG4607L): a multicentre, open-label, phase 3, randomised, controlled, non-inferiority trial. *Lancet* 2022;399:1607-1617. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/35461558>.

400. Villamizar N, Swanson SJ. Lobectomy vs. segmentectomy for NSCLC (T<2 cm). *Ann Cardiothorac Surg* 2014;3:160-166. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/24790839>.

401. Landreneau RJ, Normolle DP, Christie NA, et al. Recurrence and survival outcomes after anatomic segmentectomy versus lobectomy for clinical stage I non-small-cell lung cancer: a propensity-matched analysis. *J Clin Oncol* 2014;32:2449-2455. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/24982447>.

402. Altorki NK, Yip R, Hanaoka T, et al. Sublobar resection is equivalent to lobectomy for clinical stage 1A lung cancer in solid nodules. *J Thorac Cardiovasc Surg* 2014;147:754-762; Discussion 762-754. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/24280722>.



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

403. Sienel W, Dango S, Kirschbaum A, et al. Sublobar resections in stage IA non-small cell lung cancer: segmentectomies result in significantly better cancer-related survival than wedge resections. *Eur J Cardiothorac Surg* 2008;33:728-734. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/18261918>.

404. Sienel W, Stremmel C, Kirschbaum A, et al. Frequency of local recurrence following segmentectomy of stage IA non-small cell lung cancer is influenced by segment localisation and width of resection margins--implications for patient selection for segmentectomy. *Eur J Cardiothorac Surg* 2007;31:522-527; discussion 527-528. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17229574>.

405. Narsule CK, Ebright MI, Fernando HC. Sublobar versus lobar resection: current status. *Cancer J* 2011;17:23-27. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21263263>.

406. Videtic GMM, Donington J, Giuliani M, et al. Stereotactic body radiation therapy for early-stage non-small cell lung cancer: Executive Summary of an ASTRO Evidence-Based Guideline. *Pract Radiat Oncol* 2017;7:295-301. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28596092>.

407. Timmerman R, Paulus R, Galvin J, et al. Stereotactic body radiation therapy for inoperable early stage lung cancer. *JAMA* 2010;303:1070-1076. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20233825>.

408. Woody NM, Stephans KL, Marwaha G, et al. Stereotactic Body Radiation Therapy for Non-Small Cell Lung Cancer Tumors Greater Than 5 cm: Safety and Efficacy. *Int J Radiat Oncol Biol Phys* 2015;92:325-331. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25841625>.

409. Grills IS, Mangona VS, Welsh R, et al. Outcomes after stereotactic lung radiotherapy or wedge resection for stage I non-small-cell lung cancer. *J Clin Oncol* 2010;28:928-935. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20065181>.

410. Donington J, Ferguson M, Mazzone P, et al. American College of Chest Physicians and Society of Thoracic Surgeons consensus statement for evaluation and management for high-risk patients with stage I non-small cell lung cancer. *Chest* 2012;142:1620-1635. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23208335>.

411. Darling GE, Allen MS, Decker PA, et al. Randomized trial of mediastinal lymph node sampling versus complete lymphadenectomy during pulmonary resection in the patient with N0 or N1 (less than hilar) non-small cell carcinoma: results of the American College of Surgery Oncology Group Z0030 Trial. *J Thorac Cardiovasc Surg* 2011;141:662-670. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21335122>.

412. Allen MS, Darling GE, Pechet TT, et al. Morbidity and mortality of major pulmonary resections in patients with early-stage lung cancer: initial results of the randomized, prospective ACOSOG Z0030 trial. *Ann Thorac Surg* 2006;81:1013-1019; discussion 1019-1020. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/16488712>.

413. Rusch VW, Asamura H, Watanabe H, et al. The IASLC lung cancer staging project: a proposal for a new international lymph node map in the forthcoming seventh edition of the TNM classification for lung cancer. *J Thorac Oncol* 2009;4:568-577. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19357537>.

414. Swanson SJ, Batirel HF. Video-assisted thoracic surgery (VATS) resection for lung cancer. *Surg Clin North Am* 2002;82:541-559. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/12371584>.

415. Mahtabifard A, Fuller CB, McKenna RJ, Jr. Video-assisted thoracic surgery sleeve lobectomy: a case series. *Ann Thorac Surg* 2008;85:S729-732. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/18222205>.

416. Shaw JP, Dembitzer FR, Wisnivesky JP, et al. Video-assisted thoracoscopic lobectomy: state of the art and future directions. *Ann Thorac Surg* 2008;85:S705-709. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/18222201>.



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

417. Cheng D, Downey RJ, Kernstine K, et al. Video-assisted thoracic surgery in lung cancer resection: a meta-analysis and systematic review of controlled trials. *Innovations (Phila)* 2007;2:261-292. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22437196>.
418. Alam N, Flores RM. Video-assisted thoracic surgery (VATS) lobectomy: the evidence base. *JSLs* 2007;11:368-374. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17931521>.
419. Whitson BA, Andrade RS, Boettcher A, et al. Video-assisted thoracoscopic surgery is more favorable than thoracotomy for resection of clinical stage I non-small cell lung cancer. *Ann Thorac Surg* 2007;83:1965-1970. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17532379>.
420. Whitson BA, Groth SS, Duval SJ, et al. Surgery for early-stage non-small cell lung cancer: a systematic review of the video-assisted thoracoscopic surgery versus thoracotomy approaches to lobectomy. *Ann Thorac Surg* 2008;86:2008-2016; discussion 2016-2008. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19022040>.
421. Scott WJ, Allen MS, Darling G, et al. Video-assisted thoracic surgery versus open lobectomy for lung cancer: a secondary analysis of data from the American College of Surgeons Oncology Group Z0030 randomized clinical trial. *J Thorac Cardiovasc Surg* 2010;139:976-981; discussion 981-973. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20172539>.
422. Atkins BZ, Harpole DH, Jr., Mangum JH, et al. Pulmonary segmentectomy by thoracotomy or thoracoscopy: reduced hospital length of stay with a minimally-invasive approach. *Ann Thorac Surg* 2007;84:1107-1112; discussion 1112-1103. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17888955>.
423. Swanson SJ, Herndon JE, 2nd, D'Amico TA, et al. Video-assisted thoracic surgery lobectomy: report of CALGB 39802--a prospective, multi-institution feasibility study. *J Clin Oncol* 2007;25:4993-4997. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17971599>.
424. Ohtsuka T, Nomori H, Horio H, et al. Is major pulmonary resection by video-assisted thoracic surgery an adequate procedure in clinical stage I lung cancer? *Chest* 2004;125:1742-1746. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/15136385>.
425. McKenna RJ, Jr. New approaches to the minimally invasive treatment of lung cancer. *Cancer J* 2005;11:73-76. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/15831227>.
426. Demmy TL, Nwogu C. Is video-assisted thoracic surgery lobectomy better? Quality of life considerations. *Ann Thorac Surg* 2008;85:S719-728. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/18222204>.
427. Cattaneo SM, Park BJ, Wilton AS, et al. Use of video-assisted thoracic surgery for lobectomy in the elderly results in fewer complications. *Ann Thorac Surg* 2008;85:231-235; discussion 235-236. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/18154816>.
428. Cao C, Manganas C, Ang SC, et al. Video-assisted thoracic surgery versus open thoracotomy for non-small cell lung cancer: a meta-analysis of propensity score-matched patients. *Interact Cardiovasc Thorac Surg* 2013;16:244-249. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23169877>.
429. Ilonen IK, Rasanen JV, Knuuttila A, et al. Anatomic thoracoscopic lung resection for non-small cell lung cancer in stage I is associated with less morbidity and shorter hospitalization than thoracotomy. *Acta Oncol* 2011;50:1126-1132. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21314296>.
430. Villamizar NR, Darrabie MD, Burfeind WR, et al. Thoracoscopic lobectomy is associated with lower morbidity compared with thoracotomy. *J Thorac Cardiovasc Surg* 2009;138:419-425. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19619789>.
431. Paul S, Altorki NK, Sheng S, et al. Thoracoscopic lobectomy is associated with lower morbidity than open lobectomy: a propensity-matched analysis from the STS database. *J Thorac Cardiovasc*



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

Surg 2010;139:366-378. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/20106398>.

432. Su S, Scott WJ, Allen MS, et al. Patterns of survival and recurrence after surgical treatment of early stage non-small cell lung carcinoma in the ACOSOG Z0030 (ALLIANCE) trial. J Thorac Cardiovasc Surg 2014;147:747-752; Discussion 752-743. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/24290575>.

433. Lee PC, Nasar A, Port JL, et al. Long-term survival after lobectomy for non-small cell lung cancer by video-assisted thoracic surgery versus thoracotomy. Ann Thorac Surg 2013;96:951-960; discussion 960-951. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/23866808>.

434. Thomas P, Doddoli C, Yena S, et al. VATS is an adequate oncological operation for stage I non-small cell lung cancer. Eur J Cardiothorac Surg 2002;21:1094-1099. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/12048091>.

435. Roviato G, Varoli F, Vergani C, et al. Long-term survival after videothoroscopic lobectomy for stage I lung cancer. Chest 2004;126:725-732. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/15364748>.

436. Solaini L, Prusciano F, Bagioni P, Poddie DB. Long-term results of video-assisted thoracic surgery lobectomy for stage I non-small cell lung cancer: a single-centre study of 104 cases. Interact Cardiovasc Thorac Surg 2004;3:57-62. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/17670176>.

437. Demmy TL, Plante AJ, Nwogu CE, et al. Discharge independence with minimally invasive lobectomy. Am J Surg 2004;188:698-702. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/15619486>.

438. Demmy TL. VATS lobectomy for frail or complex patients. Chest Meeting Abstracts 2003;124:234S. Available at:

<http://meeting.chestpubs.org/cgi/reprint/124/4/234S.pdf>.

439. Nicastrì DG, Wisnivesky JP, Little VR, et al. Thoracoscopic lobectomy: report on safety, discharge independence, pain, and chemotherapy tolerance. J Thorac Cardiovasc Surg 2008;135:642-647. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/18329487>.

440. Petersen RP, Pham D, Burfeind WR, et al. Thoracoscopic lobectomy facilitates the delivery of chemotherapy after resection for lung cancer. Ann Thorac Surg 2007;83:1245-1249; discussion 1250. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/17383320>.

441. Hanna JM, Berry MF, D'Amico TA. Contraindications of video-assisted thoracoscopic surgical lobectomy and determinants of conversion to open. J Thorac Dis 2013;5 Suppl 3:S182-189. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/24040521>.

442. Yan TD, Cao C, D'Amico TA, et al. Video-assisted thoracoscopic surgery lobectomy at 20 years: a consensus statement. Eur J Cardiothorac Surg 2014;45:633-639. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/24130372>.

443. Yan TD, Black D, Bannon PG, McCaughan BC. Systematic review and meta-analysis of randomized and nonrandomized trials on safety and efficacy of video-assisted thoracic surgery lobectomy for early-stage non-small-cell lung cancer. J Clin Oncol 2009;27:2553-2562. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/19289625>.

444. Cao C, Manganas C, Ang SC, Yan TD. A meta-analysis of unmatched and matched patients comparing video-assisted thoracoscopic lobectomy and conventional open lobectomy. Ann Cardiothorac Surg 2012;1:16-23. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/23977459>.

445. Nakamura H. Systematic review of published studies on safety and efficacy of thoracoscopic and robot-assisted lobectomy for lung cancer. Ann Thorac Cardiovasc Surg 2014;20:93-98. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/24583699>.

446. Swanson SJ, Miller DL, McKenna RJ, Jr., et al. Comparing robot-assisted thoracic surgical lobectomy with conventional



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

video-assisted thoracic surgical lobectomy and wedge resection: results from a multihospital database (Premier). *J Thorac Cardiovasc Surg* 2014;147:929-937. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/24210834>.

447. Martins RG, D'Amico TA, Loo BW, Jr., et al. The management of patients with stage IIIA non-small cell lung cancer with N2 mediastinal node involvement. *J Natl Compr Canc Netw* 2012;10:599-613. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22570291>.

448. Farjah F, Flum DR, Varghese TK, Jr., et al. Surgeon specialty and long-term survival after pulmonary resection for lung cancer. *Ann Thorac Surg* 2009;87:995-1004; discussion 1005-1006. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19324119>.

449. Albain KS, Swann RS, Rusch VW, et al. Radiotherapy plus chemotherapy with or without surgical resection for stage III non-small-cell lung cancer: a phase III randomised controlled trial. *Lancet* 2009;374:379-386. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19632716>.

450. van Meerbeeck JP, Kramer GW, Van Schil PE, et al. Randomized controlled trial of resection versus radiotherapy after induction chemotherapy in stage IIIA-N2 non-small-cell lung cancer. *J Natl Cancer Inst* 2007;99:442-450. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17374834>.

451. Cerfolio RJ, Bryant AS. Survival of patients with unsuspected N2 (stage IIIA) nonsmall-cell lung cancer. *Ann Thorac Surg* 2008;86:362-366; discussion 366-367. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/18640297>.

452. Le Pechoux C, Pourel N, Barlesi F, et al. Postoperative radiotherapy versus no postoperative radiotherapy in patients with completely resected non-small-cell lung cancer and proven mediastinal N2 involvement (Lung ART): an open-label, randomised, phase 3 trial. *Lancet Oncol* 2022;23:104-114. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/34919827>.

453. Sher DJ, Fidler MJ, Liptay MJ, Koshy M. Comparative effectiveness of neoadjuvant chemoradiotherapy versus chemotherapy alone followed by surgery for patients with stage IIIA non-small cell lung cancer. *Lung Cancer* 2015;88:267-274. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25862147>.

454. Shah AA, Berry MF, Tzao C, et al. Induction chemoradiation is not superior to induction chemotherapy alone in stage IIIA lung cancer. *Ann Thorac Surg* 2012;93:1807-1812. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22632486>.

455. Hui Z, Men Y, Hu C, et al. Effect of postoperative radiotherapy for patients with pIIIA-N2 non-small cell lung cancer after complete resection and adjuvant chemotherapy: the phase 3 PORT-C randomized clinical trial. *JAMA Oncol* 2021;7:1178-1185. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/34165501>.

456. Gopal RS, Dubey S, Rosenzweig KE, et al. ACR Appropriateness Criteria(R) on Induction and Adjuvant Therapy for Stage N2 Non-Small-Cell Lung Cancer: expert panel on radiation oncology-lung. *Int J Radiat Oncol Biol Phys* 2010;78:969-974. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20813465>.

457. Stefani A, Alifano M, Bobbio A, et al. Which patients should be operated on after induction chemotherapy for N2 non-small cell lung cancer? Analysis of a 7-year experience in 175 patients. *J Thorac Cardiovasc Surg* 2010;140:356-363. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20381815>.

458. Evans NR, 3rd, Li S, Wright CD, et al. The impact of induction therapy on morbidity and operative mortality after resection of primary lung cancer. *J Thorac Cardiovasc Surg* 2010;139:991-996 e991-992. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20304144>.

459. Gaissert HA, Keum DY, Wright CD, et al. POINT: Operative risk of pneumonectomy--influence of preoperative induction therapy. *J Thorac Cardiovasc Surg* 2009;138:289-294. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19619768>.



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

460. Mansour Z, Kochetkova EA, Ducrocq X, et al. Induction chemotherapy does not increase the operative risk of pneumonectomy! Eur J Cardiothorac Surg 2007;31:181-185. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17141515>.

461. Weder W, Collaud S, Eberhardt WE, et al. Pneumonectomy is a valuable treatment option after neoadjuvant therapy for stage III non-small-cell lung cancer. J Thorac Cardiovasc Surg 2010;139:1424-1430. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20416887>.

462. Kappers I, van Sandick JW, Burgers SA, et al. Surgery after induction chemotherapy in stage IIIA-N2 non-small cell lung cancer: why pneumonectomy should be avoided. Lung Cancer 2010;68:222-227. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19664843>.

463. Decaluwe H, De Leyn P, Vansteenkiste J, et al. Surgical multimodality treatment for baseline resectable stage IIIA-N2 non-small cell lung cancer. Degree of mediastinal lymph node involvement and impact on survival. Eur J Cardiothorac Surg 2009;36:433-439. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19502079>.

464. Konig D, Schar S, Vuong D, et al. Long-term outcomes of operable stage III NSCLC in the pre-immunotherapy era: results from a pooled analysis of the SAKK 16/96, SAKK 16/00, SAKK 16/01, and SAKK 16/08 trials. ESMO Open 2022;7:100455. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/35398718>.

465. Thomas M, Rube C, Hoffknecht P, et al. Effect of preoperative chemoradiation in addition to preoperative chemotherapy: a randomised trial in stage III non-small-cell lung cancer. Lancet Oncol 2008;9:636-648. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/18583190>.

466. Higgins K, Chino JP, Marks LB, et al. Preoperative chemotherapy versus preoperative chemoradiotherapy for stage III (N2) non-small-cell lung cancer. Int J Radiat Oncol Biol Phys 2009;75:1462-1467. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19467798>.

467. Lung Cancer Study G. Effects of postoperative mediastinal radiation on completely resected stage II and stage III epidermoid cancer of the lung. N Engl J Med 1986;315:1377-1381. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/2877397>.

468. Keller SM, Adak S, Wagner H, et al. A randomized trial of postoperative adjuvant therapy in patients with completely resected stage II or IIIA non-small-cell lung cancer. Eastern Cooperative Oncology Group. N Engl J Med 2000;343:1217-1222. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/11071672>.

469. Douillard JY, Rosell R, De Lena M, et al. Impact of postoperative radiation therapy on survival in patients with complete resection and stage I, II, or IIIA non-small-cell lung cancer treated with adjuvant chemotherapy: the adjuvant Navelbine International Trialist Association (ANITA) Randomized Trial. Int J Radiat Oncol Biol Phys 2008;72:695-701. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/18439766>.

470. Bradley JD, Paulus R, Graham MV, et al. Phase II trial of postoperative adjuvant paclitaxel/carboplatin and thoracic radiotherapy in resected stage II and IIIA non-small-cell lung cancer: promising long-term results of the Radiation Therapy Oncology Group--RTOG 9705. J Clin Oncol 2005;23:3480-3487. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/15908657>.

471. Feigenberg SJ, Hanlon AL, Langer C, et al. A phase II study of concurrent carboplatin and paclitaxel and thoracic radiotherapy for completely resected stage II and IIIA non-small cell lung cancer. J Thorac Oncol 2007;2:287-292. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17409799>.

472. Jaklitsch MT, Herndon JE, 2nd, DeCamp MM, Jr., et al. Nodal downstaging predicts survival following induction chemotherapy for stage IIIA (N2) non-small cell lung cancer in CALGB protocol #8935. J Surg Oncol 2006;94:599-606. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17039491>.

473. Chang JY, Mehran RJ, Feng L, et al. Stereotactic ablative radiotherapy for operable stage I non-small-cell lung cancer (revised



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

STARS): long-term results of a single-arm, prospective trial with prespecified comparison to surgery. *Lancet Oncol* 2021;22:1448-1457. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/34529930>.

474. Gjyshi O, Xu T, Elhammali A, et al. Toxicity and Survival After Intensity-Modulated Proton Therapy Versus Passive Scattering Proton Therapy for NSCLC. *J Thorac Oncol* 2021;16:269-277. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/33198942>.

475. Kamran SC, Yeap BY, Ulysse CA, et al. Assessment of a Contralateral Esophagus-Sparing Technique in Locally Advanced Lung Cancer Treated With High-Dose Chemoradiation: A Phase 1 Nonrandomized Clinical Trial. *JAMA Oncol* 2021;7:910-914. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/33830168>.

476. Kong F-MS, Hu C, Ten Haken R, et al. NRG-RT0G 1106/ACRIN 6697: A phase IIR trial of standard versus adaptive (mid-treatment PET-based) chemoradiotherapy for stage III NSCLC—Results and comparison to NRG-RT0G 0617 (non-personalized RT dose escalation). *Journal of Clinical Oncology* 2021;39:8548-8548. Available at: https://ascopubs.org/doi/abs/10.1200/JCO.2021.39.15_suppl.8548.

477. Theelen W, Chen D, Verma V, et al. Pembrolizumab with or without radiotherapy for metastatic non-small-cell lung cancer: a pooled analysis of two randomised trials. *Lancet Respir Med* 2021;9:467-475. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/33096027>.

478. Lutz ST, Jones J, Chow E. Role of radiation therapy in palliative care of the patient with cancer. *J Clin Oncol* 2014;32:2913-2919. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25113773>.

479. McAvoy S, Ciura K, Wei C, et al. Definitive reirradiation for locoregionally recurrent non-small cell lung cancer with proton beam therapy or intensity modulated radiation therapy: predictors of high-grade toxicity and survival outcomes. *Int J Radiat Oncol Biol Phys* 2014;90:819-827. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25220718>.

480. Expert Panel on Radiation Oncology-Brain M, Lo SS, Gore EM, et al. ACR Appropriateness Criteria(R) pre-irradiation evaluation and management of brain metastases. *J Palliat Med* 2014;17:880-886. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24971478>.

481. Expert Panel on Radiation Oncology-Bone M, Lo SS, Lutz ST, et al. ACR Appropriateness Criteria (R) spinal bone metastases. *J Palliat Med* 2013;16:9-19. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23167547>.

482. Expert Panel On Radiation Oncology-Bone M, Lutz ST, Lo SS, et al. ACR Appropriateness Criteria(R) non-spine bone metastases. *J Palliat Med* 2012;15:521-526. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22536988>.

483. Patel SH, Robbins JR, Gore EM, et al. ACR Appropriateness Criteria(R) follow-up and retreatment of brain metastases. *Am J Clin Oncol* 2012;35:302-306. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22609733>.

484. Chang JY, Kestin LL, Barriger RB, et al. ACR Appropriateness Criteria(R) nonsurgical treatment for locally advanced non-small-cell lung cancer: good performance status/definitive intent. *Oncology (Williston Park)* 2014;28:706-710, 712, 714 passim. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25140629>.

485. Rosenzweig KE, Chang JY, Chetty IJ, et al. ACR appropriateness criteria nonsurgical treatment for non-small-cell lung cancer: poor performance status or palliative intent. *J Am Coll Radiol* 2013;10:654-664. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23890874>.

486. Sejpal S, Komaki R, Tsao A, et al. Early findings on toxicity of proton beam therapy with concurrent chemotherapy for nonsmall cell lung cancer. *Cancer* 2011;117:3004-3013. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21264827>.

487. Chang JY, Verma V, Li M, et al. Proton Beam Radiotherapy and Concurrent Chemotherapy for Unresectable Stage III Non-Small Cell Lung



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

Cancer: Final Results of a Phase 2 Study. JAMA Oncol 2017;3:e172032. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28727865>.

488. Gregoire V, Mackie TR. State of the art on dose prescription, reporting and recording in Intensity-Modulated Radiation Therapy (ICRU report No. 83). Cancer Radiother 2011;15:555-559. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21802333>.

489. Teoh M, Clark CH, Wood K, et al. Volumetric modulated arc therapy: a review of current literature and clinical use in practice. Br J Radiol 2011;84:967-996. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22011829>.

490. Chen AB, Neville BA, Sher DJ, et al. Survival outcomes after radiation therapy for stage III non-small-cell lung cancer after adoption of computed tomography-based simulation. J Clin Oncol 2011;29:2305-2311. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21537034>.

491. Liao ZX, Komaki RR, Thames HD, Jr., et al. Influence of technologic advances on outcomes in patients with unresectable, locally advanced non-small-cell lung cancer receiving concomitant chemoradiotherapy. Int J Radiat Oncol Biol Phys 2010;76:775-781. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19515503>.

492. Terasawa T, Dvorak T, Ip S, et al. Systematic review: charged-particle radiation therapy for cancer. Ann Intern Med 2009;151:556-565. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19755348>.

493. Chun SG, Hu C, Choy H, et al. Impact of Intensity-Modulated Radiation Therapy Technique for Locally Advanced Non-Small-Cell Lung Cancer: A Secondary Analysis of the NRG Oncology RTOG 0617 Randomized Clinical Trial. J Clin Oncol 2017;35:56-62. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28034064>.

494. Ung Y, Gu C-S, Cline K, et al. An Ontario Clinical Oncology Group (OCOG) randomized trial (PET SMART) of FDG PET/CT in patients with stage 3 non-small cell lung cancer (NSCLC): impact of PET on radiation

treatment volumes [Abstract O35.01]. J Thorac Oncol 2011;6:S428. Available at: <http://journals.lww.com/jto/toc/2011/06001>.

495. MacManus M, Nestle U, Rosenzweig KE, et al. Use of PET and PET/CT for radiation therapy planning: IAEA expert report 2006-2007. Radiother Oncol 2009;91:85-94. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19100641>.

496. Everitt S, Herschtal A, Callahan J, et al. High rates of tumor growth and disease progression detected on serial pretreatment fluorodeoxyglucose-positron emission tomography/computed tomography scans in radical radiotherapy candidates with nonsmall cell lung cancer. Cancer 2010;116:5030-5037. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20623786>.

497. Mohammed N, Kestin LL, Grills IS, et al. Rapid disease progression with delay in treatment of non-small-cell lung cancer. Int J Radiat Oncol Biol Phys 2011;79:466-472. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20471184>.

498. Chang JY, Zhang X, Wang X, et al. Significant reduction of normal tissue dose by proton radiotherapy compared with three-dimensional conformal or intensity-modulated radiation therapy in Stage I or Stage III non-small-cell lung cancer. Int J Radiat Oncol Biol Phys 2006;65:1087-1096. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/16682145>.

499. Bush DA, Slater JD, Shin BB, et al. Hypofractionated proton beam radiotherapy for stage I lung cancer. Chest 2004;126:1198-1203. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/15486383>.

500. Nihei K, Ogino T, Ishikura S, Nishimura H. High-dose proton beam therapy for Stage I non-small-cell lung cancer. Int J Radiat Oncol Biol Phys 2006;65:107-111. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/16458447>.

501. Grutters JP, Kessels AG, Pijls-Johannesma M, et al. Comparison of the effectiveness of radiotherapy with photons, protons and carbon-ions for non-small cell lung cancer: a meta-analysis. Radiother Oncol



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

2010;95:32-40. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/19733410>.

502. Keall PJ, Mageras GS, Balter JM, et al. The management of respiratory motion in radiation oncology report of AAPM Task Group 76. Med Phys 2006;33:3874-3900. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/17089851>.

503. Kong FM, Pan C, Eisbruch A, Ten Haken RK. Physical models and simpler dosimetric descriptors of radiation late toxicity. Semin Radiat Oncol 2007;17:108-120. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/17395041>.

504. Timmerman RD. An overview of hypofractionation and introduction to this issue of seminars in radiation oncology. Semin Radiat Oncol 2008;18:215-222. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/18725106>.

505. Rusch VW, Giroux DJ, Kraut MJ, et al. Induction chemoradiation and surgical resection for superior sulcus non-small-cell lung carcinomas: long-term results of Southwest Oncology Group Trial 9416 (Intergroup Trial 0160). J Clin Oncol 2007;25:313-318. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/17235046>.

506. Cerfolio RJ, Bryant AS, Jones VL, Cerfolio RM. Pulmonary resection after concurrent chemotherapy and high dose (60Gy) radiation for non-small cell lung cancer is safe and may provide increased survival. Eur J Cardiothorac Surg 2009;35:718-723; discussion 723. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/19233668>.

507. Kwong KF, Edelman MJ, Suntharalingam M, et al. High-dose radiotherapy in trimodality treatment of Pancoast tumors results in high pathologic complete response rates and excellent long-term survival. J Thorac Cardiovasc Surg 2005;129:1250-1257. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/15942564>.

508. Sonett JR, Suntharalingam M, Edelman MJ, et al. Pulmonary resection after curative intent radiotherapy (>59 Gy) and concurrent chemotherapy in non-small-cell lung cancer. Ann Thorac Surg

2004;78:1200-1205; discussion 1206. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/15464470>.

509. ICRU. ICRU Report 50. Prescribing, Recording and Reporting Photon Beam Therapy. Bethesda, MD: International Commission on Radiation Units and Measurements; 1993.

510. ICRU. Prescribing, Recording and Reporting Photon Beam Therapy (Report 62) (Supplement to ICRU Report 50). Bethesda, MD: ICRU; 1999.

511. ICRU Report 83: prescribing, recording, and reporting intensity-modulated photon-beam therapy (IMRT). Bethesda, MD: International Commission on Radiation Units and Measurements; 2010. Available at: <http://bit.ly/2pBwGkl>.

512. Group IDW, Holmes T, Das R, et al. American Society of Radiation Oncology recommendations for documenting intensity-modulated radiation therapy treatments. Int J Radiat Oncol Biol Phys 2009;74:1311-1318. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19616738>.

513. Kong FM, Ritter T, Quint DJ, et al. Consideration of dose limits for organs at risk of thoracic radiotherapy: atlas for lung, proximal bronchial tree, esophagus, spinal cord, ribs, and brachial plexus. Int J Radiat Oncol Biol Phys 2011;81:1442-1457. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/20934273>.

514. Marks LB, Yorke ED, Jackson A, et al. Use of normal tissue complication probability models in the clinic. Int J Radiat Oncol Biol Phys 2010;76:S10-19. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/20171502>.

515. Marks LB, Bentzen SM, Deasy JO, et al. Radiation dose-volume effects in the lung. Int J Radiat Oncol Biol Phys 2010;76:S70-76. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20171521>.

516. Werner-Wasik M, Yorke E, Deasy J, et al. Radiation dose-volume effects in the esophagus. Int J Radiat Oncol Biol Phys 2010;76:S86-93. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20171523>.



517. Gagliardi G, Constone LS, Moiseenko V, et al. Radiation dose-volume effects in the heart. *Int J Radiat Oncol Biol Phys* 2010;76:S77-85. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20171522>.

518. Kirkpatrick JP, van der Kogel AJ, Schultheiss TE. Radiation dose-volume effects in the spinal cord. *Int J Radiat Oncol Biol Phys* 2010;76:S42-49. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20171517>.

519. Speirs CK, DeWees TA, Rehman S, et al. Heart Dose Is an Independent Dosimetric Predictor of Overall Survival in Locally Advanced Non-Small Cell Lung Cancer. *J Thorac Oncol* 2017;12:293-301. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27743888>.

520. Wang K, Eblan MJ, Deal AM, et al. Cardiac Toxicity After Radiotherapy for Stage III Non-Small-Cell Lung Cancer: Pooled Analysis of Dose-Escalation Trials Delivering 70 to 90 Gy. *J Clin Oncol* 2017;35:1387-1394. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28113017>.

521. Al-Halabi H, Paetzold P, Sharp GC, et al. A contralateral esophagus-sparing technique to limit severe esophagitis associated with concurrent high-dose radiation and chemotherapy in patients with thoracic malignancies. *Int J Radiat Oncol Biol Phys* 2015;92:803-810. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26104934>.

522. Amini A, Yang J, Williamson R, et al. Dose constraints to prevent radiation-induced brachial plexopathy in patients treated for lung cancer. *Int J Radiat Oncol Biol Phys* 2012;82:e391-398. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22284035>.

523. Graham MV, Purdy JA, Emami B, et al. Clinical dose-volume histogram analysis for pneumonitis after 3D treatment for non-small cell lung cancer (NSCLC). *Int J Radiat Oncol Biol Phys* 1999;45:323-329. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/10487552>.

524. Palma DA, Senan S, Tsujino K, et al. Predicting radiation pneumonitis after chemoradiation therapy for lung cancer: an international individual patient data meta-analysis. *Int J Radiat Oncol Biol Phys*

2013;85:444-450. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22682812>.

525. Kong FM, Hayman JA, Griffith KA, et al. Final toxicity results of a radiation-dose escalation study in patients with non-small-cell lung cancer (NSCLC): predictors for radiation pneumonitis and fibrosis. *Int J Radiat Oncol Biol Phys* 2006;65:1075-1086. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/16647222>.

526. Hernando ML, Marks LB, Bentel GC, et al. Radiation-induced pulmonary toxicity: a dose-volume histogram analysis in 201 patients with lung cancer. *Int J Radiat Oncol Biol Phys* 2001;51:650-659. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/11597805>.

527. Kim TH, Cho KH, Pyo HR, et al. Dose-volumetric parameters for predicting severe radiation pneumonitis after three-dimensional conformal radiation therapy for lung cancer. *Radiology* 2005;235:208-215. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/15703313>.

528. Wang S, Liao Z, Wei X, et al. Analysis of clinical and dosimetric factors associated with treatment-related pneumonitis (TRP) in patients with non-small-cell lung cancer (NSCLC) treated with concurrent chemotherapy and three-dimensional conformal radiotherapy (3D-CRT). *Int J Radiat Oncol Biol Phys* 2006;66:1399-1407. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/16997503>.

529. Rose J, Rodrigues G, Yaremko B, et al. Systematic review of dose-volume parameters in the prediction of esophagitis in thoracic radiotherapy. *Radiother Oncol* 2009;91:282-287. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/18950881>.

530. Hall WH, Guiou M, Lee NY, et al. Development and validation of a standardized method for contouring the brachial plexus: preliminary dosimetric analysis among patients treated with IMRT for head-and-neck cancer. *Int J Radiat Oncol Biol Phys* 2008;72:1362-1367. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/18448267>.

531. Bezjak A, Temin S, Franklin G, et al. Definitive and Adjuvant Radiotherapy in Locally Advanced Non-Small-Cell Lung Cancer: American



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

Society of Clinical Oncology Clinical Practice Guideline Endorsement of the American Society for Radiation Oncology Evidence-Based Clinical Practice Guideline. J Clin Oncol 2015;33:2100-2105. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25944914>.

532. Bradley J, Graham MV, Winter K, et al. Toxicity and outcome results of RTOG 9311: a phase I-II dose-escalation study using three-dimensional conformal radiotherapy in patients with inoperable non-small-cell lung carcinoma. Int J Radiat Oncol Biol Phys 2005;61:318-328. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/15667949>.

533. Dess RT, Sun Y, Matuszak MM, et al. Cardiac Events After Radiation Therapy: Combined Analysis of Prospective Multicenter Trials for Locally Advanced Non-Small-Cell Lung Cancer. J Clin Oncol 2017;35:1395-1402. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28301264>.

534. Bradley JD, Paulus R, Komaki R, et al. Standard-dose versus high-dose conformal radiotherapy with concurrent and consolidation carboplatin plus paclitaxel with or without cetuximab for patients with stage IIIA or IIIB non-small-cell lung cancer (RTOG 0617): a randomised, two-by-two factorial phase 3 study. The Lancet Oncology 2015;16:187-199. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25601342>.

535. Cox JD. Are the results of RTOG 0617 mysterious? Int J Radiat Oncol Biol Phys 2012;82:1042-1044. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22284026>.

536. Bradley JD, Bae K, Graham MV, et al. Primary analysis of the phase II component of a phase I/II dose intensification study using three-dimensional conformal radiation therapy and concurrent chemotherapy for patients with inoperable non-small-cell lung cancer: RTOG 0117. J Clin Oncol 2010;28:2475-2480. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20368547>.

537. Bradley JD, Moughan J, Graham MV, et al. A phase I/II radiation dose escalation study with concurrent chemotherapy for patients with inoperable stages I to III non-small-cell lung cancer: phase I results of

RTOG 0117. Int J Radiat Oncol Biol Phys 2010;77:367-372. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20457350>.

538. Schild SE, Fan W, Stinchcombe TE, et al. Toxicity Related to Radiotherapy Dose and Targeting Strategy: A Pooled Analysis of Cooperative Group Trials of Combined Modality Therapy for Locally Advanced Non-Small Cell Lung Cancer. J Thorac Oncol 2019;14:298-303. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30292852>.

539. Kong FM, Ten Haken RK, Schipper MJ, et al. High-dose radiation improved local tumor control and overall survival in patients with inoperable/unresectable non-small-cell lung cancer: long-term results of a radiation dose escalation study. Int J Radiat Oncol Biol Phys 2005;63:324-333. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/16168827>.

540. Zhao L, West BT, Hayman JA, et al. High radiation dose may reduce the negative effect of large gross tumor volume in patients with medically inoperable early-stage non-small cell lung cancer. Int J Radiat Oncol Biol Phys 2007;68:103-110. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17363189>.

541. Wang L, Correa CR, Zhao L, et al. The effect of radiation dose and chemotherapy on overall survival in 237 patients with Stage III non-small-cell lung cancer. Int J Radiat Oncol Biol Phys 2009;73:1383-1390. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/18929449>.

542. Rosenman JG, Halle JS, Socinski MA, et al. High-dose conformal radiotherapy for treatment of stage IIIA/IIIB non-small-cell lung cancer: technical issues and results of a phase I/II trial. Int J Radiat Oncol Biol Phys 2002;54:348-356. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/12243807>.

543. Schild SE, McGinnis WL, Graham D, et al. Results of a Phase I trial of concurrent chemotherapy and escalating doses of radiation for unresectable non-small-cell lung cancer. Int J Radiat Oncol Biol Phys 2006;65:1106-1111. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/16730134>.



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

544. Schild SE, Pang HH, Fan W, et al. Exploring Radiotherapy Targeting Strategy and Dose: A Pooled Analysis of Cooperative Group Trials of Combined Modality Therapy for Stage III NSCLC. *J Thorac Oncol* 2018;13:1171-1182. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29689435>.

545. Socinski MA, Blackstock AW, Bogart JA, et al. Randomized phase II trial of induction chemotherapy followed by concurrent chemotherapy and dose-escalated thoracic conformal radiotherapy (74 Gy) in stage III non-small-cell lung cancer: CALGB 30105. *J Clin Oncol* 2008;26:2457-2463. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/18487565>.

546. Stinchcombe TE, Lee CB, Moore DT, et al. Long-term follow-up of a phase I/II trial of dose escalating three-dimensional conformal thoracic radiation therapy with induction and concurrent carboplatin and paclitaxel in unresectable stage IIIA/B non-small cell lung cancer. *J Thorac Oncol* 2008;3:1279-1285. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/18978563>.

547. Machtay M, Bae K, Movsas B, et al. Higher biologically effective dose of radiotherapy is associated with improved outcomes for locally advanced non-small cell lung carcinoma treated with chemoradiation: an analysis of the Radiation Therapy Oncology Group. *Int J Radiat Oncol Biol Phys* 2012;82:425-434. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20980108>.

548. Sun B, Brooks ED, Komaki RU, et al. 7-year follow-up after stereotactic ablative radiotherapy for patients with stage I non-small cell lung cancer: Results of a phase 2 clinical trial. *Cancer* 2017;123:3031-3039. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28346656>.

549. Taremi M, Hope A, Dahele M, et al. Stereotactic body radiotherapy for medically inoperable lung cancer: prospective, single-center study of 108 consecutive patients. *Int J Radiat Oncol Biol Phys* 2012;82:967-973. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21377293>.

550. Verma V, McMillan MT, Grover S, Simone CB, 2nd. Stereotactic body radiation therapy and the influence of chemotherapy on overall survival for large (≥ 5 centimeter) non-small cell lung cancer. *Int J Radiat Oncol Biol Phys* 2017;97:146-154. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27843029>.

551. Verma V, Shostrom VK, Kumar SS, et al. Multi-institutional experience of stereotactic body radiotherapy for large (≥ 5 centimeters) non-small cell lung tumors. *Cancer* 2017;123:688-696. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27741355>.

552. Allibhai Z, Taremi M, Bezjak A, et al. The impact of tumor size on outcomes after stereotactic body radiation therapy for medically inoperable early-stage non-small cell lung cancer. *Int J Radiat Oncol Biol Phys* 2013;87:1064-1070. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24210082>.

553. Genshaft SJ, Suh RD, Abtin F, et al. Society of Interventional Radiology Quality Improvement Standards on Percutaneous Ablation of Non-Small Cell Lung Cancer and Metastatic Disease to the Lungs. *J Vasc Interv Radiol* 2021;32:1242 e1241-1242 e1210. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/34000388>.

554. Abtin F, De Baere T, Dupuy DE, et al. Updates on current role and practice of lung ablation. *J Thorac Imaging* 2019;34:266-277. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31094898>.

555. Palussiere J, Chomy F, Savina M, et al. Radiofrequency ablation of stage IA non-small cell lung cancer in patients ineligible for surgery: results of a prospective multicenter phase II trial. *J Cardiothorac Surg* 2018;13:91. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30143031>.

556. Dupuy DE, Fernando HC, Hillman S, et al. Radiofrequency ablation of stage IA non-small cell lung cancer in medically inoperable patients: Results from the American College of Surgeons Oncology Group Z4033 (Alliance) trial. *Cancer* 2015;121:3491-3498. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26096694>.



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

557. Huang BY, Li XM, Song XY, et al. Long-term results of CT-guided percutaneous radiofrequency ablation of inoperable patients with stage Ia non-small cell lung cancer: A retrospective cohort study. *Int J Surg* 2018;53:143-150. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29555533>.

558. Ambrogi MC, Fanucchi O, Cioni R, et al. Long-term results of radiofrequency ablation treatment of stage I non-small cell lung cancer: a prospective intention-to-treat study. *J Thorac Oncol* 2011;6:2044-2051. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22052222>.

559. Bilal H, Mahmood S, Rajashanker B, Shah R. Is radiofrequency ablation more effective than stereotactic ablative radiotherapy in patients with early stage medically inoperable non-small cell lung cancer? *Interact Cardiovasc Thorac Surg* 2012;15:258-265. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22581864>.

560. Shirvani SM, Jiang J, Chang JY, et al. Comparative effectiveness of 5 treatment strategies for early-stage non-small cell lung cancer in the elderly. *Int J Radiat Oncol Biol Phys* 2012;84:1060-1070. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22975611>.

561. Gewanter RM, Rosenzweig KE, Chang JY, et al. ACR Appropriateness Criteria: nonsurgical treatment for non-small-cell lung cancer: good performance status/definitive intent. *Curr Probl Cancer* 2010;34:228-249. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20541060>.

562. Perez CA, Pajak TF, Rubin P, et al. Long-term observations of the patterns of failure in patients with unresectable non-oat cell carcinoma of the lung treated with definitive radiotherapy. Report by the Radiation Therapy Oncology Group. *Cancer* 1987;59:1874-1881. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/3032394>.

563. Bradley J, Bae K, Choi N, et al. A phase II comparative study of gross tumor volume definition with or without PET/CT fusion in dosimetric planning for non-small-cell lung cancer (NSCLC): primary analysis of Radiation Therapy Oncology Group (RTOG) 0515. *Int J Radiat Oncol Biol*

Phys 2012;82:435-441 e431. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21075551>.

564. Belderbos JS, Kepka L, Spring Kong FM, et al. Report from the International Atomic Energy Agency (IAEA) consultants' meeting on elective nodal irradiation in lung cancer: non-small-Cell lung cancer (NSCLC). *Int J Radiat Oncol Biol Phys* 2008;72:335-342. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/18793953>.

565. Sanuki-Fujimoto N, Sumi M, Ito Y, et al. Relation between elective nodal failure and irradiated volume in non-small-cell lung cancer (NSCLC) treated with radiotherapy using conventional fields and doses. *Radiother Oncol* 2009;91:433-437. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19167118>.

566. Sulman EP, Komaki R, Klopp AH, et al. Exclusion of elective nodal irradiation is associated with minimal elective nodal failure in non-small cell lung cancer. *Radiat Oncol* 2009;4:5. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19183471>.

567. Rosenzweig KE, Sura S, Jackson A, Yorke E. Involved-field radiation therapy for inoperable non small-cell lung cancer. *J Clin Oncol* 2007;25:5557-5561. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17984185>.

568. Yuan S, Sun X, Li M, et al. A randomized study of involved-field irradiation versus elective nodal irradiation in combination with concurrent chemotherapy for inoperable stage III nonsmall cell lung cancer. *Am J Clin Oncol* 2007;30:239-244. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17551299>.

569. Fernandes AT, Shen J, Finlay J, et al. Elective nodal irradiation (ENI) vs. involved field radiotherapy (IFRT) for locally advanced non-small cell lung cancer (NSCLC): A comparative analysis of toxicities and clinical outcomes. *Radiother Oncol* 2010;95:178-184. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20356642>.

570. Chen M, Bao Y, Ma HL, et al. Involved-field radiotherapy versus elective nodal irradiation in combination with concurrent chemotherapy for



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

locally advanced non-small cell lung cancer: a prospective randomized study. Biomed Res Int 2013;2013:371819. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23762840>.

571. Albain KS, Rusch VW, Crowley JJ, et al. Concurrent cisplatin/etoposide plus chest radiotherapy followed by surgery for stages IIIA (N2) and IIIB non-small-cell lung cancer: mature results of Southwest Oncology Group phase II study 8805. J Clin Oncol 1995;13:1880-1892. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/7636530>.

572. Sher DJ, Fidler MJ, Seder CW, et al. Relationship Between Radiation Therapy Dose and Outcome in Patients Treated With Neoadjuvant Chemoradiation Therapy and Surgery for Stage IIIA Non-Small Cell Lung Cancer: A Population-Based, Comparative Effectiveness Analysis. Int J Radiat Oncol Biol Phys 2015;92:307-316. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25838187>.

573. Suntharalingam M, Paulus R, Edelman MJ, et al. Radiation therapy oncology group protocol 02-29: a phase II trial of neoadjuvant therapy with concurrent chemotherapy and full-dose radiation therapy followed by surgical resection and consolidative therapy for locally advanced non-small cell carcinoma of the lung. Int J Radiat Oncol Biol Phys 2012;84:456-463. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22543206>.

574. Kelsey CR, Light KL, Marks LB. Patterns of failure after resection of non-small-cell lung cancer: implications for postoperative radiation therapy volumes. Int J Radiat Oncol Biol Phys 2006;65:1097-1105. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/16682136>.

575. Corso CD, Rutter CE, Wilson LD, et al. Re-evaluation of the role of postoperative radiotherapy and the impact of radiation dose for non-small-cell lung cancer using the National Cancer Database. J Thorac Oncol 2015;10:148-155. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25325781>.

576. Lally BE, Zelterman D, Colasanto JM, et al. Postoperative radiotherapy for stage II or III non-small-cell lung cancer using the surveillance, epidemiology, and end results database. J Clin Oncol

2006;24:2998-3006. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/16769986>.

577. Spoelstra FO, Senan S, Le Pechoux C, et al. Variations in target volume definition for postoperative radiotherapy in stage III non-small-cell lung cancer: analysis of an international contouring study. Int J Radiat Oncol Biol Phys 2010;76:1106-1113. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19560881>.

578. Rodrigues G, Videtic GM, Sur R, et al. Palliative thoracic radiotherapy in lung cancer: An American Society for Radiation Oncology evidence-based clinical practice guideline. Pract Radiat Oncol 2011;1:60-71. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25740118>.

579. Rodrigues G, Macbeth F, Burmeister B, et al. Consensus statement on palliative lung radiotherapy: third international consensus workshop on palliative radiotherapy and symptom control. Clin Lung Cancer 2012;13:1-5. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21729656>.

580. Chen AB, Cronin A, Weeks JC, et al. Palliative radiation therapy practice in patients with metastatic non-small-cell lung cancer: a Cancer Care Outcomes Research and Surveillance Consortium (CanCORS) Study. J Clin Oncol 2013;31:558-564. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23295799>.

581. Chow E, Harris K, Fan G, et al. Palliative radiotherapy trials for bone metastases: a systematic review. J Clin Oncol 2007;25:1423-1436. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17416863>.

582. Lutz S, Berk L, Chang E, et al. Palliative radiotherapy for bone metastases: an ASTRO evidence-based guideline. Int J Radiat Oncol Biol Phys 2011;79:965-976. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21277118>.

583. Cross CK, Berman S, Buswell L, et al. Prospective study of palliative hypofractionated radiotherapy (8.5 Gy x 2) for patients with symptomatic non-small-cell lung cancer. Int J Radiat Oncol Biol Phys



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

2004;58:1098-1105. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/15001250>.

584. A Medical Research Council (MRC) randomised trial of palliative radiotherapy with two fractions or a single fraction in patients with inoperable non-small-cell lung cancer (NSCLC) and poor performance status. Medical Research Council Lung Cancer Working Party. Br J Cancer 1992;65:934-941. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/1377484>.

585. Koshy M, Malik R, Mahmood U, et al. Comparative effectiveness of aggressive thoracic radiation therapy and concurrent chemoradiation therapy in metastatic lung cancer. Pract Radiat Oncol 2015;5:374-382.

Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26412340>.

586. Ashworth AB, Senan S, Palma DA, et al. An individual patient data metaanalysis of outcomes and prognostic factors after treatment of oligometastatic non-small-cell lung cancer. Clin Lung Cancer 2014;15:346-355. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/24894943>.

587. Milano MT, Katz AW, Okunieff P. Patterns of recurrence after curative-intent radiation for oligometastases confined to one organ. Am J Clin Oncol 2010;33:157-163. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/19770627>.

588. Salama JK, Chmura SJ, Mehta N, et al. An initial report of a radiation dose-escalation trial in patients with one to five sites of metastatic disease. Clin Cancer Res 2008;14:5255-5259. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/18698045>.

589. Gomez DR, Blumenschein GR, Jr., Lee JJ, et al. Local consolidative therapy versus maintenance therapy or observation for patients with oligometastatic non-small-cell lung cancer without progression after first-line systemic therapy: a multicentre, randomised, controlled, phase 2 study. Lancet Oncol 2016;17:1672-1682. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/27789196>.

590. Iyengar P, Wardak Z, Gerber DE, et al. Consolidative Radiotherapy for Limited Metastatic Non-Small-Cell Lung Cancer: A Phase 2 Randomized Clinical Trial. JAMA Oncol 2018;4:e173501. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/28973074>.

591. Gomez DR, Tang C, Zhang J, et al. Local Consolidative Therapy Vs. Maintenance Therapy or Observation for Patients With Oligometastatic Non-Small-Cell Lung Cancer: Long-Term Results of a Multi-Institutional, Phase II, Randomized Study. J Clin Oncol 2019;37:1558-1565. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31067138>.

592. Petty WJ, Urbanic JJ, Ahmed T, et al. Long-Term Outcomes of a Phase 2 Trial of Chemotherapy With Consolidative Radiation Therapy for Oligometastatic Non-Small Cell Lung Cancer. Int J Radiat Oncol Biol Phys 2018;102:527-535. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/30003996>.

593. Falkson CB, Vella ET, Yu E, et al. Guideline for radiotherapy with curative intent in patients with early-stage medically inoperable non-small-cell lung cancer. Curr Oncol 2017;24:e44-e49. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/28270731>.

594. Dachele M, Senan S. The role of stereotactic ablative radiotherapy for early-stage and oligometastatic non-small cell lung cancer: evidence for changing paradigms. Cancer Res Treat 2011;43:75-82. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/21811422>.

595. Heinzerling JH, Kavanagh B, Timmerman RD. Stereotactic ablative radiation therapy for primary lung tumors. Cancer J 2011;17:28-32.

Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21263264>.

596. Potters L, Kavanagh B, Galvin JM, et al. American Society for Therapeutic Radiology and Oncology (ASTRO) and American College of Radiology (ACR) practice guideline for the performance of stereotactic body radiation therapy. Int J Radiat Oncol Biol Phys 2010;76:326-332.

Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20117285>.

597. Crabtree TD, Denlinger CE, Meyers BF, et al. Stereotactic body radiation therapy versus surgical resection for stage I non-small cell lung



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

cancer. J Thorac Cardiovasc Surg 2010;140:377-386. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20400121>.

598. Guckenberger M, Andratschke N, Alheit H, et al. Definition of stereotactic body radiotherapy: principles and practice for the treatment of stage I non-small cell lung cancer. Strahlenther Onkol 2014;190:26-33. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24052011>.

599. Onishi H, Shirato H, Nagata Y, et al. Stereotactic body radiotherapy (SBRT) for operable stage I non-small-cell lung cancer: can SBRT be comparable to surgery? Int J Radiat Oncol Biol Phys 2011;81:1352-1358. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20638194>.

600. Baumann P, Nyman J, Hoyer M, et al. Outcome in a prospective phase II trial of medically inoperable stage I non-small-cell lung cancer patients treated with stereotactic body radiotherapy. J Clin Oncol 2009;27:3290-3296. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19414667>.

601. Iyengar P, Westover K, Timmerman RD. Stereotactic ablative radiotherapy (SABR) for non-small cell lung cancer. Semin Respir Crit Care Med 2013;34:845-854. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24258574>.

602. Nagata Y, Hiraoka M, Shibata T, et al. Stereotactic Body Radiation Therapy For T1N0M0 Non-small Cell Lung Cancer: First Report for Inoperable Population of a Phase II Trial by Japan Clinical Oncology Group (JCOG 0403). International Journal of Radiation Oncology*Biophysics*Physics 2012;84:S46. Available at: [http://www.redjournal.org/article/S0360-3016\(12\)01274-6/abstract](http://www.redjournal.org/article/S0360-3016(12)01274-6/abstract).

603. Palma D, Visser O, Lagerwaard FJ, et al. Impact of introducing stereotactic lung radiotherapy for elderly patients with stage I non-small-cell lung cancer: a population-based time-trend analysis. J Clin Oncol 2010;28:5153-5159. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21041709>.

604. Widder J, Postmus D, Ubbels JF, et al. Survival and quality of life after stereotactic or 3D-conformal radiotherapy for inoperable early-stage

lung cancer. Int J Radiat Oncol Biol Phys 2011;81:e291-297. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21640503>.

605. Bradley JD, El Naqa I, Drzymala RE, et al. Stereotactic body radiation therapy for early-stage non-small-cell lung cancer: the pattern of failure is distant. Int J Radiat Oncol Biol Phys 2010;77:1146-1150. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19800181>.

606. Senti S, Lagerwaard FJ, Haasbeek CJ, et al. Patterns of disease recurrence after stereotactic ablative radiotherapy for early stage non-small-cell lung cancer: a retrospective analysis. Lancet Oncol 2012;13:802-809. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22727222>.

607. Fakiris AJ, McGarry RC, Yiannoutsos CT, et al. Stereotactic body radiation therapy for early-stage non-small-cell lung carcinoma: four-year results of a prospective phase II study. Int J Radiat Oncol Biol Phys 2009;75:677-682. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19251380>.

608. Chang JY, Senan S, Paul MA, et al. Stereotactic ablative radiotherapy versus lobectomy for operable stage I non-small-cell lung cancer: a pooled analysis of two randomised trials. Lancet Oncol 2015;16:630-637. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25981812>.

609. Shultz DB, Filippi AR, Thariat J, et al. Stereotactic ablative radiotherapy for pulmonary oligometastases and oligometastatic lung cancer. J Thorac Oncol 2014;9:1426-1433. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25170641>.

610. Filippi AR, Badellino S, Guarneri A, et al. Outcomes of single fraction stereotactic ablative radiotherapy for lung metastases. Technol Cancer Res Treat 2014;13:37-45. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23819496>.

611. Chan NK, Abdullah KG, Lubelski D, et al. Stereotactic radiosurgery for metastatic spine tumors. J Neurosurg Sci 2014;58:37-44. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24614791>.



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

612. Ojerholm E, Lee JY, Kolker J, et al. Gamma Knife radiosurgery to four or more brain metastases in patients without prior intracranial radiation or surgery. *Cancer Med* 2014;3:565-571. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24510602>.

613. Timmerman RD, Herman J, Cho LC. Emergence of stereotactic body radiation therapy and its impact on current and future clinical practice. *J Clin Oncol* 2014;32:2847-2854. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25113761>.

614. Salazar OM, Sandhu TS, Lattin PB, et al. Once-weekly, high-dose stereotactic body radiotherapy for lung cancer: 6-year analysis of 60 early-stage, 42 locally advanced, and 7 metastatic lung cancers. *Int J Radiat Oncol Biol Phys* 2008;72:707-715. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/18455322>.

615. Guckenberger M, Wulf J, Mueller G, et al. Dose-response relationship for image-guided stereotactic body radiotherapy of pulmonary tumors: relevance of 4D dose calculation. *Int J Radiat Oncol Biol Phys* 2009;74:47-54. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/18977095>.

616. Kunkler IH, Audisio R, Belkacemi Y, et al. Review of current best practice and priorities for research in radiation oncology for elderly patients with cancer: the International Society of Geriatric Oncology (SIOG) task force. *Ann Oncol* 2014;25:2134-2146. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24625455>.

617. MA IJ, Shoni M, Siegert C, et al. Survival After Stereotactic Body Radiation Therapy for Clinically Diagnosed or Biopsy-Proven Early-Stage NSCLC: A Systematic Review and Meta-Analysis. *J Thorac Oncol* 2019;14:583-595. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30721798>.

618. Rusthoven CG, Kavanagh BD, Karam SD. Improved survival with stereotactic ablative radiotherapy (SABR) over lobectomy for early stage non-small cell lung cancer (NSCLC): addressing the fallout of disruptive randomized data. *Ann Transl Med* 2015;3:149. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26244136>.

619. Berman AT, Jabbour SK, Vachani A, et al. Empiric Radiotherapy for Lung Cancer Collaborative Group multi-institutional evidence-based guidelines for the use of empiric stereotactic body radiation therapy for non-small cell lung cancer without pathologic confirmation. *Transl Lung Cancer Res* 2019;8:5-14. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30788230>.

620. Dautruche A, Filion E, Mathieu D, et al. To Biopsy or Not to Biopsy?: A Matched Cohort Analysis of Early-Stage Lung Cancer Treated with Stereotactic Radiation with or Without Histologic Confirmation. *Int J Radiat Oncol Biol Phys* 2020;107:88-97. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32004581>.

621. van den Berg LL, Klinkenberg TJ, Groen HJM, Widder J. Patterns of Recurrence and Survival after Surgery or Stereotactic Radiotherapy for Early Stage NSCLC. *J Thorac Oncol* 2015;10:826-831. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25629639>.

622. Versteegen NE, Oosterhuis JW, Palma DA, et al. Stage I-II non-small-cell lung cancer treated using either stereotactic ablative radiotherapy (SABR) or lobectomy by video-assisted thoracoscopic surgery (VATS): outcomes of a propensity score-matched analysis. *Ann Oncol* 2013;24:1543-1548. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23425947>.

623. Nagata Y, Hiraoka M, Shibata T, et al. A Phase II Trial of Stereotactic Body Radiation Therapy for Operable T1N0M0 Non-small Cell Lung Cancer: Japan Clinical Oncology Group (JCOG0403). *International Journal of Radiation Oncology*Biophysics* 2010;78:S27-S28. Available at: [http://www.redjournal.org/article/S0360-3016\(10\)01078-3/abstract](http://www.redjournal.org/article/S0360-3016(10)01078-3/abstract).

624. Lagerwaard FJ, Versteegen NE, Haasbeek CJ, et al. Outcomes of stereotactic ablative radiotherapy in patients with potentially operable stage I non-small cell lung cancer. *Int J Radiat Oncol Biol Phys* 2012;83:348-353. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22104360>.



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

625. Shirvani SM, Jiang J, Chang JY, et al. Lobectomy, sublobar resection, and stereotactic ablative radiotherapy for early-stage non-small cell lung cancers in the elderly. *JAMA Surg* 2014;149:1244-1253. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25321323>.

626. Timmerman RD, Paulus R, Pass HI, et al. RTOG 0618: Stereotactic body radiation therapy (SBRT) to treat operable early-stage lung cancer patients [abstract]. *J Clin Oncol* 2013;31(Suppl 15):Abstract 7523. Available at:

627. Brooks ED, Sun B, Feng L, et al. Association of Long-term Outcomes and Survival With Multidisciplinary Salvage Treatment for Local and Regional Recurrence After Stereotactic Ablative Radiotherapy for Early-Stage Lung Cancer. *JAMA Netw Open* 2018;1:e181390. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30646121>.

628. Matsuo Y, Shibuya K, Nagata Y, et al. Preliminary report of late recurrences, at 5 years or more, after stereotactic body radiation therapy for non-small cell lung cancer. *J Thorac Oncol* 2012;7:453-456. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22252562>.

629. Zhang X, Liu H, Balter P, et al. Positron emission tomography for assessing local failure after stereotactic body radiotherapy for non-small-cell lung cancer. *Int J Radiat Oncol Biol Phys* 2012;83:1558-1565. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22572078>.

630. Hoopes DJ, Tann M, Fletcher JW, et al. FDG-PET and stereotactic body radiotherapy (SBRT) for stage I non-small-cell lung cancer. *Lung Cancer* 2007;56:229-234. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17353064>.

631. Chen F, Matsuo Y, Yoshizawa A, et al. Salvage lung resection for non-small cell lung cancer after stereotactic body radiotherapy in initially operable patients. *J Thorac Oncol* 2010;5:1999-2002. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21102261>.

632. Neri S, Takahashi Y, Terashi T, et al. Surgical treatment of local recurrence after stereotactic body radiotherapy for primary and metastatic

lung cancers. *J Thorac Oncol* 2010;5:2003-2007. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21102262>.

633. Hearn JW, Videtic GM, Djemil T, Stephans KL. Salvage stereotactic body radiation therapy (SBRT) for local failure after primary lung SBRT. *Int J Radiat Oncol Biol Phys* 2014;90:402-406. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25017480>.

634. Trakul N, Harris JP, Le QT, et al. Stereotactic ablative radiotherapy for reirradiation of locally recurrent lung tumors. *J Thorac Oncol* 2012;7:1462-1465. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22895143>.

635. Kilburn JM, Kuremsky JG, Blackstock AW, et al. Thoracic re-irradiation using stereotactic body radiotherapy (SBRT) techniques as first or second course of treatment. *Radiother Oncol* 2014;110:505-510. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24444530>.

636. Zhao L, Zhou S, Balter P, et al. Planning Target Volume D95 and Mean Dose Should Be Considered for Optimal Local Control for Stereotactic Ablative Radiation Therapy. *Int J Radiat Oncol Biol Phys* 2016;95:1226-1235. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27209498>.

637. Baker R, Han G, Sarangkasiri S, et al. Clinical and dosimetric predictors of radiation pneumonitis in a large series of patients treated with stereotactic body radiation therapy to the lung. *Int J Radiat Oncol Biol Phys* 2013;85:190-195. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22929858>.

638. Chang JY, Bezjak A, Mornex F, Committee IART. Stereotactic ablative radiotherapy for centrally located early stage non-small-cell lung cancer: what we have learned. *J Thorac Oncol* 2015;10:577-585. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25514807>.

639. Chang JY, Li QQ, Xu QY, et al. Stereotactic ablative radiation therapy for centrally located early stage or isolated parenchymal recurrences of non-small cell lung cancer: how to fly in a "no fly zone". *Int J Radiat Oncol*



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

Biol Phys 2014;88:1120-1128. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/24661665>.

640. Hadziahmetovic M, Loo BW, Timmerman RD, et al. Stereotactic body radiation therapy (stereotactic ablative radiotherapy) for stage I non-small cell lung cancer--updates of radiobiology, techniques, and clinical outcomes. *Discov Med* 2010;9:411-417. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/20515609>.

641. Hara R, Itami J, Kondo T, et al. Clinical outcomes of single-fraction stereotactic radiation therapy of lung tumors. *Cancer* 2006;106:1347-1352. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/16475150>.

642. Chang JY, Balter PA, Dong L, et al. Stereotactic body radiation therapy in centrally and superiorly located stage I or isolated recurrent non-small-cell lung cancer. *Int J Radiat Oncol Biol Phys* 2008;72:967-971. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/18954709>.

643. Takeda A, Sanuki N, Kunieda E, et al. Stereotactic body radiotherapy for primary lung cancer at a dose of 50 Gy total in five fractions to the periphery of the planning target volume calculated using a superposition algorithm. *Int J Radiat Oncol Biol Phys* 2009;73:442-448. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/18990507>.

644. Stephans KL, Djemil T, Reddy CA, et al. A comparison of two stereotactic body radiation fractionation schedules for medically inoperable stage I non-small cell lung cancer: the Cleveland Clinic experience. *J Thorac Oncol* 2009;4:976-982. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19633473>.

645. Jin JY, Kong FM, Chetty IJ, et al. Impact of fraction size on lung radiation toxicity: hypofractionation may be beneficial in dose escalation of radiotherapy for lung cancers. *Int J Radiat Oncol Biol Phys* 2010;76:782-788. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19577855>.

646. Onishi H, Shirato H, Nagata Y, et al. Hypofractionated stereotactic radiotherapy (HypoFXSRT) for stage I non-small cell lung cancer: updated results of 257 patients in a Japanese multi-institutional study. *J Thorac*

Oncol 2007;2:S94-100. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/17603311>.

647. Lagerwaard FJ, Haasbeek CJ, Smit EF, et al. Outcomes of risk-adapted fractionated stereotactic radiotherapy for stage I non-small-cell lung cancer. *Int J Radiat Oncol Biol Phys* 2008;70:685-692. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/18164849>.

648. Liu MB, Eclov NC, Trakul N, et al. Clinical impact of dose overestimation by effective path length calculation in stereotactic ablative radiation therapy of lung tumors. *Pract Radiat Oncol* 2013;3:294-300. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24674401>.

649. Xiao Y, Papiez L, Paulus R, et al. Dosimetric evaluation of heterogeneity corrections for RTOG 0236: stereotactic body radiotherapy of inoperable stage I-II non-small-cell lung cancer. *Int J Radiat Oncol Biol Phys* 2009;73:1235-1242. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19251095>.

650. Timmerman R, McGarry R, Yiannoutsos C, et al. Excessive toxicity when treating central tumors in a phase II study of stereotactic body radiation therapy for medically inoperable early-stage lung cancer. *J Clin Oncol* 2006;24:4833-4839. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17050868>.

651. Chaudhuri AA, Tang C, Binkley MS, et al. Stereotactic ablative radiotherapy (SABR) for treatment of central and ultra-central lung tumors. *Lung Cancer* 2015;89:50-56. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25997421>.

652. Haseltine JM, Rimner A, Gelblum DY, et al. Fatal complications after stereotactic body radiation therapy for central lung tumors abutting the proximal bronchial tree. *Pract Radiat Oncol* 2016;6:e27-33. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26577006>.

653. Bezjak A, Paulus R, Gaspar LE, et al. Safety and Efficacy of a Five-Fraction Stereotactic Body Radiotherapy Schedule for Centrally Located Non-Small-Cell Lung Cancer: NRG Oncology/RTOG 0813 Trial. *J*



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

Clin Oncol 2019;37:1316-1325. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/30943123>.

654. Bezjak A, Paulus R, Gaspar LE, et al. Primary Study Endpoint Analysis for NRG Oncology/RTOG 0813 Trial of Stereotactic Body Radiation Therapy (SBRT) for Centrally Located Non-Small Cell Lung Cancer (NSCLC). International Journal of Radiation Oncology*Biophysics*Physics 2016;94:5-6. Available at:
<http://dx.doi.org/10.1016/j.ijrobp.2015.10.040>.

655. Fleckenstein J, Petroff A, Schafers HJ, et al. Long-term outcomes in radically treated synchronous vs. metachronous oligometastatic non-small-cell lung cancer. BMC Cancer 2016;16:348. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/27255302>.

656. De Ruysscher D, Wanders R, van Baardwijk A, et al. Radical treatment of non-small-cell lung cancer patients with synchronous oligometastases: long-term results of a prospective phase II trial (Nct01282450). J Thorac Oncol 2012;7:1547-1555. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/22982655>.

657. Chawla S, Chen Y, Katz AW, et al. Stereotactic body radiotherapy for treatment of adrenal metastases. Int J Radiat Oncol Biol Phys 2009;75:71-75. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/19250766>.

658. Scorsetti M, Alongi F, Filippi AR, et al. Long-term local control achieved after hypofractionated stereotactic body radiotherapy for adrenal gland metastases: a retrospective analysis of 34 patients. Acta Oncol 2012;51:618-623. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/22263925>.

659. Weickhardt AJ, Scheier B, Burke JM, et al. Local ablative therapy of oligoprogressive disease prolongs disease control by tyrosine kinase inhibitors in oncogene-addicted non-small-cell lung cancer. J Thorac Oncol 2012;7:1807-1814. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/23154552>.

660. Gan GN, Weickhardt AJ, Scheier B, et al. Stereotactic radiation therapy can safely and durably control sites of extra-central nervous system oligoprogressive disease in anaplastic lymphoma kinase-positive lung cancer patients receiving crizotinib. Int J Radiat Oncol Biol Phys 2014;88:892-898. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/24462383>.

661. Iyengar P, Kavanagh BD, Wardak Z, et al. Phase II trial of stereotactic body radiation therapy combined with erlotinib for patients with limited but progressive metastatic non-small-cell lung cancer. J Clin Oncol 2014;32:3824-3830. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/25349291>.

662. Takeda M, Okamoto I, Nakagawa K. Clinical impact of continued crizotinib administration after isolated central nervous system progression in patients with lung cancer positive for ALK rearrangement. J Thorac Oncol 2013;8:654-657. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/23584297>.

663. Bogart JA, Hodgson L, Seagren SL, et al. Phase I study of accelerated conformal radiotherapy for stage I non-small-cell lung cancer in patients with pulmonary dysfunction: CALGB 39904. J Clin Oncol 2010;28:202-206. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/19933904>.

664. Cheung P, Faria S, Ahmed S, et al. Phase II study of accelerated hypofractionated three-dimensional conformal radiotherapy for stage T1-3 N0 M0 non-small cell lung cancer: NCIC CTG BR.25. J Natl Cancer Inst 2014;106. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25074417>.

665. Sura S, Yorke E, Jackson A, Rosenzweig KE. High-dose radiotherapy for the treatment of inoperable non-small cell lung cancer. Cancer J 2007;13:238-242. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/17762758>.

666. Hu C, Chang EL, Hassenbusch SJ, 3rd, et al. Nonsmall cell lung cancer presenting with synchronous solitary brain metastasis. Cancer 2006;106:1998-2004. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/16572401>.



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

667. Li J, Bentzen SM, Renschler M, Mehta MP. Regression after whole-brain radiation therapy for brain metastases correlates with survival and improved neurocognitive function. *J Clin Oncol* 2007;25:1260-1266. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17401015>.

668. Aoyama H, Tago M, Kato N, et al. Neurocognitive function of patients with brain metastasis who received either whole brain radiotherapy plus stereotactic radiosurgery or radiosurgery alone. *Int J Radiat Oncol Biol Phys* 2007;68:1388-1395. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17674975>.

669. Wolfson AH, Bae K, Komaki R, et al. Primary analysis of a phase II randomized trial Radiation Therapy Oncology Group (RTOG) 0212: impact of different total doses and schedules of prophylactic cranial irradiation on chronic neurotoxicity and quality of life for patients with limited-disease small-cell lung cancer. *Int J Radiat Oncol Biol Phys* 2011;81:77-84. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20800380>.

670. Sun A, Bae K, Gore EM, et al. Phase III trial of prophylactic cranial irradiation compared with observation in patients with locally advanced non-small-cell lung cancer: neurocognitive and quality-of-life analysis. *J Clin Oncol* 2011;29:279-286. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21135267>.

671. Tallet AV, Azria D, Barlesi F, et al. Neurocognitive function impairment after whole brain radiotherapy for brain metastases: actual assessment. *Radiat Oncol* 2012;7:77. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22640600>.

672. Chang EL, Wefel JS, Hess KR, et al. Neurocognition in patients with brain metastases treated with radiosurgery or radiosurgery plus whole-brain irradiation: a randomised controlled trial. *Lancet Oncol* 2009;10:1037-1044. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19801201>.

673. Suh JH, Videtic GM, Aref AM, et al. ACR Appropriateness Criteria: single brain metastasis. *Curr Probl Cancer* 2010;34:162-174. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20541055>.

674. Marsh JC, Gielda BT, Herskovic AM, Abrams RA. Cognitive Sparing during the Administration of Whole Brain Radiotherapy and Prophylactic Cranial Irradiation: Current Concepts and Approaches. *J Oncol* 2010;2010:198208. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20671962>.

675. Gondi V, Pugh SL, Tome WA, et al. Preservation of memory with conformal avoidance of the hippocampal neural stem-cell compartment during whole-brain radiotherapy for brain metastases (RTOG 0933): a phase II multi-institutional trial. *J Clin Oncol* 2014;32:3810-3816. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25349290>.

676. Mulvenna P, Nankivell M, Barton R, et al. Dexamethasone and supportive care with or without whole brain radiotherapy in treating patients with non-small cell lung cancer with brain metastases unsuitable for resection or stereotactic radiotherapy (QUARTZ): results from a phase 3, non-inferiority, randomised trial. *Lancet* 2016;388:2004-2014. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27604504>.

677. Chen L, Douglass J, Kleinberg L, et al. Concurrent Immune Checkpoint Inhibitors and Stereotactic Radiosurgery for Brain Metastases in Non-Small Cell Lung Cancer, Melanoma, and Renal Cell Carcinoma. *Int J Radiat Oncol Biol Phys* 2018;100:916-925. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29485071>.

678. Koenig JL, Shi S, Sborov K, et al. Adverse Radiation Effect and Disease Control in Patients Undergoing Stereotactic Radiosurgery and Immune Checkpoint Inhibitor Therapy for Brain Metastases. *World Neurosurg* 2019;126:e1399-e1411. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30902777>.

679. Kalkanis SN, Kondziolka D, Gaspar LE, et al. The role of surgical resection in the management of newly diagnosed brain metastases: a systematic review and evidence-based clinical practice guideline. *J Neurooncol* 2010;96:33-43. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19960230>.

680. Gaspar LE, Mehta MP, Patchell RA, et al. The role of whole brain radiation therapy in the management of newly diagnosed brain



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

metastases: a systematic review and evidence-based clinical practice guideline. J Neurooncol 2010;96:17-32. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19960231>.

681. Mintz A, Perry J, Spithoff K, et al. Management of single brain metastasis: a practice guideline. Curr Oncol 2007;14:131-143. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17710205>.

682. Patchell RA, Tibbs PA, Walsh JW, et al. A randomized trial of surgery in the treatment of single metastases to the brain. N Engl J Med 1990;322:494-500. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/2405271>.

683. Linskey ME, Andrews DW, Asher AL, et al. The role of stereotactic radiosurgery in the management of patients with newly diagnosed brain metastases: a systematic review and evidence-based clinical practice guideline. J Neurooncol 2010;96:45-68. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19960227>.

684. Aoyama H, Shirato H, Tago M, et al. Stereotactic radiosurgery plus whole-brain radiation therapy vs stereotactic radiosurgery alone for treatment of brain metastases: a randomized controlled trial. JAMA 2006;295:2483-2491. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/16757720>.

685. Abe E, Aoyama H. The role of whole brain radiation therapy for the management of brain metastases in the era of stereotactic radiosurgery. Curr Oncol Rep 2012;14:79-84. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22006098>.

686. Mehta MP, Paleologos NA, Mikkelsen T, et al. The role of chemotherapy in the management of newly diagnosed brain metastases: a systematic review and evidence-based clinical practice guideline. J Neurooncol 2010;96:71-83. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19960229>.

687. Ellis TL, Neal MT, Chan MD. The role of surgery, radiosurgery and whole brain radiation therapy in the management of patients with

metastatic brain tumors. Int J Surg Oncol 2012;2012:952345. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22312545>.

688. Patchell RA, Tibbs PA, Regine WF, et al. Postoperative radiotherapy in the treatment of single metastases to the brain: a randomized trial. JAMA 1998;280:1485-1489. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/9809728>.

689. Ammirati M, Cobbs CS, Linskey ME, et al. The role of retreatment in the management of recurrent/progressive brain metastases: a systematic review and evidence-based clinical practice guideline. J Neurooncol 2010;96:85-96. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19957016>.

690. Bradbury P, Sivajohanathan D, Chan A, et al. Postoperative Adjuvant Systemic Therapy in Completely Resected Non-Small-Cell Lung Cancer: A Systematic Review. Clin Lung Cancer 2017;18:259-273 e258. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28162945>.

691. Arriagada R, Bergman B, Dunant A, et al. Cisplatin-based adjuvant chemotherapy in patients with completely resected non-small-cell lung cancer. N Engl J Med 2004;350:351-360. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/14736927>.

692. Winton T, Livingston R, Johnson D, et al. Vinorelbine plus cisplatin vs. observation in resected non-small-cell lung cancer. N Engl J Med 2005;352:2589-2597. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/15972865>.

693. Douillard JY, Rosell R, De Lena M, et al. Adjuvant vinorelbine plus cisplatin versus observation in patients with completely resected stage IB-IIIA non-small-cell lung cancer (Adjuvant Navelbine International Trialist Association [ANITA]): a randomised controlled trial. Lancet Oncol 2006;7:719-727. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/16945766>.

694. Song WA, Zhou NK, Wang W, et al. Survival benefit of neoadjuvant chemotherapy in non-small cell lung cancer: an updated meta-analysis of



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

13 randomized control trials. J Thorac Oncol 2010;5:510-516. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20107424>.

695. Scagliotti GV, Pastorino U, Vansteenkiste JF, et al. Randomized phase III study of surgery alone or surgery plus preoperative cisplatin and gemcitabine in stages IB to IIIA non-small-cell lung cancer. J Clin Oncol 2012;30:172-178. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22124104>.

696. Depierre A, Milleron B, Moro-Sibilot D, et al. Preoperative chemotherapy followed by surgery compared with primary surgery in resectable stage I (except T1N0), II, and IIIa non-small-cell lung cancer. J Clin Oncol 2002;20:247-253. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/11773176>.

697. Rosell R, Gomez-Codina J, Camps C, et al. Preresectional chemotherapy in stage IIIA non-small-cell lung cancer: a 7-year assessment of a randomized controlled trial. Lung Cancer 1999;26:7-14. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/10574676>.

698. Roth JA, Fossella F, Komaki R, et al. A randomized trial comparing perioperative chemotherapy and surgery with surgery alone in resectable stage IIIA non-small-cell lung cancer. J Natl Cancer Inst 1994;86:673-680. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/8158698>.

699. Felip E, Rosell R, Maestre JA, et al. Preoperative chemotherapy plus surgery versus surgery plus adjuvant chemotherapy versus surgery alone in early-stage non-small-cell lung cancer. J Clin Oncol 2010;28:3138-3145. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20516435>.

700. Pisters KM, Vallieres E, Crowley JJ, et al. Surgery with or without preoperative paclitaxel and carboplatin in early-stage non-small-cell lung cancer: Southwest Oncology Group Trial S9900, an intergroup, randomized, phase III trial. J Clin Oncol 2010;28:1843-1849. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20231678>.

701. Westeel V, Quoix E, Puyraveau M, et al. A randomised trial comparing preoperative to perioperative chemotherapy in early-stage

non-small-cell lung cancer (IFCT 0002 trial). Eur J Cancer 2013;49:2654-2664. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23735703>.

702. Group NM-aC. Preoperative chemotherapy for non-small-cell lung cancer: a systematic review and meta-analysis of individual participant data. Lancet 2014;383:1561-1571. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24576776>.

703. Curran WJ, Jr., Paulus R, Langer CJ, et al. Sequential vs. concurrent chemoradiation for stage III non-small cell lung cancer: randomized phase III trial RTOG 9410. J Natl Cancer Inst 2011;103:1452-1460. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21903745>.

704. Auperin A, Le Pechoux C, Rolland E, et al. Meta-analysis of concomitant versus sequential radiochemotherapy in locally advanced non-small-cell lung cancer. J Clin Oncol 2010;28:2181-2190. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20351327>.

705. Socinski MA, Rosenman JG, Halle J, et al. Dose-escalating conformal thoracic radiation therapy with induction and concurrent carboplatin/paclitaxel in unresectable stage IIIA/B nonsmall cell lung carcinoma: a modified phase I/II trial. Cancer 2001;92:1213-1223. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/11571735>.

706. Furuse K, Fukuoka M, Kawahara M, et al. Phase III study of concurrent versus sequential thoracic radiotherapy in combination with mitomycin, vindesine, and cisplatin in unresectable stage III non-small-cell lung cancer. J Clin Oncol 1999;17:2692-2699. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/10561343>.

707. Rugo HS, Klein P, Melin SA, et al. Association Between Use of a Scalp Cooling Device and Alopecia After Chemotherapy for Breast Cancer. JAMA 2017;317:606-614. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28196257>.

708. Rugo HS, Melin SA, Voigt J. Scalp cooling with adjuvant/neoadjuvant chemotherapy for breast cancer and the risk of scalp metastases: systematic review and meta-analysis. Breast Cancer Res Treat



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

2017;163:199-205. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/28275922>.

709. Nangia J, Wang T, Osborne C, et al. Effect of a Scalp Cooling Device on Alopecia in Women Undergoing Chemotherapy for Breast Cancer: The SCALP Randomized Clinical Trial. JAMA 2017;317:596-605. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/28196254>.

710. Smetanay K, Junio P, Feisst M, et al. COOLHAIR: a prospective randomized trial to investigate the efficacy and tolerability of scalp cooling in patients undergoing (neo)adjuvant chemotherapy for early breast cancer. Breast Cancer Res Treat 2019;173:135-143. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/30255454>.

711. Lemieux J, Provencher L, Perron L, et al. No effect of scalp cooling on survival among women with breast cancer. Breast Cancer Res Treat 2015;149:263-268. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/25511368>.

712. Gandhi L, Rodriguez-Abreu D, Gadgeel S, et al. Pembrolizumab plus chemotherapy in metastatic non-small-cell lung cancer. N Engl J Med 2018;378:2078-2092. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/29658856>.

713. Langer CJ, Gadgeel SM, Borghaei H, et al. Carboplatin and pemetrexed with or without pembrolizumab for advanced, non-squamous non-small-cell lung cancer: a randomised, phase 2 cohort of the open-label KEYNOTE-021 study. Lancet Oncol 2016;17:1497-1508. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27745820>.

714. Socinski MA, Evans T, Gettinger S, et al. Treatment of stage IV non-small cell lung cancer: Diagnosis and management of lung cancer, 3rd ed: American College of Chest Physicians evidence-based clinical practice guidelines. Chest 2013;143:e341S-e368S. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/23649446>.

715. Azzoli CG, Temin S, Aliff T, et al. 2011 focused update of 2009 American Society of Clinical Oncology Clinical Practice Guideline Update on chemotherapy for stage IV non-small-cell lung cancer. J Clin Oncol

2011;29:3825-3831. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/21900105>.

716. Azzoli CG, Baker S, Jr., Temin S, et al. American Society of Clinical Oncology Clinical Practice Guideline update on chemotherapy for stage IV non-small-cell lung cancer. J Clin Oncol 2009;27:6251-6266. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/19917871>.

717. Group NM-AC. Chemotherapy in addition to supportive care improves survival in advanced non-small-cell lung cancer: a systematic review and meta-analysis of individual patient data from 16 randomized controlled trials. J Clin Oncol 2008;26:4617-4625. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/18678835>.

718. Souquet PJ, Chauvin F, Boissel JP, et al. Polychemotherapy in advanced non small cell lung cancer: a meta-analysis. Lancet 1993;342:19-21. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/8100290>.

719. Chemotherapy in non-small cell lung cancer: a meta-analysis using updated data on individual patients from 52 randomised clinical trials. Non-small Cell Lung Cancer Collaborative Group. BMJ 1995;311:899-909. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/7580546>.

720. Bakitas MA, El-Jawahri A, Farquhar M, et al. The TEAM approach to improving oncology outcomes by incorporating palliative care in practice. J Oncol Pract 2017;13:557-566. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/28898605>.

721. Temel JS, Greer JA, Muzikansky A, et al. Early palliative care for patients with metastatic non-small-cell lung cancer. N Engl J Med 2010;363:733-742. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/20818875>.

722. Yates P, Schofield P, Zhao I, Currow D. Supportive and palliative care for lung cancer patients. J Thorac Dis 2013;5 Suppl 5:S623-628. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24163753>.



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

723. Ford DW, Koch KA, Ray DE, Selecky PA. Palliative and end-of-life care in lung cancer: Diagnosis and management of lung cancer, 3rd ed: American College of Chest Physicians evidence-based clinical practice guidelines. Chest 2013;143:e498S-e512S. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23649453>.

724. Aizer AA, Chen MH, McCarthy EP, et al. Marital status and survival in patients with cancer. J Clin Oncol 2013;31:3869-3876. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24062405>.

725. Basch E, Barbera L, Kerrigan CL, Velikova G. Implementation of Patient-Reported Outcomes in Routine Medical Care. Am Soc Clin Oncol Educ Book 2018;38:122-134. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30231381>.

726. Stover AM, Tompkins Stricker C, Hammelef K, et al. Using Stakeholder Engagement to Overcome Barriers to Implementing Patient-reported Outcomes (PROs) in Cancer Care Delivery: Approaches From 3 Prospective Studies. Med Care 2019;57 Suppl 5 Suppl 1:S92-S99. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30985602>.

727. Basch EM, Deal AM, Dueck AC, et al. Overall survival results of a randomized trial assessing patient-reported outcomes for symptom monitoring during routine cancer treatment [abstract]. J Clin Oncol 2017;35:Abstract LBA2. Available at: <https://meetinglibrary.asco.org/record/147027/abstract>.

728. Magilligan DJ, Jr., Duvernoy C, Malik G, et al. Surgical approach to lung cancer with solitary cerebral metastasis: twenty-five years' experience. Ann Thorac Surg 1986;42:360-364. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/3767508>.

729. Arriagada R, Dunant A, Pignon JP, et al. Long-term results of the international adjuvant lung cancer trial evaluating adjuvant Cisplatin-based chemotherapy in resected lung cancer. J Clin Oncol 2010;28:35-42. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19933916>.

730. Butts CA, Ding K, Seymour L, et al. Randomized phase III trial of vinorelbine plus cisplatin compared with observation in completely

resected stage IB and II non-small-cell lung cancer: updated survival analysis of JBR-10. J Clin Oncol 2010;28:29-34. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19933915>.

731. Douillard JY, Tribodet H, Aubert D, et al. Adjuvant cisplatin and vinorelbine for completely resected non-small cell lung cancer: subgroup analysis of the Lung Adjuvant Cisplatin Evaluation. J Thorac Oncol 2010;5:220-228. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20027124>.

732. Kreuter M, Vansteenkiste J, Fischer JR, et al. Randomized phase 2 trial on refinement of early-stage NSCLC adjuvant chemotherapy with cisplatin and pemetrexed versus cisplatin and vinorelbine: the TREAT study. Ann Oncol 2013;24:986-992. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23161898>.

733. Petrelli F, Barni S. Non-cancer-related mortality after cisplatin-based adjuvant chemotherapy for non-small cell lung cancer: a study-level meta-analysis of 16 randomized trials. Med Oncol 2013;30:641. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23813019>.

734. Pignon JP, Tribodet H, Scagliotti GV, et al. Lung adjuvant cisplatin evaluation: a pooled analysis by the LACE Collaborative Group. J Clin Oncol 2008;26:3552-3559. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/18506026>.

735. Wisnivesky JP, Smith CB, Packer S, et al. Survival and risk of adverse events in older patients receiving postoperative adjuvant chemotherapy for resected stages II-IIIa lung cancer: observational cohort study. BMJ 2011;343:d4013. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21757436>.

736. Strauss GM, Herndon J, Maddaus MA, et al. Randomized clinical trial of adjuvant chemotherapy with paclitaxel and carboplatin following resection in stage IB non-small cell lung cancer (NSCLC): Report of Cancer and Leukemia Group B (CALGB) Protocol 9633 [abstract]. J Clin Oncol 2004;22 (Suppl 14):Abstract 7019. Available at: http://meeting.ascopubs.org/cgi/content/abstract/22/14_suppl/7019.



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

737. Strauss GM, Herndon JE, II, Maddaus MA, et al. Adjuvant chemotherapy in stage IB non-small cell lung cancer (NSCLC): Update of Cancer and Leukemia Group B (CALGB) protocol 9633 [abstract]. J Clin Oncol 2006;24 (Suppl 18):Abstract 7007. Available at: http://meeting.ascopubs.org/cgi/content/abstract/24/18_suppl/7007.

738. Strauss GM, Herndon JE, 2nd, Maddaus MA, et al. Adjuvant paclitaxel plus carboplatin compared with observation in stage IB non-small-cell lung cancer: CALGB 9633 with the Cancer and Leukemia Group B, Radiation Therapy Oncology Group, and North Central Cancer Treatment Group Study Groups. J Clin Oncol 2008;26:5043-5051. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/18809614>.

739. Ohe Y, Ohashi Y, Kubota K, et al. Randomized phase III study of cisplatin plus irinotecan versus carboplatin plus paclitaxel, cisplatin plus gemcitabine, and cisplatin plus vinorelbine for advanced non-small-cell lung cancer: Four-Arm Cooperative Study in Japan. Ann Oncol 2007;18:317-323. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17079694>.

740. Katz A, Saad ED. CALGB 9633: an underpowered trial with a methodologically questionable conclusion. J Clin Oncol 2009;27:2300-2301; author reply 2301-2302. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19332712>.

741. Kreuter M, Vansteenkiste J, Fischer JR, et al. Three-Year Follow-Up of a Randomized Phase II Trial on Refinement of Early-Stage NSCLC Adjuvant Chemotherapy with Cisplatin and Pemetrexed versus Cisplatin and Vinorelbine (the TREAT Study). J Thorac Oncol 2016;11:85-93. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26762743>.

742. Kris MG, Gaspar LE, Chaft JE, et al. Adjuvant Systemic Therapy and Adjuvant Radiation Therapy for Stage I to IIIA Completely Resected Non-Small-Cell Lung Cancers: American Society of Clinical Oncology/Cancer Care Ontario Clinical Practice Guideline Update. J Clin Oncol 2017;35:2960-2974. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28437162>.

743. Perol M, Chouaid C, Perol D, et al. Randomized, phase III study of gemcitabine or erlotinib maintenance therapy versus observation, with predefined second-line treatment, after cisplatin-gemcitabine induction chemotherapy in advanced non-small-cell lung cancer. J Clin Oncol 2012;30:3516-3524. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22949150>.

744. Fossella F, Pereira JR, von Pawel J, et al. Randomized, multinational, phase III study of docetaxel plus platinum combinations versus vinorelbine plus cisplatin for advanced non-small-cell lung cancer: the TAX 326 study group. J Clin Oncol 2003;21:3016-3024. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/12837811>.

745. Usami N, Yokoi K, Hasegawa Y, et al. Phase II study of carboplatin and gemcitabine as adjuvant chemotherapy in patients with completely resected non-small cell lung cancer: a report from the Central Japan Lung Study Group, CJLSG 0503 trial. Int J Clin Oncol 2010;15:583-587. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20714770>.

746. Zhang L, Ou W, Liu Q, et al. Pemetrexed plus carboplatin as adjuvant chemotherapy in patients with curative resected non-squamous non-small cell lung cancer. Thorac Cancer 2014;5:50-56. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26766972>.

747. Schmid-Bindert G, Engel-Riedel W, Reck M, et al. A randomized Phase 2 study of pemetrexed in combination with cisplatin or carboplatin as adjuvant chemotherapy in patients with completely resected stage IB or II Non-Small-Cell Lung Cancer. Lung Cancer 2015;90:397-404. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26791798>.

748. Kim YH, Hirabayashi M, Togashi Y, et al. Phase II study of carboplatin and pemetrexed in advanced non-squamous, non-small-cell lung cancer: Kyoto Thoracic Oncology Research Group Trial 0902. Cancer Chemother Pharmacol 2012;70:271-276. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22752216>.

749. Wu YL, Tsuboi M, He J, et al. Osimertinib in resected EGFR-mutated non-small-cell lung cancer. N Engl J Med 2020;383:1711-1723. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32955177>.



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

750. Wu YL, John T, Grohe C, et al. Postoperative chemotherapy use and outcomes from ADAURA: osimertinib as adjuvant therapy for resected EGFR-mutated NSCLC. *J Thorac Oncol* 2022;17:423-433. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/34740861>.

751. Felip E, Altorki N, Zhou C, et al. Adjuvant atezolizumab after adjuvant chemotherapy in resected stage IB-IIIA non-small-cell lung cancer (IMpower010): a randomised, multicentre, open-label, phase 3 trial. *Lancet* 2021;398:1344-1357. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/34555333>.

752. Leonetti A, Minari R, Boni L, et al. Phase II, open-label, single-arm, multicenter study to assess the activity and safety of alectinib as neoadjuvant treatment in surgically resectable stage III ALK-positive NSCLC: ALNEO Trial. *Clin Lung Cancer* 2021;22:473-477. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/33762169>.

753. Gu R, Shi Z, Duan T, Song M. Feasibility and safety of neoadjuvant alectinib in pulmonary invasive mucinous adenocarcinoma with ALK rearrangement: case report and literature review. *Onco Targets Ther* 2021;14:5107-5113. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/34707369>.

754. Hu Y, Ren S, Wang R, et al. Case Report: pathological complete response to neoadjuvant alectinib in a patient with resectable ALK-positive non-small cell lung cancer. *Front Pharmacol* 2022;13:816683. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/35873553>.

755. Bing Z, Jia Z, Wang Y, et al. Pathological complete response to neoadjuvant ceritinib of a crizotinib-resistant, stage IIIB non-small cell lung cancer with ALK rearrangement: A case report. *Thorac Cancer* 2021;12:2130-2133. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/34105864>.

756. de Scordilli M, Michelotti A, Bertoli E, et al. Targeted therapy and immunotherapy in early-stage non-small cell lung cancer: Current evidence and ongoing trials. *Int J Mol Sci* 2022;23. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/35806230>.

757. Chaft JE, Dagogo-Jack I, Santini FC, et al. Clinical outcomes of patients with resected, early-stage ALK-positive lung cancer. *Lung Cancer* 2018;122:67-71. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30032847>.

758. Gallina FT, Bertolaccini L, Forcella D, et al. Analysis of molecular biomarkers in resected early-stage non-small cells lung cancer: A narrative review. *Cancers (Basel)* 2022;14. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/35454856>.

759. Provencio M, Serna-Blasco R, Nadal E, et al. Overall survival and biomarker analysis of neoadjuvant nivolumab plus chemotherapy in operable stage IIIA non-small-cell lung cancer (NADIM phase II trial). *J Clin Oncol* 2022;40:2924-2933. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/35576508>.

760. Provencio M, Nadal E, Insa A, et al. Neoadjuvant chemotherapy and nivolumab in resectable non-small-cell lung cancer (NADIM): an open-label, multicentre, single-arm, phase 2 trial. *Lancet Oncol* 2020;21:1413-1422. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32979984>.

761. Daly ME, Singh N, Ismaila N, et al. Management of stage III non-small-cell lung cancer: ASCO Guideline. *J Clin Oncol* 2022;40:1356-1384. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/34936470>.

762. Dillman RO, Seagren SL, Propert KJ, et al. A randomized trial of induction chemotherapy plus high-dose radiation versus radiation alone in stage III non-small-cell lung cancer. *N Engl J Med* 1990;323:940-945. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/2169587>.

763. Le Chevalier T, Arriagada R, Quoix E, et al. Radiotherapy alone versus combined chemotherapy and radiotherapy in nonresectable non-small-cell lung cancer: first analysis of a randomized trial in 353 patients. *J Natl Cancer Inst* 1991;83:417-423. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/1847977>.



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

764. Schaake-Koning C, van den Bogaert W, Dalesio O, et al. Effects of concomitant cisplatin and radiotherapy on inoperable non-small-cell lung cancer. *N Engl J Med* 1992;326:524-530. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/1310160>.

765. Dillman RO, Seagren SL, Herndon J, et al. A randomized trial of induction chemotherapy plus high-dose radiation versus radiation alone in stage III non-small-cell lung cancer: Five-year follow-up of cancer and leukemia group B (CALGB) 8433 trial. *J Clin Oncol (Meeting Abstracts)* 1993;12:329. Available at:

766. Dillman RO, Herndon J, Seagren SL, et al. Improved survival in stage III non-small-cell lung cancer: seven-year follow-up of cancer and leukemia group B (CALGB) 8433 trial. *J Natl Cancer Inst* 1996;88:1210-1215. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/8780630>.

767. Liu T, Mu Y, Dang J, Li G. The role of postoperative radiotherapy for completely resected pIIIA-N2 non-small cell lung cancer patients with different clinicopathological features: a systemic review and meta-analysis. *J Cancer* 2019;10:3941-3949. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31417638>.

768. Hancock JG, Rosen JE, Antonicelli A, et al. Impact of adjuvant treatment for microscopic residual disease after non-small cell lung cancer surgery. *Ann Thorac Surg* 2015;99:406-413. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25528723>.

769. Burdett S, Stewart L, Group PM-a. Postoperative radiotherapy in non-small-cell lung cancer: update of an individual patient data meta-analysis. *Lung Cancer* 2005;47:81-83. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/15603857>.

770. Sause W, Kolesar P, Taylor SI, et al. Final results of phase III trial in regionally advanced unresectable non-small cell lung cancer: Radiation Therapy Oncology Group, Eastern Cooperative Oncology Group, and Southwest Oncology Group. *Chest* 2000;117:358-364. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/10669675>.

771. O'Rourke N, Roque IFM, Farre Bernado N, Macbeth F. Concurrent chemoradiotherapy in non-small cell lung cancer. *Cochrane Database Syst Rev* 2010:CD002140. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20556756>.

772. Baumann M, Herrmann T, Koch R, et al. Final results of the randomized phase III CHARTWEL-trial (ARO 97-1) comparing hyperfractionated-accelerated versus conventionally fractionated radiotherapy in non-small cell lung cancer (NSCLC). *Radiother Oncol* 2011;100:76-85. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21757247>.

773. Mauguen A, Le Pechoux C, Saunders MI, et al. Hyperfractionated or accelerated radiotherapy in lung cancer: an individual patient data meta-analysis. *J Clin Oncol* 2012;30:2788-2797. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22753901>.

774. Ramnath N, Dilling TJ, Harris LJ, et al. Treatment of stage III non-small cell lung cancer: Diagnosis and management of lung cancer, 3rd ed: American College of Chest Physicians evidence-based clinical practice guidelines. *Chest* 2013;143:e314S-e340S. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23649445>.

775. Atagi S, Kawahara M, Yokoyama A, et al. Thoracic radiotherapy with or without daily low-dose carboplatin in elderly patients with non-small-cell lung cancer: a randomised, controlled, phase 3 trial by the Japan Clinical Oncology Group (JCOG0301). *Lancet Oncol* 2012;13:671-678. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22622008>.

776. Atagi S, Mizusawa J, Ishikura S, et al. Chemoradiotherapy in elderly patients with non-small-cell lung cancer: Long-term follow-up of a randomized trial (JCOG0301). *Clin Lung Cancer* 2018;19:e619-e627. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29887243>.

777. Rosell R, Gatzemeier U, Betticher DC, et al. Phase III randomised trial comparing paclitaxel/carboplatin with paclitaxel/cisplatin in patients with advanced non-small-cell lung cancer: a cooperative multinational trial. *Ann Oncol* 2002;13:1539-1549. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/12377641>.



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

778. Senan S, Brade A, Wang LH, et al. PROCLAIM: Randomized Phase III Trial of Pemetrexed-Cisplatin or Etoposide-Cisplatin Plus Thoracic Radiation Therapy Followed by Consolidation Chemotherapy in Locally Advanced Nonsquamous Non-Small-Cell Lung Cancer. *J Clin Oncol* 2016;34:953-962. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26811519>.

779. Ezer N, Smith CB, Galsky MD, et al. Cisplatin vs. carboplatin-based chemoradiotherapy in patients >65 years of age with stage III non-small cell lung cancer. *Radiother Oncol* 2014;112:272-278. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25150635>.

780. Albain KS, Crowley JJ, Turrisi AT, 3rd, et al. Concurrent cisplatin, etoposide, and chest radiotherapy in pathologic stage IIIB non-small-cell lung cancer: a Southwest Oncology Group phase II study, SWOG 9019. *J Clin Oncol* 2002;20:3454-3460. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/12177106>.

781. Belani CP, Choy H, Bonomi P, et al. Combined chemoradiotherapy regimens of paclitaxel and carboplatin for locally advanced non-small-cell lung cancer: a randomized phase II locally advanced multi-modality protocol. *J Clin Oncol* 2005;23:5883-5891. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/16087941>.

782. Santana-Davila R, DeVossety K, Szabo A, et al. Cisplatin and etoposide versus carboplatin and paclitaxel with concurrent radiotherapy for stage III non-small-cell lung cancer: an analysis of Veterans Health Administration data. *J Clin Oncol* 2015;33:567-574. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25422491>.

783. Choy H, Gerber DE, Bradley JD, et al. Concurrent pemetrexed and radiation therapy in the treatment of patients with inoperable stage III non-small cell lung cancer: a systematic review of completed and ongoing studies. *Lung Cancer* 2015;87:232-240. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25650301>.

784. Govindan R, Bogart J, Stinchcombe T, et al. Randomized phase II study of pemetrexed, carboplatin, and thoracic radiation with or without cetuximab in patients with locally advanced unresectable non-small-cell

lung cancer: Cancer and Leukemia Group B trial 30407. *J Clin Oncol* 2011;29:3120-3125. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21747084>.

785. Vokes EE, Senan S, Treat JA, Iscoe NA. PROCLAIM: A phase III study of pemetrexed, cisplatin, and radiation therapy followed by consolidation pemetrexed versus etoposide, cisplatin, and radiation therapy followed by consolidation cytotoxic chemotherapy of choice in locally advanced stage III non-small-cell lung cancer of other than predominantly squamous cell histology. *Clin Lung Cancer* 2009;10:193-198. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19443340>.

786. Antonia SJ, Villegas A, Daniel D, et al. Overall survival with durvalumab after chemoradiotherapy in stage III NSCLC. *N Engl J Med* 2018;379:2342-2350. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30280658>.

787. Hui R, Ozguroglu M, Villegas A, et al. Patient-reported outcomes with durvalumab after chemoradiotherapy in stage III, unresectable non-small-cell lung cancer (PACIFIC): a randomised, controlled, phase 3 study. *Lancet Oncol* 2019;20:1670-1680. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31601496>.

788. Smit EF, van Meerbeeck JP, Lianes P, et al. Three-arm randomized study of two cisplatin-based regimens and paclitaxel plus gemcitabine in advanced non-small-cell lung cancer: a phase III trial of the European Organization for Research and Treatment of Cancer Lung Cancer Group--EORTC 08975. *J Clin Oncol* 2003;21:3909-3917. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/14581415>.

789. Zatloukal P, Petruzelka L, Zemanova M, et al. Concurrent versus sequential chemoradiotherapy with cisplatin and vinorelbine in locally advanced non-small cell lung cancer: a randomized study. *Lung Cancer* 2004;46:87-98. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/15364136>.

790. Scagliotti GV, Parikh P, von Pawel J, et al. Phase III study comparing cisplatin plus gemcitabine with cisplatin plus pemetrexed in



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

chemotherapy-naïve patients with advanced-stage non-small-cell lung cancer. J Clin Oncol 2008;26:3543-3551. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/18506025>.

791. Kelly K, Crowley J, Bunn PA, Jr., et al. Randomized phase III trial of paclitaxel plus carboplatin versus vinorelbine plus cisplatin in the treatment of patients with advanced non-small-cell lung cancer: a Southwest Oncology Group trial. J Clin Oncol 2001;19:3210-3218. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/11432888>.

792. Schiller JH, Harrington D, Belani CP, et al. Comparison of four chemotherapy regimens for advanced non-small-cell lung cancer. N Engl J Med 2002;346:92-98. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/11784875>.

793. Santana-Davila R, Szabo A, Arce-Lara C, et al. Cisplatin versus carboplatin-based regimens for the treatment of patients with metastatic lung cancer. An analysis of Veterans Health Administration data. J Thorac Oncol 2014;9:702-709. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24662458>.

794. Grossi F, Kubota K, Cappuzzo F, et al. Future scenarios for the treatment of advanced non-small cell lung cancer: focus on taxane-containing regimens. Oncologist 2010;15:1102-1112. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20930102>.

795. Leighl NB. Treatment paradigms for patients with metastatic non-small-cell lung cancer: first-, second-, and third-line. Curr Oncol 2012;19:S52-58. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22787411>.

796. de Marinis F, Rossi A, Di Maio M, et al. Treatment of advanced non-small-cell lung cancer: Italian Association of Thoracic Oncology (AIOT) clinical practice guidelines. Lung Cancer 2011;73:1-10. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21440325>.

797. Danson S, Middleton MR, O'Byrne KJ, et al. Phase III trial of gemcitabine and carboplatin versus mitomycin, ifosfamide, and cisplatin or mitomycin, vinblastine, and cisplatin in patients with advanced nonsmall

cell lung carcinoma. Cancer 2003;98:542-553. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/12879472>.

798. Booton R, Lorigan P, Anderson H, et al. A phase III trial of docetaxel/carboplatin versus mitomycin C/ifosfamide/cisplatin (MIC) or mitomycin C/vinblastine/cisplatin (MVP) in patients with advanced non-small-cell lung cancer: a randomised multicentre trial of the British Thoracic Oncology Group (BTOG1). Ann Oncol 2006;17:1111-1119. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/16603599>.

799. Gronberg BH, Bremnes RM, Flotten O, et al. Phase III study by the Norwegian lung cancer study group: pemetrexed plus carboplatin compared with gemcitabine plus carboplatin as first-line chemotherapy in advanced non-small-cell lung cancer. J Clin Oncol 2009;27:3217-3224. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19433683>.

800. D'Addario G, Pintilie M, Leighl NB, et al. Platinum-based versus non-platinum-based chemotherapy in advanced non-small-cell lung cancer: a meta-analysis of the published literature. J Clin Oncol 2005;23:2926-2936. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/15728229>.

801. Greco FA, Spigel DR, Kuzur ME, et al. Paclitaxel/Carboplatin/gemcitabine versus gemcitabine/vinorelbine in advanced non-small-cell lung cancer: a phase II/III study of the Minnie Pearl Cancer Research Network. Clin Lung Cancer 2007;8:483-487. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17922972>.

802. Herbst RS, Khuri FR, Lu C, et al. The novel and effective nonplatinum, nontaxane combination of gemcitabine and vinorelbine in advanced nonsmall cell lung carcinoma: potential for decreased toxicity and combination with biological therapy. Cancer 2002;95:340-353. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/12124835>.

803. Pujol JL, Breton JL, Gervais R, et al. Gemcitabine-docetaxel versus cisplatin-vinorelbine in advanced or metastatic non-small-cell lung cancer: a phase III study addressing the case for cisplatin. Ann Oncol 2005;16:602-610. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/15741225>.



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

804. Zornosa C, Vandergrift JL, Kalemkerian GP, et al. First-line systemic therapy practice patterns and concordance with NCCN guidelines for patients diagnosed with metastatic NSCLC treated at NCCN institutions. *J Natl Compr Canc Netw* 2012;10:847-856. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22773800>.

805. Pennell NA. Selection of chemotherapy for patients with advanced non-small cell lung cancer. *Cleve Clin J Med* 2012;79 Electronic Suppl 1:eS46-50. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22614966>.

806. Sandler A, Gray R, Perry MC, et al. Paclitaxel-carboplatin alone or with bevacizumab for non-small-cell lung cancer. *N Engl J Med* 2006;355:2542-2550. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17167137>.

807. Socinski MA, Schell MJ, Peterman A, et al. Phase III trial comparing a defined duration of therapy versus continuous therapy followed by second-line therapy in advanced-stage IIIB/IV non-small-cell lung cancer. *J Clin Oncol* 2002;20:1335-1343. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/11870177>.

808. Fidias P, Novello S. Strategies for prolonged therapy in patients with advanced non-small-cell lung cancer. *J Clin Oncol* 2010;28:5116-5123. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21041704>.

809. Scagliotti GV, Park K, Patil S, et al. Survival without toxicity for cisplatin plus pemetrexed versus cisplatin plus gemcitabine in chemonaive patients with advanced non-small cell lung cancer: a risk-benefit analysis of a large phase III study. *Eur J Cancer* 2009;45:2298-2303. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19473833>.

810. Edelman MJ, Le Chevalier T, Soria JC. Maintenance therapy and advanced non-small-cell lung cancer: a skeptic's view. *J Thorac Oncol* 2012;7:1331-1336. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22895137>.

811. Hershman DL, Lacchetti C, Dworkin RH, et al. Prevention and management of chemotherapy-induced peripheral neuropathy in survivors of adult cancers: American Society of Clinical Oncology clinical practice

guideline. *J Clin Oncol* 2014;32:1941-1967. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24733808>.

812. Patel JD, Socinski MA, Garon EB, et al. PointBreak: a randomized phase III study of pemetrexed plus carboplatin and bevacizumab followed by maintenance pemetrexed and bevacizumab versus paclitaxel plus carboplatin and bevacizumab followed by maintenance bevacizumab in patients with stage IIIB or IV nonsquamous non-small-cell lung cancer. *J Clin Oncol* 2013;31:4349-4357. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24145346>.

813. Zhu J, Sharma DB, Gray SW, et al. Carboplatin and paclitaxel with vs without bevacizumab in older patients with advanced non-small cell lung cancer. *JAMA* 2012;307:1593-1601. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22511687>.

814. Langer C, Ravelo A, Hazard SJ, et al. Comparison of survival and hospitalization rates between Medicare patients with advanced NSCLC treated with bevacizumab-carboplatin-paclitaxel and carboplatin-paclitaxel: a retrospective cohort study. *Lung Cancer* 2014;86:350-357. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25439437>.

815. Langer CJ, Socinski MA, Patel JD, et al. Isolating the Role of Bevacizumab in Elderly Patients With Previously Untreated Nonsquamous Non-Small Cell Lung Cancer: Secondary Analyses of the ECOG 4599 and PointBreak Trials. *Am J Clin Oncol* 2016;39:441-447. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25628268>.

816. Rizvi NA, Riely GJ, Azzoli CG, et al. Phase I/II trial of weekly intravenous 130-nm albumin-bound paclitaxel as initial chemotherapy in patients with stage IV non-small-cell lung cancer. *J Clin Oncol* 2008;26:639-643. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/18235124>.

817. Green MR, Manikhas GM, Orlov S, et al. Abraxane, a novel Cremophor-free, albumin-bound particle form of paclitaxel for the treatment of advanced non-small-cell lung cancer. *Ann Oncol* 2006;17:1263-1268. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/16740598>.



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

818. Socinski MA, Bondarenko I, Karaseva NA, et al. Weekly nab-paclitaxel in combination with carboplatin versus solvent-based paclitaxel plus carboplatin as first-line therapy in patients with advanced non-small-cell lung cancer: final results of a phase III trial. *J Clin Oncol* 2012;30:2055-2062. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/22547591>.

819. Felip E, Gridelli C, Baas P, et al. Metastatic non-small-cell lung cancer: consensus on pathology and molecular tests, first-line, second-line, and third-line therapy: 1st ESMO Consensus Conference in Lung Cancer; Lugano 2010. *Ann Oncol* 2011;22:1507-1519. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/21536661>.

820. Lilenbaum R, Zukin M, Pereira JR, et al. A randomized phase III trial of single-agent pemetrexed (P) versus carboplatin and pemetrexed (CP) in patients with advanced non-small cell lung cancer (NSCLC) and performance status (PS) of 2 [abstract]. *J Clin Oncol* 2012;30(Suppl 15):Abstract 7506. Available at:

http://meeting.ascopubs.org/cgi/content/abstract/30/15_suppl/7506.

821. Langer CJ, O'Byrne KJ, Socinski MA, et al. Phase III trial comparing paclitaxel poliglumex (CT-2103, PPX) in combination with carboplatin versus standard paclitaxel and carboplatin in the treatment of PS 2 patients with chemotherapy-naïve advanced non-small cell lung cancer. *J Thorac Oncol* 2008;3:623-630. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/18520802>.

822. Lilenbaum R, Villalflor VM, Langer C, et al. Single-agent versus combination chemotherapy in patients with advanced non-small cell lung cancer and a performance status of 2: prognostic factors and treatment selection based on two large randomized clinical trials. *J Thorac Oncol* 2009;4:869-874. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/19487960>.

823. Roth BJ, Krilov L, Adams S, et al. Clinical cancer advances 2012: annual report on progress against cancer from the american society of clinical oncology. *J Clin Oncol* 2013;31:131-161. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/23213095>.

824. Zukin M, Barrios CH, Pereira JR, et al. Randomized phase III trial of single-agent pemetrexed versus carboplatin and pemetrexed in patients with advanced non-small-cell lung cancer and Eastern Cooperative Oncology Group performance status of 2. *J Clin Oncol* 2013;31:2849-2853. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/23775961>.

825. Thatcher N, Hirsch FR, Luft AV, et al. Necitumumab plus gemcitabine and cisplatin versus gemcitabine and cisplatin alone as first-line therapy in patients with stage IV squamous non-small-cell lung cancer (SQUIRE): an open-label, randomised, controlled phase 3 trial. *Lancet Oncol* 2015;16:763-774. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/26045340>.

826. Goldstein DA, Chen Q, Ayer T, et al. Necitumumab in Metastatic Squamous Cell Lung Cancer: Establishing a Value-Based Cost. *JAMA Oncol* 2015;1:1293-1300. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/26313558>.

827. Sandler AB, Johnson DH, Herbst RS. Anti-vascular endothelial growth factor monoclonals in non-small cell lung cancer. *Clin Cancer Res* 2004;10:4258s-4262s. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/15217970>.

828. Giaccone G. Epidermal growth factor receptor inhibitors in the treatment of non-small-cell lung cancer. *J Clin Oncol* 2005;23:3235-3242. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/15886311>.

829. Lin JJ, Kennedy E, Sequist LV, et al. Clinical Activity of Alectinib in Advanced RET-Rearranged Non-Small Cell Lung Cancer. *J Thorac Oncol* 2016;11:2027-2032. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/27544060>.

830. Sabari JK, Santini FC, Schram AM, et al. The activity, safety, and evolving role of brigatinib in patients with ALK-rearranged non-small cell lung cancers. *Onco Targets Ther* 2017;10:1983-1992. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/28435288>.



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

831. Solomon BJ, Besse B, Bauer TM, et al. Lorlatinib in patients with ALK-positive non-small-cell lung cancer: results from a global phase 2 study. *Lancet Oncol* 2018;19:1654-1667. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30413378>.

832. Besse B, Solomon BJ, Felip E, et al. Lorlatinib in patients (Pts) with previously treated ALK+ advanced non-small cell lung cancer (NSCLC): Updated efficacy and safety [abstract]. *J Clin Oncol* 2018;36(15_suppl):Abstract 9032. Available at: http://ascopubs.org/doi/abs/10.1200/JCO.2018.36.15_suppl.9032.

833. Shaw AT, Solomon BJ, Besse B, et al. ALK Resistance Mutations and Efficacy of Lorlatinib in Advanced Anaplastic Lymphoma Kinase-Positive Non-Small-Cell Lung Cancer. *J Clin Oncol* 2019;37:1370-1379. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30892989>.

834. Lassen U, Albert CM, Kummar S, et al. Larotrectinib efficacy and safety in TRK fusion cancer: an expanded clinical dataset showing consistency in an age and tumor agnostic approach [abstract] [abstract]. Presented at the ESMO Congress; October 19-23; Munich, Germany. Abstract 4090.

835. Paik PK, Felip E, Veillon R, et al. Tepotinib in Non-Small-Cell Lung Cancer with MET Exon 14 Skipping Mutations. *N Engl J Med* 2020;383:931-943. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32469185>.

836. Riely GJ, Kris MG, Zhao B, et al. Prospective assessment of discontinuation and reinitiation of erlotinib or gefitinib in patients with acquired resistance to erlotinib or gefitinib followed by the addition of everolimus. *Clin Cancer Res* 2007;13:5150-5155. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17785570>.

837. Kuriyama Y, Kim YH, Nagai H, et al. Disease flare after discontinuation of crizotinib in anaplastic lymphoma kinase-positive lung cancer. *Case Rep Oncol* 2013;6:430-433. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24019783>.

838. Chaft JE, Oxnard GR, Sima CS, et al. Disease flare after tyrosine kinase inhibitor discontinuation in patients with EGFR-mutant lung cancer and acquired resistance to erlotinib or gefitinib: implications for clinical trial design. *Clin Cancer Res* 2011;17:6298-6303. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21856766>.

839. Pop O, Pirvu A, Toffart AC, Moro-Sibilot D. Disease flare after treatment discontinuation in a patient with EML4-ALK lung cancer and acquired resistance to crizotinib. *J Thorac Oncol* 2012;7:e1-2. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22797152>.

840. Gainor JF, Shaw AT, Sequist LV, et al. EGFR Mutations and ALK Rearrangements Are Associated with Low Response Rates to PD-1 Pathway Blockade in Non-Small Cell Lung Cancer: A Retrospective Analysis. *Clin Cancer Res* 2016;22:4585-4593. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27225694>.

841. Hida T, Nokihara H, Kondo M, et al. Alectinib versus crizotinib in patients with ALK-positive non-small-cell lung cancer (J-ALEX): an open-label, randomised phase 3 trial. *Lancet* 2017;390:29-39. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28501140>.

842. Larkins E, Blumenthal GM, Chen H, et al. FDA Approval: Alectinib for the Treatment of Metastatic, ALK-Positive Non-Small Cell Lung Cancer Following Crizotinib. *Clin Cancer Res* 2016;22:5171-5176. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27413075>.

843. Shaw AT, Gandhi L, Gadgeel S, et al. Alectinib in ALK-positive, crizotinib-resistant, non-small-cell lung cancer: a single-group, multicentre, phase 2 trial. *Lancet Oncol* 2016;17:234-242. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26708155>.

844. Camidge DR, Kim HR, Ahn MJ, et al. Brigatinib versus crizotinib in advanced ALK inhibitor-naïve ALK-positive non-small cell lung cancer: Second interim analysis of the phase III ALTA-1L trial. *J Clin Oncol* 2020;38:3592-3603. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32780660>.



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

845. Camidge DR, Kim HR, Ahn MJ, et al. Brigatinib versus crizotinib in ALK inhibitor-naïve advanced ALK-positive NSCLC: final results of phase 3 ALTA-1L trial. *J Thorac Oncol* 2021;16:2091-2108. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/34537440>.

846. Kim DW, Tiseo M, Ahn MJ, et al. Brigatinib in Patients With Crizotinib-Refractory Anaplastic Lymphoma Kinase-Positive Non-Small-Cell Lung Cancer: A Randomized, Multicenter Phase II Trial. *J Clin Oncol* 2017;35:2490-2498. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28475456>.

847. Camidge DR, Tiseo M, Ahn M-J, et al. P3.02a-013 Brigatinib in crizotinib-refractory ALK+ NSCLC: central assessment and updates from ALTA, a pivotal randomized phase 2 trial [abstract]. *J Thorac Oncol* 2017;12:S1167-S1169. Available at: <http://bit.ly/2pCxxvKu>.

848. Lim SM, Kim HR, Lee JS, et al. Open-Label, Multicenter, Phase II Study of Ceritinib in Patients With Non-Small-Cell Lung Cancer Harboring ROS1 Rearrangement. *J Clin Oncol* 2017;35:2613-2618. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28520527>.

849. Shaw AT, Kim TM, Crino L, et al. Ceritinib versus chemotherapy in patients with ALK-rearranged non-small-cell lung cancer previously given chemotherapy and crizotinib (ASCEND-5): a randomised, controlled, open-label, phase 3 trial. *Lancet Oncol* 2017;18:874-886. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28602779>.

850. Crino L, Ahn MJ, De Marinis F, et al. Multicenter Phase II Study of Whole-Body and Intracranial Activity With Ceritinib in Patients With ALK-Rearranged Non-Small-Cell Lung Cancer Previously Treated With Chemotherapy and Crizotinib: Results From ASCEND-2. *J Clin Oncol* 2016;34:2866-2873. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27432917>.

851. Kazandjian D, Blumenthal GM, Chen HY, et al. FDA approval summary: crizotinib for the treatment of metastatic non-small cell lung cancer with anaplastic lymphoma kinase rearrangements. *Oncologist* 2014;19:e5-11. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25170012>.

852. Shaw AT, Kim DW, Nakagawa K, et al. Crizotinib versus chemotherapy in advanced ALK-positive lung cancer. *N Engl J Med* 2013;368:2385-2394. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23724913>.

853. Crino L, Kim D, Riely GJ, et al. Initial phase II results with crizotinib in advanced ALK-positive non-small cell lung cancer (NSCLC): PROFILE 1005 [abstract]. *J Clin Oncol* 2011;29 (Suppl 15):Abstract 7514. Available at: http://meeting.ascopubs.org/cgi/content/abstract/29/15_suppl/7514.

854. Camidge DR, Bang Y, Kwak EL, et al. Progression-free survival (PFS) from a phase I study of crizotinib (PF-02341066) in patients with ALK-positive non-small cell lung cancer (NSCLC) [abstract]. *J Clin Oncol* 2011;29(Suppl 15):Abstract 2501. Available at: http://meeting.ascopubs.org/cgi/content/abstract/29/15_suppl/2501.

855. Rodig SJ, Shapiro GI. Crizotinib, a small-molecule dual inhibitor of the c-Met and ALK receptor tyrosine kinases. *Curr Opin Investig Drugs* 2010;11:1477-1490. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21154129>.

856. Frampton JE. Crizotinib: a review of its use in the treatment of anaplastic lymphoma kinase-positive, advanced non-small cell lung cancer. *Drugs* 2013;73:2031-2051. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24288180>.

857. Costa DB, Shaw AT, Ou SH, et al. Clinical Experience With Crizotinib in Patients With Advanced ALK-Rearranged Non-Small-Cell Lung Cancer and Brain Metastases. *J Clin Oncol* 2015;33:1881-1888. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25624436>.

858. Camidge DR, Bang YJ, Kwak EL, et al. Activity and safety of crizotinib in patients with ALK-positive non-small-cell lung cancer: updated results from a phase 1 study. *Lancet Oncol* 2012;13:1011-1019. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22954507>.

859. Shaw AT, Yeap BY, Solomon BJ, et al. Impact of crizotinib on survival in patients with advanced, ALK-positive NSCLC compared with historical controls [abstract]. *J Clin Oncol* 2011;29(Suppl 15):Abstract



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

7507. Available at:

http://meeting.ascopubs.org/cgi/content/abstract/29/15_suppl/7507.

860. Bang YJ. Treatment of ALK-positive non-small cell lung cancer. Arch Pathol Lab Med 2012;136:1201-1204. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/23020724>.

861. Choi YL, Soda M, Yamashita Y, et al. EML4-ALK mutations in lung cancer that confer resistance to ALK inhibitors. N Engl J Med 2010;363:1734-1739. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/20979473>.

862. Rothenstein JM, Letarte N. Managing treatment-related adverse events associated with Alk inhibitors. Curr Oncol 2014;21:19-26. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/24523601>.

863. Brosnan EM, Weickhardt AJ, Lu X, et al. Drug-induced reduction in estimated glomerular filtration rate in patients with ALK-positive non-small cell lung cancer treated with the ALK inhibitor crizotinib. Cancer 2014;120:664-674. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/24258622>.

864. Gandhi L, Ou SI, Shaw AT, et al. Efficacy of alectinib in central nervous system metastases in crizotinib-resistant ALK-positive non-small-cell lung cancer: Comparison of RECIST 1.1 and RANO-HGG criteria. Eur J Cancer 2017;82:27-33. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/28646771>.

865. Camidge DR, Kim DW, Tiseo M, et al. Exploratory Analysis of Brigatinib Activity in Patients With Anaplastic Lymphoma Kinase-Positive Non-Small-Cell Lung Cancer and Brain Metastases in Two Clinical Trials. J Clin Oncol 2018;36:2693-2701. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/29768119>.

866. Solomon BJ, Bauer TM, Ignatius Ou SH, et al. Post hoc analysis of lorlatinib intracranial efficacy and safety in patients with ALK-positive advanced non-small-cell lung cancer from the phase III CROWN study. J Clin Oncol 2022;40:3593-3602. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/35605188>.

867. Planchard D, Smit EF, Groen HJM, et al. Dabrafenib plus trametinib in patients with previously untreated BRAF(V600E)-mutant metastatic non-small-cell lung cancer: an open-label, phase 2 trial. Lancet Oncol 2017;18:1307-1316. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/28919011>.

868. Planchard D, Besse B, Kim TM. Updated survival of patients (pts) with previously treated BRAF V600E-mutant advanced non-small cell lung cancer (NSCLC) who received dabrafenib (D) or D + trametinib (T) in the phase II BR113928 study [abstract]. J Clin Oncol 2017;35:Abstract 9075. Available at:

http://ascopubs.org/doi/abs/10.1200/JCO.2017.35.15_suppl.9075.

869. Planchard D, Groen HJM, Kim TM, et al. Interim results of a phase II study of the BRAF inhibitor (BRAFi) dabrafenib (D) in combination with the MEK inhibitor trametinib (T) in patients (pts) with BRAF V600E mutated (mut) metastatic non-small cell lung cancer (NSCLC) [abstract]. J Clin Oncol 2015;33:Abstract 8006. Available at:

<http://meetinglibrary.asco.org/content/147124-156>.

870. Odogwu L, Mathieu L, Blumenthal G, et al. FDA Approval Summary: Dabrafenib and Trametinib for the Treatment of Metastatic Non-Small Cell Lung Cancers Harboring BRAF V600E Mutations. Oncologist 2018;23:740-745. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/29438093>.

871. Planchard D, Kim TM, Mazieres J, et al. Dabrafenib in patients with BRAF(V600E)-positive advanced non-small-cell lung cancer: a single-arm, multicentre, open-label, phase 2 trial. Lancet Oncol 2016;17:642-650. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/27080216>.

872. Chalmers A, Cannon L, Akerley W. Adverse Event Management in Patients with BRAF V600E-Mutant Non-Small Cell Lung Cancer Treated with Dabrafenib plus Trametinib. Oncologist 2019;24:963-972. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/30598499>.

873. Nelson V, Ziehr J, Agulnik M, Johnson M. Afatinib: emerging next-generation tyrosine kinase inhibitor for NSCLC. Onco Targets Ther



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

2013;6:135-143. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/23493883>.

874. De Greve J, Teugels E, Geers C, et al. Clinical activity of afatinib (BIBW 2992) in patients with lung adenocarcinoma with mutations in the kinase domain of HER2/neu. Lung Cancer 2012;76:123-127. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/22325357>.

875. FDA approves afatinib for advanced lung cancer. Oncology (Williston Park) 2013;27:813-814. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/24133833>.

876. Dungo RT, Keating GM. Afatinib: first global approval. Drugs 2013;73:1503-1515. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/23982599>.

877. West H, Oxnard GR, Doebele RC. Acquired resistance to targeted therapies in advanced non-small cell lung cancer: new strategies and new agents. Am Soc Clin Oncol Educ Book 2013;272-278. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/23714521>.

878. Soria JC, Felip E, Cobo M, et al. Afatinib versus erlotinib as second-line treatment of patients with advanced squamous cell carcinoma of the lung (LUX-Lung 8): an open-label randomised controlled phase 3 trial. Lancet Oncol 2015;16:897-907. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/26156651>.

879. Yang JC, Sequist LV, Geater SL, et al. Clinical activity of afatinib in patients with advanced non-small-cell lung cancer harbouring uncommon EGFR mutations: a combined post-hoc analysis of LUX-Lung 2, LUX-Lung 3, and LUX-Lung 6. Lancet Oncol 2015;16:830-838. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/26051236>.

880. Park K, Tan EH, O'Byrne K, et al. Afatinib versus gefitinib as first-line treatment of patients with EGFR mutation-positive non-small-cell lung cancer (LUX-Lung 7): a phase 2B, open-label, randomised controlled trial. Lancet Oncol 2016;17:577-589. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/27083334>.

881. Paz-Ares L, Tan EH, O'Byrne K, et al. Afatinib versus gefitinib in patients with EGFR mutation-positive advanced non-small-cell lung cancer: overall survival data from the phase IIb LUX-Lung 7 trial. Ann Oncol 2017;28:270-277. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/28426106>.

882. Urata Y, Katakami N, Morita S, et al. Randomized Phase III Study Comparing Gefitinib With Erlotinib in Patients With Previously Treated Advanced Lung Adenocarcinoma: WJOG 5108L. J Clin Oncol 2016;34:3248-3257. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/27022112>.

883. Fukuoka M, Wu YL, Thongprasert S, et al. Biomarker analyses and final overall survival results from a phase III, randomized, open-label, first-line study of gefitinib versus carboplatin/paclitaxel in clinically selected patients with advanced non-small-cell lung cancer in Asia (IPASS). J Clin Oncol 2011;29:2866-2874. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/21670455>.

884. Kazandjian D, Blumenthal GM, Yuan W, et al. FDA Approval of Gefitinib for the Treatment of Patients with Metastatic EGFR Mutation-Positive Non-Small Cell Lung Cancer. Clin Cancer Res 2016;22:1307-1312. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/26980062>.

885. Janne PA, Wang X, Socinski MA, et al. Randomized phase II trial of erlotinib alone or with carboplatin and paclitaxel in patients who were never or light former smokers with advanced lung adenocarcinoma: CALGB 30406 trial. J Clin Oncol 2012;30:2063-2069. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/22547605>.

886. Masters GA, Temin S, Azzoli CG, et al. Systemic Therapy for Stage IV Non-Small-Cell Lung Cancer: American Society of Clinical Oncology Clinical Practice Guideline Update. J Clin Oncol 2015;33:3488-3515. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/26324367>.

887. Jackman DM, Miller VA, Cioffredi LA, et al. Impact of epidermal growth factor receptor and KRAS mutations on clinical outcomes in previously untreated non-small cell lung cancer patients: results of an



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

online tumor registry of clinical trials. Clin Cancer Res 2009;15:5267-5273. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19671843>.

888. Gridelli C, Ciardiello F, Gallo C, et al. First-line erlotinib followed by second-line cisplatin-gemcitabine chemotherapy in advanced non-small-cell lung cancer: the TORCH randomized trial. J Clin Oncol 2012;30:3002-3011. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22778317>.

889. Burotto M, Manasanch EE, Wilkerson J, Fojo T. Gefitinib and erlotinib in metastatic non-small cell lung cancer: a meta-analysis of toxicity and efficacy of randomized clinical trials. Oncologist 2015;20:400-410. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25795635>.

890. Haspinger ER, Agustoni F, Torri V, et al. Is there evidence for different effects among EGFR-TKIs? Systematic review and meta-analysis of EGFR tyrosine kinase inhibitors (TKIs) versus chemotherapy as first-line treatment for patients harboring EGFR mutations. Crit Rev Oncol Hematol 2015;94:213-227. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25523487>.

891. Nakagawa K, Garon EB, Seto T, et al. Ramucirumab plus erlotinib in patients with untreated, EGFR-mutated, advanced non-small-cell lung cancer (RELAY): a randomised, double-blind, placebo-controlled, phase 3 trial. Lancet Oncol 2019;20:1655-1669. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31591063>.

892. Kawashima Y, Fukuhara T, Saito H, et al. Bevacizumab plus erlotinib versus erlotinib alone in Japanese patients with advanced, metastatic, EGFR-mutant non-small-cell lung cancer (NEJ026): overall survival analysis of an open-label, randomised, multicentre, phase 3 trial. Lancet Respir Med 2022;10:72-82. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/34454653>.

893. Saito H, Fukuhara T, Furuya N, et al. Erlotinib plus bevacizumab versus erlotinib alone in patients with EGFR-positive advanced non-squamous non-small-cell lung cancer (NEJ026): interim analysis of an open-label, randomised, multicentre, phase 3 trial. Lancet Oncol

2019;20:625-635. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30975627>.

894. Sequist LV, Joshi VA, Janne PA, et al. Response to treatment and survival of patients with non-small cell lung cancer undergoing somatic EGFR mutation testing. Oncologist 2007;12:90-98. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17285735>.

895. Inoue A, Kobayashi K, Usui K, et al. First-line gefitinib for patients with advanced non-small-cell lung cancer harboring epidermal growth factor receptor mutations without indication for chemotherapy. J Clin Oncol 2009;27:1394-1400. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19224850>.

896. Thatcher N, Goldschmidt JH, Thomas M, et al. Efficacy and Safety of the Biosimilar ABP 215 Compared with Bevacizumab in Patients with Advanced Nonsquamous Non-small Cell Lung Cancer (MAPLE): A Randomized, Double-blind, Phase III Study. Clin Cancer Res 2019;25:2088-2095. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30617139>.

897. Reinmuth N, Bryl M, Bondarenko I, et al. PF-06439535 (a Bevacizumab Biosimilar) Compared with Reference Bevacizumab (Avastin((R))), Both Plus Paclitaxel and Carboplatin, as First-Line Treatment for Advanced Non-Squamous Non-Small-Cell Lung Cancer: A Randomized, Double-Blind Study. BioDrugs 2019;33:555-570. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31338773>.

898. Melosky B, Reardon DA, Nixon AB, et al. Bevacizumab biosimilars: scientific justification for extrapolation of indications. Future Oncol 2018;14:2507-2520. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29690784>.

899. Weise M, Kurki P, Wolff-Holz E, et al. Biosimilars: the science of extrapolation. Blood 2014;124:3191-3196. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25298038>.



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

900. Weise M, Bielsky MC, De Smet K, et al. Biosimilars: what clinicians should know. *Blood* 2012;120:5111-5117. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/23093622>.

901. Mok TS, Cheng Y, Zhou X, et al. Improvement in Overall Survival in a Randomized Study That Compared Dacomitinib With Gefitinib in Patients With Advanced Non-Small-Cell Lung Cancer and EGFR-Activating Mutations. *J Clin Oncol* 2018;36:2244-2250. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/29864379>.

902. Wu YL, Cheng Y, Zhou X, et al. Dacomitinib versus gefitinib as first-line treatment for patients with EGFR-mutation-positive non-small-cell lung cancer (ARCHER 1050): a randomised, open-label, phase 3 trial. *Lancet Oncol* 2017;18:1454-1466. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/28958502>.

903. Mok TS, Cheng Y, Zhou X, et al. Updated Overall Survival in a Randomized Study Comparing Dacomitinib with Gefitinib as First-Line Treatment in Patients with Advanced Non-Small-Cell Lung Cancer and EGFR-Activating Mutations. *Drugs* 2021;81:257-266. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/33331989>.

904. Ricciuti B, Chiari R, Chiarini P, et al. Osimertinib (AZD9291) and CNS Response in Two Radiotherapy-Naive Patients with EGFR-Mutant and T790M-Positive Advanced Non-Small Cell Lung Cancer. *Clin Drug Investig* 2016;36:683-686. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/27177916>.

905. Ramalingam SS, Reungwetwattana T, Chewaskulyong B, et al. Osimertinib versus standard-of-care EGFR-TKI as first-line treatment in patients with EGFRm advanced NSCLC: FLAURA [abstract] [abstract]. Presented at the ESMO Congress; Madrid. Abstract LBA2_PR.

906. Ramalingam SS, Yang JC, Lee CK, et al. Osimertinib As First-Line Treatment of EGFR Mutation-Positive Advanced Non-Small-Cell Lung Cancer. *J Clin Oncol* 2018;36:841-849. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/28841389>.

907. Schoenfeld AJ, Arbour KC, Rizvi H, et al. Severe immune-related adverse events are common with sequential PD-(L)1 blockade and osimertinib. *Ann Oncol* 2019;30:839-844. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/30847464>.

908. Oshima Y, Tanimoto T, Yuji K, Tojo A. EGFR-TKI-Associated Interstitial Pneumonitis in Nivolumab-Treated Patients With Non-Small Cell Lung Cancer. *JAMA Oncol* 2018;4:1112-1115. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/29327061>.

909. Yang JCH, Kim SW, Kim DW, et al. Osimertinib in patients with epidermal growth factor receptor mutation-positive non-small-cell lung cancer and leptomeningeal metastases: The BLOOM study. *J Clin Oncol* 2020;38:538-547. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/31809241>.

910. Yang JC-H, Cho BC, Kim D-W, et al. Osimertinib for patients (pts) with leptomeningeal metastases (LM) from EGFR-mutant non-small cell lung cancer (NSCLC): Updated results from the BLOOM study [abstract]. *J Clin Oncol* 2017;35(15):Abstract 2020. Available at:

http://ascopubs.org/doi/abs/10.1200/JCO.2017.35.15_suppl.2020.

911. Merker JD, Oxnard GR, Compton C, et al. Circulating Tumor DNA Analysis in Patients With Cancer: American Society of Clinical Oncology and College of American Pathologists Joint Review. *J Clin Oncol* 2018;36:1631-1641. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/29504847>.

912. Oxnard GR, Thress KS, Alden RS, et al. Association Between Plasma Genotyping and Outcomes of Treatment With Osimertinib (AZD9291) in Advanced Non-Small-Cell Lung Cancer. *J Clin Oncol* 2016;34:3375-3382. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/27354477>.

913. Sacher AG, Paweletz C, Dahlberg SE, et al. Prospective Validation of Rapid Plasma Genotyping for the Detection of EGFR and KRAS Mutations in Advanced Lung Cancer. *JAMA Oncol* 2016;2:1014-1022. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/27055085>.



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

914. Hochmair MJ, Holzer S, Filipits M, et al. EGFR T790M resistance mutation in NSCLC: Real-life data of patients treated with osimertinib [abstract]. J Clin Oncol 2016;34:Abstract e20572. Available at:

915. Reichegger H, Jochum W, Forbs D, et al. Rapid Intracranial Response to Osimertinib in a Patient with Epidermal Growth Factor Receptor T790M-Positive Adenocarcinoma of the Lung. Oncol Res Treat 2016;39:461-463. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/27486808>.

916. Ballard P, Yates JW, Yang Z, et al. Preclinical comparison of osimertinib with other EGFR-TKIs in EGFR-mutant NSCLC brain metastases models, and early evidence of clinical brain metastases activity. Clin Cancer Res 2016;22:5130-5140. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/27435396>.

917. Janjigian YY, Smit EF, Groen HJ, et al. Dual inhibition of EGFR with afatinib and cetuximab in kinase inhibitor-resistant EGFR-mutant lung cancer with and without T790M mutations. Cancer Discov 2014;4:1036-1045. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/25074459>.

918. Pirker R, Pereira JR, Szczesna A, et al. Cetuximab plus chemotherapy in patients with advanced non-small-cell lung cancer (FLEX): an open-label randomised phase III trial. Lancet 2009;373:1525-1531. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/19410716>.

919. Li BT, Shen R, Buonocore D, et al. Ado-trastuzumab emtansine in patients with HER2 mutant lung cancers: Results from a phase II basket trial. J Clin Oncol 2017;35:Abstract 8510. Available at:
http://abstracts.asco.org/199/AbstView_199_193079.html.

920. Iwama E, Zenke Y, Sugawara S, et al. Trastuzumab emtansine for patients with non-small cell lung cancer positive for human epidermal growth factor receptor 2 exon-20 insertion mutations. Eur J Cancer 2022;162:99-106. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/34959152>.

921. Tsurutani J, Iwata H, Krop I, et al. Targeting HER2 with Trastuzumab Deruxtecan: A Dose-Expansion, Phase I Study in Multiple Advanced Solid Tumors. Cancer Discov 2020;10:688-701. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/32213540>.

922. Smit EF, Nakagawa K, Nagasaka M, et al. Trastuzumab deruxtecan (T-DXd; DS-8201) in patients with HER2-mutated metastatic non-small cell lung cancer (NSCLC): Interim results of DESTINY-Lung01. Journal of Clinical Oncology 2020;38:9504-9504. Available at:
https://ascopubs.org/doi/abs/10.1200/JCO.2020.38.15_suppl.9504.

923. Cappuzzo F, Bemis L, Varella-Garcia M. HER2 mutation and response to trastuzumab therapy in non-small-cell lung cancer. N Engl J Med 2006;354:2619-2621. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/16775247>.

924. Mazieres J, Peters S, Lepage B, et al. Lung cancer that harbors an HER2 mutation: epidemiologic characteristics and therapeutic perspectives. J Clin Oncol 2013;31:1997-2003. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/23610105>.

925. Guisier F, Dubos-Arvis C, Vinas F, et al. Efficacy and Safety of Anti-PD-1 Immunotherapy in Patients With Advanced NSCLC With BRAF, HER2, or MET Mutations or RET Translocation: GFPC 01-2018. J Thorac Oncol 2020;15:628-636. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/31945494>.

926. Garon EB, Heist RS, Seto T, et al. CT082 - Capmatinib in METex14-mutated (mut) advanced non-small cell lung cancer (NSCLC): Results from the phase II GEOMETRY mono-1 study, including efficacy in patients (pts) with brain metastases (BM) [abstract]. Presented at the ACCR Annual Meeting 2020 (virtual). Abstract CT082.

927. Schuler MH, Berardi R, Lim W-T, et al. Phase (Ph) I study of the safety and efficacy of the cMET inhibitor capmatinib (INC280) in patients (pts) with advanced cMET+ non-small cell lung cancer (NSCLC). Journal of Clinical Oncology 2016;34:9067-9067. Available at:
https://ascopubs.org/doi/abs/10.1200/JCO.2016.34.15_suppl.9067.



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

928. Le X, Sakai H, Felip E, et al. Tepotinib efficacy and safety in patients with MET exon 14 skipping NSCLC: Outcomes in patient subgroups from the VISION study with relevance for clinical practice. Clin Cancer Res 2022;28:1117-1126. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/34789481>.

929. Doebele RC, Ahn M-J, Siena S, et al. OA02.01: Efficacy and safety of entrectinib in locally advanced or metastatic ROS1-positive non-small cell lung cancer (NSCLC) [abstract]. J Thorac Oncol 2018;13:S321-S322. Available at: [https://www.jto.org/article/S1556-0864\(18\)31197-3/fulltext](https://www.jto.org/article/S1556-0864(18)31197-3/fulltext).

930. Drilon AE, Sima CS, Somwar R, et al. Phase II study of cabozantinib for patients with advanced RET-rearranged lung cancers [abstract]. J Clin Oncol 2015;33:Abstract 8007. Available at: <http://meetinglibrary.asco.org/content/147349-156>.

931. Drilon A, Oxnard G, Wirth L, et al. PL02.08 Registrational Results of LIBRETTO-001: A Phase 1/2 Trial of LOXO-292 in Patients with RET Fusion-Positive Lung Cancers. Journal of Thoracic Oncology 2019;14:S6-S7. Available at: <https://doi.org/10.1016/j.jtho.2019.08.059>.

932. Michels S, Massuti B, Schildhaus HU, et al. Safety and Efficacy of Crizotinib in Patients With Advanced or Metastatic ROS1-Rearranged Lung Cancer (EUCROSS): A European Phase II Clinical Trial. J Thorac Oncol 2019;14:1266-1276. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30978502>.

933. Wu YL, Yang JC, Kim DW, et al. Phase II Study of Crizotinib in East Asian Patients With ROS1-Positive Advanced Non-Small-Cell Lung Cancer. J Clin Oncol 2018;36:1405-1411. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29596029>.

934. Liu D, Offin M, Harnicar S, et al. Entrectinib: an orally available, selective tyrosine kinase inhibitor for the treatment of NTRK, ROS1, and ALK fusion-positive solid tumors. Ther Clin Risk Manag 2018;14:1247-1252. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30050303>.

935. Schneider JL, Muzikansky A, Lin JJ, et al. A phase 2 study of lorlatinib in patients with ROS1-rearranged lung cancer with brain-only progression on crizotinib. JTO Clin Res Rep 2022;3:100347. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/35815322>.

936. Lin JJ, Choudhury NJ, Yoda S, et al. Spectrum of mechanisms of resistance to crizotinib and lorlatinib in ROS1 fusion-positive lung cancer. Clin Cancer Res 2021;27:2899-2909. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/33685866>.

937. Cohen MH, Gootenberg J, Keegan P, Pazdur R. FDA drug approval summary: bevacizumab (Avastin) plus Carboplatin and Paclitaxel as first-line treatment of advanced/metastatic recurrent nonsquamous non-small cell lung cancer. Oncologist 2007;12:713-718. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17602060>.

938. Garon EB, Ciuleanu TE, Arrieta O, et al. Ramucirumab plus docetaxel versus placebo plus docetaxel for second-line treatment of stage IV non-small-cell lung cancer after disease progression on platinum-based therapy (REVEL): a multicentre, double-blind, randomised phase 3 trial. Lancet 2014;384:665-673. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24933332>.

939. Larkins E, Sceapura B, Blumenthal GM, et al. U.S. Food and Drug Administration Approval Summary: Ramucirumab for the Treatment of Metastatic Non-Small Cell Lung Cancer Following Disease Progression On or After Platinum-Based Chemotherapy. Oncologist 2015;20:1320-1325. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26446239>.

940. Choi W, Park SY, Lee Y, et al. The clinical impact of capmatinib in the treatment of advanced non-small cell lung cancer with MET Exon 14 skipping mutation or gene amplification. Cancer Res Treat 2021;53:1024-1032. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/33540494>.

941. Camidge DR, Otterson GA, Clark JW, et al. Crizotinib in patients with MET-amplified NSCLC. J Thorac Oncol 2021;16:1017-1029. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/33676017>.



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

942. Imakita T, Fujita K, Kanai O, et al. Small cell transformation of non-small cell lung cancer under immunotherapy: Case series and literature review. *Thorac Cancer* 2021;12:3062-3067. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/34622569>.

943. Hanna NH, Schneider BJ, Temin S, et al. Therapy for Stage IV Non-Small-Cell Lung Cancer Without Driver Alterations: ASCO and OH (CCO) Joint Guideline Update. *J Clin Oncol* 2020;38:1608-1632. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31990617>.

944. Mok TSK, Wu YL, Kudaba I, et al. Pembrolizumab versus chemotherapy for previously untreated, PD-L1-expressing, locally advanced or metastatic non-small-cell lung cancer (KEYNOTE-042): a randomised, open-label, controlled, phase 3 trial. *Lancet* 2019;393:1819-1830. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30955977>.

945. Rittmeyer A, Barlesi F, Waterkamp D, et al. Atezolizumab versus docetaxel in patients with previously treated non-small-cell lung cancer (OAK): a phase 3, open-label, multicentre randomised controlled trial. *Lancet* 2017;389:255-265. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27979383>.

946. Herbst RS, Baas P, Kim DW, et al. Pembrolizumab versus docetaxel for previously treated, PD-L1-positive, advanced non-small-cell lung cancer (KEYNOTE-010): a randomised controlled trial. *Lancet* 2016;387:1540-1550. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26712084>.

947. Brahmer JR, Lacchetti C, Schneider BJ, et al. Management of immune-related adverse events in patients treated with immune checkpoint inhibitor therapy: American Society of Clinical Oncology Clinical Practice Guideline. *J Clin Oncol* 2018;36:1714-1768. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29442540>.

948. Davies M, Duffield EA. Safety of checkpoint inhibitors for cancer treatment: strategies for patient monitoring and management of immune-mediated adverse events. *Immunotargets Ther* 2017;6:51-71. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28894725>.

949. Cascone T, Weissferdt A, Godoy MCB, et al. Nodal immune flare mimics nodal disease progression following neoadjuvant immune checkpoint inhibitors in non-small cell lung cancer. *Nat Commun* 2021;12:5045. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/34413300>.

950. Cousin S, Toulmonde M, Kind M, et al. Pulmonary sarcoidosis induced by the anti-PD1 monoclonal antibody pembrolizumab. *Ann Oncol* 2016;27:1178-1179. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27091806>.

951. Chiou VL, Burotto M. Pseudoprogression and Immune-Related Response in Solid Tumors. *J Clin Oncol* 2015;33:3541-3543. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26261262>.

952. Socinski MA, Jotte RM, Cappuzzo F, et al. Atezolizumab for First-Line Treatment of Metastatic Nonsquamous NSCLC. *N Engl J Med* 2018;378:2288-2301. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29863955>.

953. Herbst RS, Giaccone G, de Marinis F, et al. Atezolizumab for First-Line Treatment of PD-L1-Selected Patients with NSCLC. *N Engl J Med* 2020;383:1328-1339. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32997907>.

954. Spigel DR, De Marinis F, Giaccone G, et al. IMpower110: Interim OS analysis of a phase III study of atezolizumab (atezo)vs platinum-based chemotherapy (chemo) as 1L treatment (tx) in PD-L1–selected NSCLC [abstract] *Ann Oncol* 2019;30(suppl_5):Abstract 6256. Available at: <https://tinyurl.com/yacn39pe>.

955. Herbst R, De Marinis F, Giaccone G, et al. Clinical Efficacy of Atezolizumab in Biomarker Subgroups by SP142, SP263 and 22C3 PD-L1 Immunohistochemistry Assays and by Blood Tumour Mutational Burden: Results From the IMpower110 Study [abstract]. *European Society For Medical Oncology (ESMO) Immuno-Oncology Congress Geneva, Switzerland; 2019:Abstract 325*. Available at: <https://tinyurl.com/y8wryaku>.



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

956. Reck M, Mok TSK, Nishio M, et al. Atezolizumab plus bevacizumab and chemotherapy in non-small-cell lung cancer (IMpower150): key subgroup analyses of patients with EGFR mutations or baseline liver metastases in a randomised, open-label phase 3 trial. *Lancet Respir Med* 2019;7:387-401. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30922878>.

957. West H, McCleod M, Hussein M, et al. Atezolizumab in combination with carboplatin plus nab-paclitaxel chemotherapy compared with chemotherapy alone as first-line treatment for metastatic non-squamous non-small-cell lung cancer (IMpower130): a multicentre, randomised, open-label, phase 3 trial. *Lancet Oncol* 2019;20:924-937. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31122901>.

958. Barlesi F, Park K, Ciardiello F. Primary analysis from OAK, a randomized phase III study comparing atezolizumab with docetaxel in 2L/3L NSCLC [abstract] [abstract]. Presented at the 2016 Annual Meeting European Society for Medical Oncology (ESMO) Copenhagen, Denmark. Abstract LBA44.

959. Gogishvili M, Melkadze T, Makharadze T, et al. Cemiplimab plus chemotherapy versus chemotherapy alone in non-small cell lung cancer: a randomized, controlled, double-blind phase 3 trial. *Nat Med* 2022;28:2374-2380. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/36008722>.

960. Wei J, van der Wekken AJ, Saber A, et al. Mutations in EMT-Related Genes in ALK Positive Crizotinib Resistant Non-Small Cell Lung Cancers. *Cancers (Basel)* 2018;10. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29300322>.

961. Paz-Ares LG, Ramalingam SS, Ciuleanu TE, et al. First-line nivolumab plus ipilimumab in advanced NSCLC: 4-year outcomes from the randomized, open-label, phase 3 CheckMate 227 part 1 trial. *J Thorac Oncol* 2022;17:289-308. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/34648948>.

962. Reck M, Ciuleanu T-E, Dols MC, et al. Nivolumab (NIVO) + ipilimumab (IPI) + 2 cycles of platinum-doublet chemotherapy (chemo) vs

4 cycles chemo as first-line (1L) treatment (tx) for stage IV/recurrent non-small cell lung cancer (NSCLC): CheckMate 9LA. *Journal of Clinical Oncology* 2020;38:9501-9501. Available at: https://ascopubs.org/doi/abs/10.1200/JCO.2020.38.15_suppl.9501.

963. Reck M, Ciuleanu TE, Cobo M, et al. First-line nivolumab plus ipilimumab with two cycles of chemotherapy versus chemotherapy alone (four cycles) in advanced non-small-cell lung cancer: CheckMate 9LA 2-year update. *ESMO Open* 2021;6:100273. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/34607285>.

964. Paz-Ares L, Ciuleanu TE, Cobo M, et al. First-line nivolumab plus ipilimumab combined with two cycles of chemotherapy in patients with non-small-cell lung cancer (CheckMate 9LA): an international, randomised, open-label, phase 3 trial. *Lancet Oncol* 2021;22:198-211. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/33476593>.

965. Horn L, Spigel DR, Vokes EE, et al. Nivolumab versus docetaxel in previously treated patients with advanced non-small-cell lung cancer: Two-year outcomes from two randomized, open-label, phase III trials (CheckMate 017 and CheckMate 057). *J Clin Oncol* 2017;35:3924-3933. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29023213>.

966. Kazandjian D, Suzman DL, Blumenthal G, et al. FDA Approval Summary: Nivolumab for the Treatment of Metastatic Non-Small Cell Lung Cancer With Progression On or After Platinum-Based Chemotherapy. *Oncologist* 2016;21:634-642. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26984449>.

967. Melosky B, Chu Q, Juergens R, et al. Pointed Progress in Second-Line Advanced Non-Small-Cell Lung Cancer: The Rapidly Evolving Field of Checkpoint Inhibition. *J Clin Oncol* 2016;34:1676-1688. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26884577>.

968. Nishino M, Giobbie-Hurder A, Hatabu H, et al. Incidence of Programmed Cell Death 1 Inhibitor-Related Pneumonitis in Patients With Advanced Cancer: A Systematic Review and Meta-analysis. *JAMA Oncol* 2016;2:1607-1616. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27540850>.



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

969. Naidoo J, Wang X, Woo KM, et al. Pneumonitis in Patients Treated With Anti-Programmed Death-1/Programmed Death Ligand 1 Therapy. *J Clin Oncol* 2017;35:709-717. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27646942>.

970. Sgambato A, Casaluce F, Sacco PC, et al. Anti PD-1 and PDL-1 Immunotherapy in the Treatment of Advanced Non- Small Cell Lung Cancer (NSCLC): A Review on Toxicity Profile and its Management. *Curr Drug Saf* 2016;11:62-68. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26412670>.

971. Gettinger SN, Horn L, Gandhi L, et al. Overall Survival and Long-Term Safety of Nivolumab (Anti-Programmed Death 1 Antibody, BMS-936558, ONO-4538) in Patients With Previously Treated Advanced Non-Small-Cell Lung Cancer. *J Clin Oncol* 2015;33:2004-2012. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25897158>.

972. Rizvi NA, Mazieres J, Planchard D, et al. Activity and safety of nivolumab, an anti-PD-1 immune checkpoint inhibitor, for patients with advanced, refractory squamous non-small-cell lung cancer (CheckMate 063): a phase 2, single-arm trial. *Lancet Oncol* 2015;16:257-265. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25704439>.

973. Postow MA, Callahan MK, Wolchok JD. Immune Checkpoint Blockade in Cancer Therapy. *J Clin Oncol* 2015;33:1974-1982. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25605845>.

974. Chapman PB, D'Angelo SP, Wolchok JD. Rapid eradication of a bulky melanoma mass with one dose of immunotherapy. *N Engl J Med* 2015;372:2073-2074. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25891305>.

975. Khoja L, Butler MO, Kang SP, et al. Pembrolizumab. *J Immunother Cancer* 2015;3:36. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26288737>.

976. Pai-Scherf L, Blumenthal GM, Li H, et al. FDA Approval Summary: Pembrolizumab for Treatment of Metastatic Non-Small Cell Lung Cancer:

First-Line Therapy and Beyond. *Oncologist* 2017;22:1392-1399. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28835513>.

977. Garassino M, Rodriguez-Abreu D, Gadgeel S, et al. OA04.06 Evaluation of TMB IN KEYNOTE-189: pembrolizumab plus chemotherapy vs placebo plus chemotherapy for nonsquamous NSCLC [abstract]. 2019 World Conference on Lung Cancer (WCLC). Barcelona, Spain: International Association for the Study of Lung Cancer (IASLC) 2019:Abstract: OA04.06. Available at: <https://www.iaslc.org/About-IASLC/News-Detail/keynote-189-tumor-mutational-burden-not-significantly-associated-with-efficacy-of-pembrolizumab>.

978. Paz-Ares LG, Luft A, Tafreshi A, et al. Phase 3 study of carboplatin-paclitaxel/nab-paclitaxel (Chemo) with or without pembrolizumab for patients with metastatic squamous non-small cell lung cancer [abstract]. *J Clin Oncol* 2018;36:Abstract 105. Available at: http://abstracts.asco.org/214/AbstView_214_228023.html.

979. Borghaei H, Langer CJ, Paz-Ares L, et al. Pembrolizumab plus chemotherapy versus chemotherapy alone in patients with advanced non-small cell lung cancer without tumor PD-L1 expression: A pooled analysis of 3 randomized controlled trials. *Cancer* 2020;126:4867-4877. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32914866>.

980. Paz-Ares L, Luft A, Vicente D, et al. Pembrolizumab plus Chemotherapy for Squamous Non-Small-Cell Lung Cancer. *N Engl J Med* 2018;379:2040-2051. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30280635>.

981. Garon EB, Rizvi NA, Hui R, et al. Pembrolizumab for the treatment of non-small-cell lung cancer. *N Engl J Med* 2015;372:2018-2028. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25891174>.

982. Sul J, Blumenthal GM, Jiang X, et al. FDA Approval Summary: Pembrolizumab for the Treatment of Patients With Metastatic Non-Small Cell Lung Cancer Whose Tumors Express Programmed Death-Ligand 1. *Oncologist* 2016;21:643-650. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27026676>.



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

983. Rizvi NA, Cho BC, Reinmuth N, et al. Durvalumab with or without tremelimumab vs standard chemotherapy in first-line treatment of metastatic non-small cell lung cancer: the MYSTIC phase 3 randomized clinical trial. *JAMA Oncol* 2020;6:661-674. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32271377>.

984. Johnson ML, Cho BC, Luft A, et al. Durvalumab with or without tremelimumab in combination with chemotherapy as first-line therapy for metastatic non-small-cell lung cancer: The phase III POSEIDON study. *J Clin Oncol* 2023;41:1213-1227. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/36327426>.

985. Peters S, Cho BC, Luft A, et al. OA15.04 Association between KRAS/STK11/KEAP1 mutations and outcomes in POSEIDON: Durvalumab ± tremelimumab + chemotherapy in mNSCLC [abstract]. *J Thorac Oncol* 2022;17:S39-S41. Available at: <https://www.sciencedirect.com/science/article/pii/S1556086422004142>.

986. Gridelli C, de Marinis F, Di Maio M, et al. Maintenance treatment of advanced non-small-cell lung cancer: results of an international expert panel meeting of the Italian association of thoracic oncology. *Lung Cancer* 2012;76:269-279. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22266040>.

987. Hashemi-Sadraei N, Pennell NA. Advanced non-small cell lung cancer (NSCLC): maintenance therapy for all? *Curr Treat Options Oncol* 2012;13:478-490. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22972369>.

988. Paz-Ares L, de Marinis F, Dediu M, et al. Maintenance therapy with pemetrexed plus best supportive care versus placebo plus best supportive care after induction therapy with pemetrexed plus cisplatin for advanced non-squamous non-small-cell lung cancer (PARAMOUNT): a double-blind, phase 3, randomised controlled trial. *Lancet Oncol* 2012;13:247-255. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22341744>.

989. Paz-Ares LG, de Marinis F, Dediu M, et al. PARAMOUNT: Final overall survival results of the phase III study of maintenance pemetrexed versus placebo immediately after induction treatment with pemetrexed

plus cisplatin for advanced nonsquamous non-small-cell lung cancer. *J Clin Oncol* 2013;31:2895-2902. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23835707>.

990. Patel JD, Hensing TA, Rademaker A, et al. Phase II study of pemetrexed and carboplatin plus bevacizumab with maintenance pemetrexed and bevacizumab as first-line therapy for nonsquamous non-small-cell lung cancer. *J Clin Oncol* 2009;27:3284-3289. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19433684>.

991. Barlesi F, Scherpereel A, Gorbunova V, et al. Maintenance bevacizumab-pemetrexed after first-line cisplatin-pemetrexed-bevacizumab for advanced nonsquamous non-small-cell lung cancer: updated survival analysis of the AVAPERL (MO22089) randomized phase III trial. *Ann Oncol* 2014;25:1044-1052. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24585722>.

992. Barlesi F, Scherpereel A, Rittmeyer A, et al. Randomized phase III trial of maintenance bevacizumab with or without pemetrexed after first-line induction with bevacizumab, cisplatin, and pemetrexed in advanced nonsquamous non-small-cell lung cancer: AVAPERL (MO22089). *J Clin Oncol* 2013;31:3004-3011. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23835708>.

993. Nadler E, Yu E, Ravelo A, et al. Bevacizumab treatment to progression after chemotherapy: outcomes from a U.S. community practice network. *Oncologist* 2011;16:486-496. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21441299>.

994. Perol M, Chouaid C, Milleron BJ, et al. Maintenance with either gemcitabine or erlotinib versus observation with predefined second-line treatment after cisplatin-gemcitabine induction chemotherapy in advanced NSCLC: IFCT-GFPC 0502 phase III study [abstract]. *J Clin Oncol* 2010;28(Suppl 15):Abstract 7507. Available at: http://meeting.ascopubs.org/cgi/content/abstract/28/15_suppl/7507.

995. Brodowicz T, Krzakowski M, Zwitter M, et al. Cisplatin and gemcitabine first-line chemotherapy followed by maintenance gemcitabine or best supportive care in advanced non-small cell lung cancer: a phase III



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

trial. Lung Cancer 2006;52:155-163. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/16569462>.

996. Gerber DE, Schiller JH. Maintenance chemotherapy for advanced non-small-cell lung cancer: new life for an old idea. J Clin Oncol 2013;31:1009-1020. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/23401441>.

997. Cappuzzo F, Ciuleanu T, Stelmakh L, et al. Erlotinib as maintenance treatment in advanced non-small-cell lung cancer: a multicentre, randomised, placebo-controlled phase 3 study. Lancet Oncol 2010;11:521-529. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/20493771>.

998. Ciuleanu T, Brodowicz T, Zielinski C, et al. Maintenance pemetrexed plus best supportive care versus placebo plus best supportive care for non-small-cell lung cancer: a randomised, double-blind, phase 3 study. Lancet 2009;374:1432-1440. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/19767093>.

999. Cohen MH, Cortazar P, Justice R, Pazdur R. Approval summary: pemetrexed maintenance therapy of advanced/metastatic nonsquamous, non-small cell lung cancer (NSCLC). Oncologist 2010;15:1352-1358. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21148615>.

1000. Cicen S, Geater SL, Petrov P, et al. Maintenance erlotinib versus erlotinib at disease progression in patients with advanced non-small-cell lung cancer who have not progressed following platinum-based chemotherapy (IUNO study). Lung Cancer 2016;102:30-37. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/27987585>.

1001. Rittmeyer A. Quality of Life in Patients with NSCLC Receiving Maintenance Therapy. Cancers (Basel) 2015;7:950-962. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/26035509>.

1002. Fidias PM, Dakhil SR, Lyss AP, et al. Phase III study of immediate compared with delayed docetaxel after front-line therapy with gemcitabine plus carboplatin in advanced non-small-cell lung cancer. J Clin Oncol

2009;27:591-598. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/19075278>.

1003. Coate LE, Shepherd FA. Maintenance therapy in advanced non-small cell lung cancer: evolution, tolerability and outcomes. Ther Adv Med Oncol 2011;3:139-157. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/21904577>.

1004. Rigotti NA, Regan S, Levy DE, et al. Sustained care intervention and postdischarge smoking cessation among hospitalized adults: a randomized clinical trial. JAMA 2014;312:719-728. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/25138333>.

1005. Stead LF, Hartmann-Boyce J, Perera R, Lancaster T. Telephone counselling for smoking cessation. Cochrane Database Syst Rev 2013;8:CD002850. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/23934971>.

1006. Stead LF, Lancaster T. Combined pharmacotherapy and behavioural interventions for smoking cessation. Cochrane Database Syst Rev 2012;10:CD008286. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/23076944>.

1007. Patterson GA, Ginsberg RJ, Poon PY, et al. A prospective evaluation of magnetic resonance imaging, computed tomography, and mediastinoscopy in the preoperative assessment of mediastinal node status in bronchogenic carcinoma. J Thorac Cardiovasc Surg 1987;94:679-684. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/3669696>.

1008. Gonzalez-Stawinski GV, Lemaire A, Merchant F, et al. A comparative analysis of positron emission tomography and mediastinoscopy in staging non-small cell lung cancer. J Thorac Cardiovasc Surg 2003;126:1900-1905. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/14688703>.

1009. Tournoy KG, Maddens S, Gosselin R, et al. Integrated FDG-PET/CT does not make invasive staging of the intrathoracic lymph nodes in non-small cell lung cancer redundant: a prospective study. Thorax



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

2007;62:696-701. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/17687098>.

1010. Meyers BF, Haddad F, Siegel BA, et al. Cost-effectiveness of routine mediastinoscopy in computed tomography- and positron emission tomography-screened patients with stage I lung cancer. J Thorac Cardiovasc Surg 2006;131:822-829; discussion 822-829. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/16580440>.

1011. Dillemans B, Deneffe G, Verschakelen J, Decramer M. Value of computed tomography and mediastinoscopy in preoperative evaluation of mediastinal nodes in non-small cell lung cancer. A study of 569 patients. Eur J Cardiothorac Surg 1994;8:37-42. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/8136168>.

1012. Arita T, Kuramitsu T, Kawamura M, et al. Bronchogenic carcinoma: incidence of metastases to normal sized lymph nodes. Thorax 1995;50:1267-1269. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/8553299>.

1013. McCloud TC, Bourgouin PM, Greenberg RW, et al. Bronchogenic carcinoma: analysis of staging in the mediastinum with CT by correlative lymph node mapping and sampling. Radiology 1992;182:319-323. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/1732943>.

1014. Seely JM, Mayo JR, Miller RR, Muller NL. T1 lung cancer: prevalence of mediastinal nodal metastases and diagnostic accuracy of CT. Radiology 1993;186:129-132. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/8416552>.

1015. Kerr KM, Lamb D, Wathen CG, et al. Pathological assessment of mediastinal lymph nodes in lung cancer: implications for non-invasive mediastinal staging. Thorax 1992;47:337-341. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/1609375>.

1016. Chin R, Jr., Ward R, Keyes JW, et al. Mediastinal staging of non-small-cell lung cancer with positron emission tomography. Am J Respir Crit Care Med 1995;152:2090-2096. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/8520780>.

1017. Kernstine KH, Stanford W, Mullan BF, et al. PET, CT, and MRI with Combidex for mediastinal staging in non-small cell lung carcinoma. Ann Thorac Surg 1999;68:1022-1028. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/10510001>.

1018. De Leyn P, Stroobants S, De Wever W, et al. Prospective comparative study of integrated positron emission tomography-computed tomography scan compared with remediastinoscopy in the assessment of residual mediastinal lymph node disease after induction chemotherapy for mediastinoscopy-proven stage IIIA-N2 Non-small-cell lung cancer: a Leuven Lung Cancer Group Study. J Clin Oncol 2006;24:3333-3339. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/16849747>.

1019. Cerfolio RJ, Bryant AS, Ojha B. Restaging patients with N2 (stage IIIa) non-small cell lung cancer after neoadjuvant chemoradiotherapy: a prospective study. J Thorac Cardiovasc Surg 2006;131:1229-1235. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/16733150>.

1020. Maziak DE, Darling GE, Inculet RI, et al. Positron emission tomography in staging early lung cancer: a randomized trial. Ann Intern Med 2009;151:221-228, W-248. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/19581636>.

1021. Pieterman RM, van Putten JW, Meuzelaar JJ, et al. Preoperative staging of non-small-cell lung cancer with positron-emission tomography. N Engl J Med 2000;343:254-261. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/10911007>.

1022. Manente P, Vicario G, Piazza F, et al. Does PET/CT modify the therapeutic approach in medical oncology [abstract]? . J Clin Oncol 2008;26(Suppl 15):Abstract 17525. Available at:

http://meeting.ascopubs.org/cgi/content/abstract/26/15_suppl/17525.

1023. Fischer B, Lassen U, Mortensen J, et al. Preoperative staging of lung cancer with combined PET-CT. N Engl J Med 2009;361:32-39. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/19571281>.

1024. De Wever W, Stroobants S, Coolen J, Verschakelen JA. Integrated PET/CT in the staging of nonsmall cell lung cancer: technical aspects and



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

clinical integration. Eur Respir J 2009;33:201-212. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19118231>.

1025. Darling GE, Maziak DE, Inculet RI, et al. Positron emission tomography-computed tomography compared with invasive mediastinal staging in non-small cell lung cancer: results of mediastinal staging in the early lung positron emission tomography trial. J Thorac Oncol 2011;6:1367-1372. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21587082>.

1026. Yasufuku K, Pierre A, Darling G, et al. A prospective controlled trial of endobronchial ultrasound-guided transbronchial needle aspiration compared with mediastinoscopy for mediastinal lymph node staging of lung cancer. J Thorac Cardiovasc Surg 2011;142:1393-1400 e1391. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21963329>.

1027. Annema JT, van Meerbeeck JP, Rintoul RC, et al. Mediastinoscopy vs endosonography for mediastinal nodal staging of lung cancer: a randomized trial. JAMA 2010;304:2245-2252. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21098770>.

1028. Tournoy KG, Keller SM, Annema JT. Mediastinal staging of lung cancer: novel concepts. Lancet Oncol 2012;13:e221-229. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22554550>.

1029. Vilman P, Krasnik M, Larsen SS, et al. Transesophageal endoscopic ultrasound-guided fine-needle aspiration (EUS-FNA) and endobronchial ultrasound-guided transbronchial needle aspiration (EBUS-TBNA) biopsy: a combined approach in the evaluation of mediastinal lesions. Endoscopy 2005;37:833-839. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/16116534>.

1030. Yasufuku K, Nakajima T, Motoori K, et al. Comparison of endobronchial ultrasound, positron emission tomography, and CT for lymph node staging of lung cancer. Chest 2006;130:710-718. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/16963667>.

1031. Ernst A, Eberhardt R, Krasnik M, Herth FJ. Efficacy of endobronchial ultrasound-guided transbronchial needle aspiration of hilar

lymph nodes for diagnosing and staging cancer. J Thorac Oncol 2009;4:947-950. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19590457>.

1032. Rintoul RC, Tournoy KG, El Daly H, et al. EBUS-TBNA for the clarification of PET positive intra-thoracic lymph nodes-an international multi-centre experience. J Thorac Oncol 2009;4:44-48. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19096305>.

1033. Defranchi SA, Edell ES, Daniels CE, et al. Mediastinoscopy in patients with lung cancer and negative endobronchial ultrasound guided needle aspiration. Ann Thorac Surg 2010;90:1753-1757. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21095301>.

1034. Medford AR, Bennett JA, Free CM, Agrawal S. Mediastinal staging procedures in lung cancer: EBUS, TBNA and mediastinoscopy. Curr Opin Pulm Med 2009;15:334-342. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19395972>.

1035. Mayr NA, Hussey DH, Yuh WT. Cost-effectiveness of high-contrast-dose MR screening of asymptomatic brain metastasis. AJNR Am J Neuroradiol 1995;16:215-217. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/7755752>.

1036. Videtic GM, Chang JY, Chetty IJ, et al. ACR appropriateness Criteria(R) early-stage non-small-cell lung cancer. Am J Clin Oncol 2014;37:201-207. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25180631>.

1037. Barnes JB, Johnson SB, Dahiya RS, et al. Concomitant weekly cisplatin and thoracic radiotherapy for Pancoast tumors of the lung: pilot experience of the San Antonio Cancer Institute. Am J Clin Oncol 2002;25:90-92. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/11823705>.

1038. Rusch VW, Giroux DJ, Kraut MJ, et al. Induction chemoradiation and surgical resection for non-small cell lung carcinomas of the superior sulcus: Initial results of Southwest Oncology Group Trial 9416 (Intergroup



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

Trial 0160). J Thorac Cardiovasc Surg 2001;121:472-483. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/11241082>.

1039. Pourel N, Santelmo N, Naafa N, et al. Concurrent cisplatin/etoposide plus 3D-conformal radiotherapy followed by surgery for stage IIB (superior sulcus T3N0)/III non-small cell lung cancer yields a high rate of pathological complete response. Eur J Cardiothorac Surg 2008;33:829-836. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/18367406>.

1040. Gandara DR, Chansky K, Albain KS, et al. Consolidation docetaxel after concurrent chemoradiotherapy in stage IIIB non-small-cell lung cancer: phase II Southwest Oncology Group Study S9504. J Clin Oncol 2003;21:2004-2010. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/12743155>.

1041. Nakagawa T, Okumura N, Miyoshi K, et al. Prognostic factors in patients with ipsilateral pulmonary metastasis from non-small cell lung cancer. Eur J Cardiothorac Surg 2005;28:635-639. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/16126398>.

1042. Lee JG, Lee CY, Kim DJ, et al. Non-small cell lung cancer with ipsilateral pulmonary metastases: prognosis analysis and staging assessment. Eur J Cardiothorac Surg 2008;33:480-484. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/18249000>.

1043. Bhaskarla A, Tang PC, Mashtare T, et al. Analysis of second primary lung cancers in the SEER database. J Surg Res 2010;162:1-6. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20400118>.

1044. Aziz TM, Saad RA, Glasser J, et al. The management of second primary lung cancers. A single centre experience in 15 years. Eur J Cardiothorac Surg 2002;21:527-533. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/11888775>.

1045. Adebajo SA, Moritz DM, Danby CA. The results of modern surgical therapy for multiple primary lung cancers. Chest 1997;112:693-701. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/9315801>.

1046. Nakata M, Sawada S, Yamashita M, et al. Surgical treatments for multiple primary adenocarcinoma of the lung. Ann Thorac Surg 2004;78:1194-1199. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/15464469>.

1047. Ginsberg MS, Griff SK, Go BD, et al. Pulmonary nodules resected at video-assisted thoracoscopic surgery: etiology in 426 patients. Radiology 1999;213:277-282. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/10540672>.

1048. Allen MS. Multiple benign lung tumors. Semin Thorac Cardiovasc Surg 2003;15:310-314. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/12973710>.

1049. Asamura H. Multiple primary cancers or multiple metastases, that is the question. J Thorac Oncol 2010;5:930-931. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20581574>.

1050. Girard N, Deshpande C, Azzoli CG, et al. Use of epidermal growth factor receptor/Kirsten rat sarcoma 2 viral oncogene homolog mutation testing to define clonal relationships among multiple lung adenocarcinomas: comparison with clinical guidelines. Chest 2010;137:46-52. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19376842>.

1051. Han HS, Eom DW, Kim JH, et al. EGFR mutation status in primary lung adenocarcinomas and corresponding metastatic lesions: discordance in pleural metastases. Clin Lung Cancer 2011;12:380-386. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21729655>.

1052. Martini N, Melamed MR. Multiple primary lung cancers. J Thorac Cardiovasc Surg 1975;70:606-612. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/170482>.

1053. Patel SB, Kadi W, Walts AE, et al. Next-Generation Sequencing: A Novel Approach to Distinguish Multifocal Primary Lung Adenocarcinomas from Intrapulmonary Metastases. J Mol Diagn 2017;19:870-880. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28866070>.



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

1054. Zheng R, Shen Q, Mardekian S, et al. Molecular profiling of key driver genes improves staging accuracy in multifocal non-small cell lung cancer. *J Thorac Cardiovasc Surg* 2020;160:e71-e79. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32007245>.

1055. Liu C, Liu C, Zou X, et al. Next-generation sequencing facilitates differentiating between multiple primary lung cancer and intrapulmonary metastasis: a case series. *Diagn Pathol* 2021;16:21. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/33706781>.

1056. Chang YL, Wu CT, Lee YC. Surgical treatment of synchronous multiple primary lung cancers: experience of 92 patients. *J Thorac Cardiovasc Surg* 2007;134:630-637. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17723810>.

1057. Tanvetyanon T, Robinson L, Sommers KE, et al. Relationship between tumor size and survival among patients with resection of multiple synchronous lung cancers. *J Thorac Oncol* 2010;5:1018-1024. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20453687>.

1058. Rea F, Zuin A, Callegaro D, et al. Surgical results for multiple primary lung cancers. *Eur J Cardiothorac Surg* 2001;20:489-495. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/11509268>.

1059. Gibbs IC, Loo BW, Jr. CyberKnife stereotactic ablative radiotherapy for lung tumors. *Technol Cancer Res Treat* 2010;9:589-596. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21070081>.

1060. Godoy MC, Naidich DP. Subsolid pulmonary nodules and the spectrum of peripheral adenocarcinomas of the lung: recommended interim guidelines for assessment and management. *Radiology* 2009;253:606-622. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19952025>.

1061. Pearson FG, DeLarue NC, Ilves R, et al. Significance of positive superior mediastinal nodes identified at mediastinoscopy in patients with resectable cancer of the lung. *J Thorac Cardiovasc Surg* 1982;83:1-11. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/7054602>.

1062. Rice TW. Thoracoscopy in the staging of thoracic malignancies. In: Kaiser LR, Daniel TM, eds, eds. *Thoracoscopic Surgery*. Philadelphia: Lippincott Williams & Wilkins; 1993:153-162.

1063. Gandara DR, Chansky K, Albain KS, et al. Long-term survival with concurrent chemoradiation therapy followed by consolidation docetaxel in stage IIIB non-small-cell lung cancer: a phase II Southwest Oncology Group Study (S9504). *Clin Lung Cancer* 2006;8:116-121. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17026812>.

1064. Mina LA, Neubauer MA, Ansari RH, et al. Phase III trial of cisplatin (P) plus etoposide (E) plus concurrent chest radiation (XRT) with or without consolidation docetaxel (D) in patients (pts) with inoperable stage III non-small cell lung cancer (NSCLC): HOG LUN 01-24/USO-023—Updated results [abstract]. *J Clin Oncol* 2008;26 (Suppl 15):Abstract 7519. Available at: http://meeting.ascopubs.org/cgi/content/abstract/26/15_suppl/7519.

1065. Hanna N, Neubauer M, Yiannoutsos C, et al. Phase III study of cisplatin, etoposide, and concurrent chest radiation with or without consolidation docetaxel in patients with inoperable stage III non-small-cell lung cancer: the Hoosier Oncology Group and U.S. Oncology. *J Clin Oncol* 2008;26:5755-5760. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19001323>.

1066. Hanna NH, Neubauer M, Ansari R, et al. Phase III trial of cisplatin (P) plus etoposide (E) plus concurrent chest radiation (XRT) with or without consolidation docetaxel (D) in patients (pts) with inoperable stage III non-small cell lung cancer (NSCLC): HOG LUN 01-24/USO-023 [abstract]. *J Clin Oncol* 2007;25 (Suppl 18):Abstract 7512. Available at: http://meeting.ascopubs.org/cgi/content/abstract/25/18_suppl/7512.

1067. Decker DA, Dines DE, Payne WS, et al. The significance of a cytologically negative pleural effusion in bronchogenic carcinoma. *Chest* 1978;74:640-642. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/216532>.

1068. Feller-Kopman DJ, Reddy CB, DeCamp MM, et al. Management of Malignant Pleural Effusions. An Official ATS/STS/STR Clinical Practice



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

Guideline. Am J Respir Crit Care Med 2018;198:839-849. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30272503>.

1069. Demmy TL, Gu L, Burkhalter JE, et al. Optimal management of malignant pleural effusions (results of CALGB 30102). J Natl Compr Canc Netw 2012;10:975-982. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22878823>.

1070. de Vin T, Engels B, Gevaert T, et al. Stereotactic radiotherapy for oligometastatic cancer: a prognostic model for survival. Ann Oncol 2014;25:467-471. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24355488>.

1071. Simone CB, 2nd, Burri SH, Heinzerling JH. Novel radiotherapy approaches for lung cancer: combining radiation therapy with targeted and immunotherapies. Transl Lung Cancer Res 2015;4:545-552. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26629423>.

1072. Campo M, Al-Halabi H, Khandekar M, et al. Integration of Stereotactic Body Radiation Therapy With Tyrosine Kinase Inhibitors in Stage IV Oncogene-Driven Lung Cancer. Oncologist 2016;21:964-973. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27354669>.

1073. Belani CP, Ramalingam S, Perry MC, et al. Randomized, phase III study of weekly paclitaxel in combination with carboplatin versus standard every-3-weeks administration of carboplatin and paclitaxel for patients with previously untreated advanced non-small-cell lung cancer. J Clin Oncol 2008;26:468-473. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/18202422>.

1074. Rosell R, Gomez-Codina J, Camps C, et al. A randomized trial comparing preoperative chemotherapy plus surgery with surgery alone in patients with non-small-cell lung cancer. N Engl J Med 1994;330:153-158. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/8043059>.

1075. Pisters K, Vallieres E, Bunn P, et al. S9900: A phase III trial of surgery alone or surgery plus preoperative (preop) paclitaxel/carboplatin (PC) chemotherapy in early stage non-small cell lung cancer (NSCLC): Preliminary results [abstract]. J Clin Oncol 2005;23 (Suppl 16):Abstract

LBA7012. Available at:

http://meeting.ascopubs.org/cgi/content/abstract/23/16_suppl/LBA7012.

1076. Pisters K, Vallieres E, Bunn PA, Jr., et al. S9900: Surgery alone or surgery plus induction (ind) paclitaxel/carboplatin (PC) chemotherapy in early stage non-small cell lung cancer (NSCLC): Follow-up on a phase III trial [abstract]. J Clin Oncol 2007;25 (Suppl 18):Abstract 7520. Available at: http://meeting.ascopubs.org/cgi/content/abstract/25/18_suppl/7520.

1077. Burkes RL, Ginsberg RJ, Shepherd FA, et al. Induction chemotherapy with mitomycin, vindesine, and cisplatin for stage III unresectable non-small-cell lung cancer: results of the Toronto Phase II Trial. J Clin Oncol 1992;10:580-586. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/1312587>.

1078. Bonomi P, Faber L. Neoadjuvant chemoradiation therapy in non-small cell lung cancer: The Rush University experience. Lung Cancer 1993;9:383-390. Available at:

1079. Rusch VW, Albain KS, Crowley JJ, et al. Surgical resection of stage IIIA and stage IIIB non-small-cell lung cancer after concurrent induction chemoradiotherapy. A Southwest Oncology Group trial. J Thorac Cardiovasc Surg 1993;105:97-104; discussion 104-106. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/8380477>.

1080. Park SY, Lee JG, Kim J, et al. Efficacy of platinum-based adjuvant chemotherapy in T2aN0 stage IB non-small cell lung cancer. J Cardiothorac Surg 2013;8:151. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23759129>.

1081. Arriagada R, Auperin A, Burdett S, et al. Adjuvant chemotherapy, with or without postoperative radiotherapy, in operable non-small-cell lung cancer: two meta-analyses of individual patient data. Lancet 2010;375:1267-1277. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20338627>.

1082. Postoperative radiotherapy in non-small-cell lung cancer: systematic review and meta-analysis of individual patient data from nine randomised controlled trials. PORT Meta-analysis Trialists Group. Lancet



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

1998;352:257-263. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/9690404>.

1083. Robinson CG, Patel AP, Bradley JD, et al. Postoperative radiotherapy for pathologic N2 non-small-cell lung cancer treated with adjuvant chemotherapy: a review of the National Cancer Data Base. *J Clin Oncol* 2015;33:870-876. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/25667283>.

1084. Patel SH, Ma Y, Wernicke AG, et al. Evidence supporting contemporary post-operative radiation therapy (PORT) using linear accelerators in N2 lung cancer. *Lung Cancer* 2014;84:156-160. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24674156>.

1085. Choy H, Schwartzberg LS, Dakhil SR, et al. Phase 2 study of pemetrexed plus carboplatin, or pemetrexed plus cisplatin with concurrent radiation therapy followed by pemetrexed consolidation in patients with favorable-prognosis inoperable stage IIIA/B non-small-cell lung cancer. *J Thorac Oncol* 2013;8:1308-1316. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/23981966>.

1086. Garrido P, Engel-Riedel W, Serke M, et al. Final results from a Phase II study of pemetrexed and cisplatin with concurrent thoracic radiation after Pem-Cis induction in patients with unresectable locally advanced non-squamous non-small cell lung cancer (NSCLC). *Lung Cancer* 2015;88:160-166. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/25758556>.

1087. Schneider BJ, Ismaila N, Aerts J, et al. Lung cancer surveillance after definitive curative-intent therapy: ASCO Guideline. *J Clin Oncol* 2020;38:753-766. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/31829901>.

1088. Crabtree TD, Puri V, Chen SB, et al. Does the method of radiologic surveillance affect survival after resection of stage I non-small cell lung cancer? *J Thorac Cardiovasc Surg* 2015;149:45-52, 53 e41-43. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25218540>.

1089. Erb CT, Su KW, Soulos PR, et al. Surveillance Practice Patterns after Curative Intent Therapy for Stage I Non-Small-Cell Lung Cancer in the Medicare Population. *Lung Cancer* 2016;99:200-207. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/27565940>.

1090. Colt HG, Murgu SD, Korst RJ, et al. Follow-up and surveillance of the patient with lung cancer after curative-intent therapy: Diagnosis and management of lung cancer, 3rd ed: American College of Chest Physicians evidence-based clinical practice guidelines. *Chest* 2013;143:e437S-e454S. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/23649451>.

1091. Lou F, Huang J, Sima CS, et al. Patterns of recurrence and second primary lung cancer in early-stage lung cancer survivors followed with routine computed tomography surveillance. *J Thorac Cardiovasc Surg* 2013;145:75-81; discussion 81-72. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/23127371>.

1092. Srikantharajah D, Ghuman A, Nagendran M, Maruthappu M. Is computed tomography follow-up of patients after lobectomy for non-small cell lung cancer of benefit in terms of survival? *Interact Cardiovasc Thorac Surg* 2012;15:893-898. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/22859511>.

1093. Hanna WC, Paul NS, Darling GE, et al. Minimal-dose computed tomography is superior to chest x-ray for the follow-up and treatment of patients with resected lung cancer. *J Thorac Cardiovasc Surg* 2014;147:30-33. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/24139896>.

1094. Calman L, Beaver K, Hind D, et al. Survival benefits from follow-up of patients with lung cancer: a systematic review and meta-analysis. *J Thorac Oncol* 2011;6:1993-2004. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/21892108>.

1095. Dane B, Grechushkin V, Plank A, et al. PET/CT vs. non-contrast CT alone for surveillance 1-year post lobectomy for stage I non-small-cell lung cancer. *Am J Nucl Med Mol Imaging* 2013;3:408-416. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/24116349>.



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

1096. Henschke CI, McCauley DI, Yankelevitz DF, et al. Early Lung Cancer Action Project: overall design and findings from baseline screening. *Lancet* 1999;354:99-105. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/10408484>.

1097. Ulaner GA, Lyall A. Identifying and distinguishing treatment effects and complications from malignancy at FDG PET/CT. *Radiographics* 2013;33:1817-1834. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24108564>.

1098. Shi Q, Smith TG, Michonski JD, et al. Symptom burden in cancer survivors 1 year after diagnosis: a report from the American Cancer Society's Studies of Cancer Survivors. *Cancer* 2011;117:2779-2790. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21495026>.

1099. Gelb AF, Tashkin DP, Epstein JD, et al. Physiologic characteristics of malignant unilateral main-stem bronchial obstruction. Diagnosis and Nd-YAG laser treatment. *Am Rev Respir Dis* 1988;138:1382-1385. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/2462389>.

1100. Howell DD, James JL, Hartsell WF, et al. Single-fraction radiotherapy versus multifraction radiotherapy for palliation of painful vertebral bone metastases-equivalent efficacy, less toxicity, more convenient: a subset analysis of Radiation Therapy Oncology Group trial 97-14. *Cancer* 2013;119:888-896. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23165743>.

1101. Griffioen GH, Toguri D, Dahele M, et al. Radical treatment of synchronous oligometastatic non-small cell lung carcinoma (NSCLC): patient outcomes and prognostic factors. *Lung Cancer* 2013;82:95-102. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23973202>.

1102. Ashworth A, Rodrigues G, Boldt G, Palma D. Is there an oligometastatic state in non-small cell lung cancer? A systematic review of the literature. *Lung Cancer* 2013;82:197-203. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24051084>.

1103. Collen C, Christian N, Schallier D, et al. Phase II study of stereotactic body radiotherapy to primary tumor and metastatic locations in

oligometastatic nonsmall-cell lung cancer patients. *Ann Oncol* 2014;25:1954-1959. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25114022>.

1104. Tree AC, Khoo VS, Eeles RA, et al. Stereotactic body radiotherapy for oligometastases. *Lancet Oncol* 2013;14:e28-37. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23276369>.

1105. Kelly P, Balter PA, Rebueno N, et al. Stereotactic body radiation therapy for patients with lung cancer previously treated with thoracic radiation. *Int J Radiat Oncol Biol Phys* 2010;78:1387-1393. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20381271>.

1106. Meijneke TR, Petit SF, Wentzler D, et al. Reirradiation and stereotactic radiotherapy for tumors in the lung: dose summation and toxicity. *Radiother Oncol* 2013;107:423-427. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23647748>.

1107. Peulen H, Karlsson K, Lindberg K, et al. Toxicity after reirradiation of pulmonary tumours with stereotactic body radiotherapy. *Radiother Oncol* 2011;101:260-266. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22056534>.

1108. Reyngold M, Wu AJ, McLane A, et al. Toxicity and outcomes of thoracic re-irradiation using stereotactic body radiation therapy (SBRT). *Radiat Oncol* 2013;8:99. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23617949>.

1109. Scagliotti GV, Hirsh V, Siena S, et al. Overall survival improvement in patients with lung cancer and bone metastases treated with denosumab versus zoledronic acid: subgroup analysis from a randomized phase 3 study. *J Thorac Oncol* 2012;7:1823-1829. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23154554>.

1110. Casas A, Llombart A, Martin M. Denosumab for the treatment of bone metastases in advanced breast cancer. *Breast* 2013;22:585-592. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23759273>.



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

1111. Ibrahim A, Scher N, Williams G, et al. Approval summary for zoledronic acid for treatment of multiple myeloma and cancer bone metastases. Clin Cancer Res 2003;9:2394-2399. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/12855610>.

1112. Henry D, Vadhan-Raj S, Hirsh V, et al. Delaying skeletal-related events in a randomized phase 3 study of denosumab versus zoledronic acid in patients with advanced cancer: an analysis of data from patients with solid tumors. Support Care Cancer 2014;22:679-687. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24162260>.

1113. Henry DH, Costa L, Goldwasser F, et al. Randomized, double-blind study of denosumab versus zoledronic acid in the treatment of bone metastases in patients with advanced cancer (excluding breast and prostate cancer) or multiple myeloma. J Clin Oncol 2011;29:1125-1132. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21343556>.

1114. Rosen LS, Gordon D, Tchekmedyan NS, et al. Long-term efficacy and safety of zoledronic acid in the treatment of skeletal metastases in patients with nonsmall cell lung carcinoma and other solid tumors: a randomized, Phase III, double-blind, placebo-controlled trial. Cancer 2004;100:2613-2621. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/15197804>.

1115. Henry DH, von Moos R, Hungria V, et al. Delaying skeletal-related events in a randomized phase III study of denosumab versus zoledronic acid in patients with advanced cancer [abstract]. J Clin Oncol 2010;28 (Suppl 15):Abstract 9133. Available at: http://meeting.ascopubs.org/cgi/content/abstract/28/15_suppl/9133.

1116. Sakuma Y, Matsukuma S, Yoshihara M, et al. Distinctive evaluation of nonmucinous and mucinous subtypes of bronchioloalveolar carcinomas in EGFR and K-ras gene-mutation analyses for Japanese lung adenocarcinomas: confirmation of the correlations with histologic subtypes and gene mutations. Am J Clin Pathol 2007;128:100-108. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17580276>.

1117. Shaw AT, Yeap BY, Solomon BJ, et al. Effect of crizotinib on overall survival in patients with advanced non-small-cell lung cancer harbouring

ALK gene rearrangement: a retrospective analysis. Lancet Oncol 2011;12:1004-1012. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21933749>.

1118. Pilkington G, Boland A, Brown T, et al. A systematic review of the clinical effectiveness of first-line chemotherapy for adult patients with locally advanced or metastatic non-small cell lung cancer. Thorax 2015;70:359-367. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25661113>.

1119. Sandler A, Yi J, Dahlberg S, et al. Treatment outcomes by tumor histology in Eastern Cooperative Group Study E4599 of bevacizumab with paclitaxel/carboplatin for advanced non-small cell lung cancer. J Thorac Oncol 2010;5:1416-1423. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20686429>.

1120. Socinski MA, Langer CJ, Huang JE, et al. Safety of bevacizumab in patients with non-small-cell lung cancer and brain metastases. J Clin Oncol 2009;27:5255-5261. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19738122>.

1121. Quoix E, Zalcman G, Oster JP, et al. Carboplatin and weekly paclitaxel doublet chemotherapy compared with monotherapy in elderly patients with advanced non-small-cell lung cancer: IFCT-0501 randomised, phase 3 trial. Lancet 2011;378:1079-1088. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21831418>.

1122. Santos FN, de Castria TB, Cruz MR, Riera R. Chemotherapy for advanced non-small cell lung cancer in the elderly population. Cochrane Database Syst Rev 2015;10:CD010463. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26482542>.

1123. Smit E, Moro-Sibilot D, Carpeno Jde C, et al. Cisplatin and carboplatin-based chemotherapy in the first-line treatment of non-small cell lung cancer: Analysis from the European FRAME study. Lung Cancer 2016;92:35-40. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26775594>.



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

1124. Kubota K, Kawahara M, Ogawara M, et al. Vinorelbine plus gemcitabine followed by docetaxel versus carboplatin plus paclitaxel in patients with advanced non-small-cell lung cancer: a randomised, open-label, phase III study. *Lancet Oncol* 2008;9:1135-1142. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19013107>.

1125. Johnson DH, Fehrenbacher L, Novotny WF, et al. Randomized phase II trial comparing bevacizumab plus carboplatin and paclitaxel with carboplatin and paclitaxel alone in previously untreated locally advanced or metastatic non-small-cell lung cancer. *J Clin Oncol* 2004;22:2184-2191. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/15169807>.

1126. Reck M, von Pawel J, Zatloukal P, et al. Phase III trial of cisplatin plus gemcitabine with either placebo or bevacizumab as first-line therapy for nonsquamous non-small-cell lung cancer: AVAIL. *J Clin Oncol* 2009;27:1227-1234. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19188680>.

1127. Mezger J, von Pawel J, Reck M. Bevacizumab (Bv) single-agent maintenance following Bv-based chemotherapy in patients with advanced non-small cell lung cancer (NSCLC): Results from an exploratory analysis of the AVAIL study [abstract]. *J Clin Oncol* 2009;27 (Suppl 15):Abstract e19001. Available at: <http://meeting.ascopubs.org/cgi/content/abstract/27/15S/e19001>.

1128. Scagliotti G, Brodowicz T, Shepherd FA, et al. Treatment-by-histology interaction analyses in three phase III trials show superiority of pemetrexed in nonsquamous non-small cell lung cancer. *J Thorac Oncol* 2011;6:64-70. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21119545>.

1129. Soon YY, Stockler MR, Askie LM, Boyer MJ. Duration of chemotherapy for advanced non-small-cell lung cancer: a systematic review and meta-analysis of randomized trials. *J Clin Oncol* 2009;27:3277-3283. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19470938>.

1130. Eisenhauer EA, Therasse P, Bogaerts J, et al. New response evaluation criteria in solid tumours: revised RECIST guideline (version

1.1). *Eur J Cancer* 2009;45:228-247. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19097774>.

1131. Therasse P, Arbuck SG, Eisenhauer EA, et al. New guidelines to evaluate the response to treatment in solid tumors. European Organization for Research and Treatment of Cancer, National Cancer Institute of the United States, National Cancer Institute of Canada. *J Natl Cancer Inst* 2000;92:205-216. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/10655437>.

1132. Nishino M, Hatabu H, Johnson BE, McCloud TC. State of the art: Response assessment in lung cancer in the era of genomic medicine. *Radiology* 2014;271:6-27. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24661292>.

1133. Pfister DG, Johnson DH, Azzoli CG, et al. American Society of Clinical Oncology treatment of unresectable non-small-cell lung cancer guideline: update 2003. *J Clin Oncol* 2004;22:330-353. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/14691125>.

1134. Park K, Yu CJ, Kim SW, et al. First-Line Erlotinib Therapy Until and Beyond Response Evaluation Criteria in Solid Tumors Progression in Asian Patients With Epidermal Growth Factor Receptor Mutation-Positive Non-Small-Cell Lung Cancer: The ASPIRATION Study. *JAMA Oncol* 2016;2:305-312. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26720423>.

1135. Becker K, Xu Y. Management of tyrosine kinase inhibitor resistance in lung cancer with EGFR mutation. *World J Clin Oncol* 2014;5:560-567. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25302160>.

1136. Choudhury NJ, Marra A, Sui JSY, et al. Molecular biomarkers of disease outcomes and mechanisms of acquired resistance to first-line osimertinib in advanced EGFR-mutant lung cancers. *J Thorac Oncol* 2022. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/36494075>.

1137. Ou SH. Second-generation irreversible epidermal growth factor receptor (EGFR) tyrosine kinase inhibitors (TKIs): a better mousetrap? A



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

review of the clinical evidence. Crit Rev Oncol Hematol 2012;83:407-421. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22257651>.

1138. Nguyen KS, Kobayashi S, Costa DB. Acquired resistance to epidermal growth factor receptor tyrosine kinase inhibitors in non-small-cell lung cancers dependent on the epidermal growth factor receptor pathway. Clin Lung Cancer 2009;10:281-289. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19632948>.

1139. Gazdar AF. Activating and resistance mutations of EGFR in non-small-cell lung cancer: role in clinical response to EGFR tyrosine kinase inhibitors. Oncogene 2009;28 Suppl 1:S24-31. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19680293>.

1140. Leonetti A, Sharma S, Minari R, et al. Resistance mechanisms to osimertinib in EGFR-mutated non-small cell lung cancer. Br J Cancer 2019;121:725-737. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31564718>.

1141. Cheema PK, Banerji SO, Blais N, et al. Canadian Consensus Recommendations on the management of MET-altered NSCLC. Curr Oncol 2021;28:4552-4576. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/34898564>.

1142. Conforti F, Catania C, Toffalorio F, et al. EGFR tyrosine kinase inhibitors beyond focal progression obtain a prolonged disease control in patients with advanced adenocarcinoma of the lung. Lung Cancer 2013;81:440-444. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23810573>.

1143. Miller VA, Hirsh V, Cadranell J, et al. Afatinib versus placebo for patients with advanced, metastatic non-small-cell lung cancer after failure of erlotinib, gefitinib, or both, and one or two lines of chemotherapy (LUX-Lung 1): a phase 2b/3 randomised trial. Lancet Oncol 2012;13:528-538. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22452896>.

1144. Katakami N, Atagi S, Goto K, et al. LUX-Lung 4: a phase II trial of afatinib in patients with advanced non-small-cell lung cancer who

progressed during prior treatment with erlotinib, gefitinib, or both. J Clin Oncol 2013;31:3335-3341. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23816963>.

1145. Hirsh V, Cadranell J, Cong XJ, et al. Symptom and quality of life benefit of afatinib in advanced non-small-cell lung cancer patients previously treated with erlotinib or gefitinib: results of a randomized phase IIb/III trial (LUX-Lung 1). J Thorac Oncol 2013;8:229-237. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23328549>.

1146. Meoni G, Cecere FL, Lucherini E, Di Costanzo F. Medical treatment of advanced non-small cell lung cancer in elderly patients: a review of the role of chemotherapy and targeted agents. J Geriatr Oncol 2013;4:282-290. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24070465>.

1147. Weiss JM, Stinchcombe TE. Second-Line Therapy for Advanced NSCLC. Oncologist 2013;18:947-953. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23918070>.

1148. van Putten JW, Baas P, Codrington H, et al. Activity of single-agent gemcitabine as second-line treatment after previous chemotherapy or radiotherapy in advanced non-small-cell lung cancer. Lung Cancer 2001;33:289-298. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/11551424>.

1149. Crino L, Mosconi AM, Scagliotti G, et al. Gemcitabine as second-line treatment for advanced non-small-cell lung cancer: A phase II trial. J Clin Oncol 1999;17:2081-2085. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/10561261>.

1150. Anderson H, Hopwood P, Stephens RJ, et al. Gemcitabine plus best supportive care (BSC) vs BSC in inoperable non-small cell lung cancer--a randomized trial with quality of life as the primary outcome. UK NSCLC Gemcitabine Group. Non-Small Cell Lung Cancer. Br J Cancer 2000;83:447-453. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/10945489>.



NCCN Guidelines Version 3.2024

Non-Small Cell Lung Cancer

1151. Sculier JP, Lafitte JJ, Berghmans T, et al. A phase II trial testing gemcitabine as second-line chemotherapy for non small cell lung cancer. The European Lung Cancer Working Party. 101473.1044@compuserve.com. Lung Cancer 2000;29:67-73. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/10880849>.

1152. Fossella FV, DeVore R, Kerr RN, et al. Randomized phase III trial of docetaxel versus vinorelbine or ifosfamide in patients with advanced non-small-cell lung cancer previously treated with platinum-containing chemotherapy regimens. The TAX 320 Non-Small Cell Lung Cancer Study Group. J Clin Oncol 2000;18:2354-2362. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/10856094>.

1153. Shepherd FA, Dancey J, Ramlau R, et al. Prospective randomized trial of docetaxel versus best supportive care in patients with non-small-cell lung cancer previously treated with platinum-based chemotherapy. J Clin Oncol 2000;18:2095-2103. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/10811675>.

1154. Hanna N, Shepherd FA, Fossella FV, et al. Randomized phase III trial of pemetrexed versus docetaxel in patients with non-small-cell lung cancer previously treated with chemotherapy. J Clin Oncol 2004;22:1589-1597. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/15117980>.

1155. Shepherd FA, Rodrigues Pereira J, Ciuleanu T, et al. Erlotinib in previously treated non-small-cell lung cancer. N Engl J Med 2005;353:123-132. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/16014882>.

1156. Schwartz LH, Litiere S, de Vries E, et al. RECIST 1.1-Update and clarification: From the RECIST committee. Eur J Cancer 2016;62:132-137. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27189322>.

1157. Ades F, Yamaguchi N. WHO, RECIST, and immune-related response criteria: is it time to revisit pembrolizumab results? Ecanermediscience 2015;9:604. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26715941>.

1158. Wolchok JD, Hoos A, O'Day S, et al. Guidelines for the evaluation of immune therapy activity in solid tumors: immune-related response criteria. Clin Cancer Res 2009;15:7412-7420. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19934295>.

1159. Cavanna L, Citterio C, Orlandi E. Immune checkpoint inhibitors in EGFR-mutation positive TKI-treated patients with advanced non-small-cell lung cancer network meta-analysis. Oncotarget 2019;10:209-215. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30719215>.

1160. Mok TSK, Kim SW, Wu YL, et al. Gefitinib Plus Chemotherapy Versus Chemotherapy in Epidermal Growth Factor Receptor Mutation-Positive Non-Small-Cell Lung Cancer Resistant to First-Line Gefitinib (IMPRESS): Overall Survival and Biomarker Analyses. J Clin Oncol 2017;35:4027-4034. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28968167>.

1161. Sacher AG, Janne PA, Oxnard GR. Management of acquired resistance to epidermal growth factor receptor kinase inhibitors in patients with advanced non-small cell lung cancer. Cancer 2014;120:2289-2298. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24752335>.

1162. Demarinis F, Paul S, Hanna N, et al. Survival update for the phase III study of pemetrexed vs docetaxel in non-small cell lung cancer (NSCLC) [abstract]. J Clin Oncol 2006;24 (Suppl 18):Abstract 7133. Available at: http://meeting.ascopubs.org/cgi/content/abstract/24/18_suppl/7133.

1163. Garassino MC, Martelli O, Brogini M, et al. Erlotinib versus docetaxel as second-line treatment of patients with advanced non-small-cell lung cancer and wild-type EGFR tumours (TAILOR): a randomised controlled trial. Lancet Oncol 2013;14:981-988. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23883922>.

1164. Kawaguchi T, Ando M, Asami K, et al. Randomized phase III trial of erlotinib versus docetaxel as second- or third-line therapy in patients with advanced non-small-cell lung cancer: Docetaxel and Erlotinib Lung Cancer Trial (DELTA). J Clin Oncol 2014;32:1902-1908. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24841974>.



NCCN Guidelines Version 3.2024 Non-Small Cell Lung Cancer

1165. Eccles BK, Geldart TR, Laurence VM, et al. Experience of first- and subsequent-line systemic therapy in the treatment of non-small cell lung cancer. *Ther Adv Med Oncol* 2011;3:163-170. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21904578>.

1166. Morgensztern D, Cobo M, Ponce Aix S, et al. ABOUND.2L+: A randomized phase 2 study of nanoparticle albumin-bound paclitaxel with or without CC-486 as second-line treatment for advanced nonsquamous non-small cell lung cancer (NSCLC). *Cancer* 2018;124:4667-4675. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30383906>.

1167. Langer CJ, Mok T, Postmus PE. Targeted agents in the third-/fourth-line treatment of patients with advanced (stage III/IV) non-small cell lung cancer (NSCLC). *Cancer Treat Rev* 2013;39:252-260. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22703830>.

1168. Noble J, Ellis PM, Mackay JA, et al. Second-line or subsequent systemic therapy for recurrent or progressive non-small cell lung cancer: a systematic review and practice guideline. *J Thorac Oncol* 2006;1:1042-1058. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17409993>.